



on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

```
=> s coenzyme q10/cn
L5          1 COENZYME Q10/CN
```

```
=> file caplus
COST IN U.S. DOLLARS
```

SINCE FILE	TOTAL
ENTRY	SESSION
5.99	20.13

```
FULL ESTIMATED COST
```

FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Apr 2010 VOL 152 ISS 17  
FILE LAST UPDATED: 21 Apr 2010 (20100421/ED)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2010  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2010

Caplus now includes complete International Patent Classification (IPC) reclassification data for the first quarter of 2010.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

```
=> s l5 and carcinoma
      5610 L5
      222424 CARCINOMA
      39195 CARCINOMAS
      179 CARCINOMATA
      1 CARCINOMATAS
      231654 CARCINOMA
          (CARCINOMA OR CARCINOMAS OR CARCINOMATA OR CARCINOMATAS)
L6          41 L5 AND CARCINOMA
```

```
=> d ti total
```

```
L6  ANSWER 1 OF 41  CAPLUS  COPYRIGHT 2010 ACS on STN
TI  Drug Effects Viewed from a Signal Transduction Network Perspective

L6  ANSWER 2 OF 41  CAPLUS  COPYRIGHT 2010 ACS on STN
TI  Application of nanoscale polysaccharide of Ganoderma as antitumor agent
```

L6 ANSWER 3 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Effect of Coenzyme Q10, Riboflavin and Niacin on Tamoxifen treated postmenopausal breast cancer women with special reference to blood chemistry profiles

L6 ANSWER 4 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Molecular modeling and experimental evidence for hypericin as a substrate for mitochondrial complex III; mitochondrial photodamage as demonstrated using specific inhibitors

L6 ANSWER 5 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Topical formulations comprising lipophilic bioactive agents having enhanced bioavailability

L6 ANSWER 6 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Method of cancer screening; method of cancer treatment; and method of auto-immune disease treatment

L6 ANSWER 7 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Composition, its use for treating systemic diseases a conditions, and product containing said composition

L6 ANSWER 8 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Foamable vehicle and vitamin and flavonoid pharmaceutical compositions thereof for treatment of skin and other disorders

L6 ANSWER 9 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Protective effect of coenzyme Q10 against cisplatin-induced nephrotoxicity in rats

L6 ANSWER 10 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Fused heterocyclic-substituted dihydroxyheptenoic acid as HMG-CoA reductase inhibitors and their preparation, pharmaceutical compositions and use in the treatment of diseases

L6 ANSWER 11 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer

L6 ANSWER 12 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Transcatheter tumor immunoembolization

L6 ANSWER 13 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Ameliorating effect of coenzyme Q10, riboflavin and niacin in tamoxifen-treated postmenopausal breast cancer patients with special reference to lipids and lipoproteins

L6 ANSWER 14 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Topically applied glucosamine sulfate and all its related, precursor, and derivative compounds significantly increases the skin's natural production of hyaluronic acid for the rejuvenation of healthier younger-looking skin; while phosphatidylcholine is required to replace its deficiency caused by topical dimethylaminoethanol (DMAE)

L6 ANSWER 15 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Composition for moderating alcohol metabolism and for reducing the risk of alcohol induced diseases

L6 ANSWER 16 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Determination of coenzyme Q10 in functional and neoplastic human renal

tissues

- L6 ANSWER 17 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Marker genes for the diagnosis of chronic fatigue syndrome by gene expression profiling
- L6 ANSWER 18 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI NAD<sup>+</sup>/NADH and/or CoQ/CoQH<sub>2</sub> ratios from plasma membrane electron transport may determine ceramide and sphingosine-1-phosphate levels accompanying G1 arrest and apoptosis
- L6 ANSWER 19 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Chemoprevention of breast cancer: current status and future prospects
- L6 ANSWER 20 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Methods and compositions for the treatment of diseases characterized by calcification and/or plaque formation
- L6 ANSWER 21 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Energy-modulating vitamins - a new combinatorial therapy prevents cancer cachexia in rat mammary carcinoma
- L6 ANSWER 22 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Combined efficacy of tamoxifen and coenzyme Q10 on the status of lipid peroxidation and antioxidants in DMBA induced breast cancer
- L6 ANSWER 23 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Effect of a nutritional supplement containing vitamin E, selenium, vitamin C and coenzyme Q10 on serum PSA in patients with hormonally untreated carcinoma of the prostate: a randomized placebo-controlled study
- L6 ANSWER 24 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Augmented efficacy of tamoxifen in rat breast tumorigenesis when gavaged along with riboflavin, niacin, and CoQ10: Effects on lipid peroxidation and antioxidants in mitochondria
- L6 ANSWER 25 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Role of mitochondria in neuronal cell death induced by oxidative stress; neuroprotection by Coenzyme Q10
- L6 ANSWER 26 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Tetracycline compounds having target therapeutic activities
- L6 ANSWER 27 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Systemic treatment of pathological conditions resulting from oxidative stress and/or redox imbalance
- L6 ANSWER 28 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Tetracycline compounds having target therapeutic activities
- L6 ANSWER 29 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Coenzyme Q10 concentrations and antioxidant status in tissues of breast cancer patients
- L6 ANSWER 30 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Effect of radiation therapy on small-cell lung cancer is reduced by ubiquinone intake
- L6 ANSWER 31 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
TI Plasma coenzyme Q10 concentrations in breast cancer. Prognosis and therapeutic consequences

L6 ANSWER 32 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Coenzyme Q10 tissular levels in colonic and gastric carcinomas

L6 ANSWER 33 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Progress on therapy of breast cancer with vitamin Q10 and the regression of metastases

L6 ANSWER 34 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Method of inhibiting carcinogenesis by treatment with dehydroepiandrosterone and analogs thereof

L6 ANSWER 35 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Vitamin E and coenzyme Q10 in normal human skin and in basal cell epitheliomas

L6 ANSWER 36 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Studies on the polyisoprenoid composition in hepatocellular carcinomas and its correlation with their differentiation

L6 ANSWER 37 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Effects of vitamins on lipid peroxidation and suppression of DNA synthesis induced by adriamycin in Ehrlich cells

L6 ANSWER 38 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Effect of BCG, coenzyme Q10, or their combination on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats

L6 ANSWER 39 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Combined effect of BCG and coenzyme Q10 on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats

L6 ANSWER 40 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Biosynthetic changes of vitamin K3 and ubiquinone 0 in man

L6 ANSWER 41 OF 41 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Histochemical studies of the effects of coenzyme Q10 and menadione on oxidative enzymes in normal and neoplastic cells

=> s l6 and py<=2004  
 25157969 PY<=2004  
 L7 16 L6 AND PY<=2004

=> d ti total

L7 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Tetracycline compounds having target therapeutic activities

L7 ANSWER 2 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Systemic treatment of pathological conditions resulting from oxidative stress and/or redox imbalance

L7 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Tetracycline compounds having target therapeutic activities

L7 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Coenzyme Q10 concentrations and antioxidant status in tissues of breast cancer patients

L7 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

TI Effect of radiation therapy on small-cell lung cancer is reduced by ubiquinone intake  
 L7 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Plasma coenzyme Q10 concentrations in breast cancer. Prognosis and therapeutic consequences  
 L7 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Coenzyme Q10 tissular levels in colonic and gastric carcinomas  
 L7 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Progress on therapy of breast cancer with vitamin Q10 and the regression of metastases  
 L7 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Method of inhibiting carcinogenesis by treatment with dehydroepiandrosterone and analogs thereof  
 L7 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Vitamin E and coenzyme Q10 in normal human skin and in basal cell epitheliomas  
 L7 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Studies on the polyisoprenoid composition in hepatocellular carcinomas and its correlation with their differentiation  
 L7 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Effects of vitamins on lipid peroxidation and suppression of DNA synthesis induced by adriamycin in Ehrlich cells  
 L7 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Effect of BCG, coenzyme Q10, or their combination on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats  
 L7 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Combined effect of BCG and coenzyme Q10 on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats  
 L7 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Biosynthetic changes of vitamin K3 and ubiquinone 0 in man  
 L7 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 TI Histochemical studies of the effects of coenzyme Q10 and menadione on oxidative enzymes in normal and neoplastic cells

=> d ibib abs 17 total

L7 ANSWER 1 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2004:633439 CAPLUS  
 DOCUMENT NUMBER: 141:167771  
 TITLE: Tetracycline compounds having target therapeutic activities  
 INVENTOR(S): Levy, Stuart B.; Draper, Michael; Nelson, Mark L.; Jones, Graham  
 PATENT ASSIGNEE(S): Paratek Pharmaceuticals, Inc., USA  
 SOURCE: PCT Int. Appl., 277 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004064728	A2	20040805	WO 2004-US1036	20040116 <--
WO 2004064728	A3	20041216		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI US 20060194773 A1 20060831 US 2004-996119 20041122				
PRIORITY APPLN. INFO.:				
US 2003-441141P P 20030116				
US 2001-305546P P 20010713				
US 2002-395741P P 20020712				
US 2002-196010 A2 20020715				
US 2004-759484 B1 20040116				

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 141:167771

AB Methods and compds. for treating diseases, e.g. inflammation process-associated states, with tetracycline compds. having a target therapeutic activity are described. Preparation of selected tetracycline compds. is described.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 2 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:1007587 CAPLUS

DOCUMENT NUMBER: 140:35996

TITLE: Systemic treatment of pathological conditions resulting from oxidative stress and/or redox imbalance

INVENTOR(S): Gojon-Romanillos, Gabriel

PATENT ASSIGNEE(S): Mex.

SOURCE: U.S. Pat. Appl. Publ., 14 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030235571	A1	20031225	US 2003-463765	20030618 <--
US 20090181081	A1	20090716	US 2009-405165	20090316
US 20090304819	A1	20091210	US 2009-543407	20090818
PRIORITY APPLN. INFO.:				
US 2002-389491P P 20020619				
US 2003-463765 B3 20030618				
US 2009-405165 A2 20090316				
AB Alterations of redox homeostasis in mammals underlie a host of symptoms, syndromes and diseases, including AIDS and cancer, which can be successfully treated by administration to a mammal of therapeutically-effective amts. of sulfide compds. and/or thiosulfate compds. and/or thionite compds. and/or sulfite compds. and/or thionate compds. and/or any organic, inorg. or organometallic precursors thereof. The unique compds. of this invention contain one or more "active sulfur compds." in combination with each other or with other therapeutic agents. The invention also encompasses the varying modes of administration of the therapeutic compds.				

L7 ANSWER 3 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:57866 CAPLUS  
 DOCUMENT NUMBER: 138:117673  
 TITLE: Tetracycline compounds having target therapeutic activities  
 INVENTOR(S): Levy, Stuart B.; Draper, Michael; Nelson, Mark L.; Jones, Graham  
 PATENT ASSIGNEE(S): Paratek Pharmaceuticals, Inc., USA  
 SOURCE: PCT Int. Appl., 158 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003005971	A2	20030123	WO 2002-US22451	20020715 <--
WO 2003005971	A3	20031127		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002318238	A1	20030129	AU 2002-318238	20020715 <--
US 20040063674	A1	20040401	US 2002-196010	20020715 <--
EP 1408987	A2	20040421	EP 2002-748169	20020715 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2004537544	T	20041216	JP 2003-511780	20020715 <--
US 20060194773	A1	20060831	US 2004-996119	20041122
JP 2009298801	A	20091224	JP 2009-187938	20090814
PRIORITY APPLN. INFO.:				
US 2001-305546P P 20010713 US 2002-395741P P 20020712 JP 2003-511780 A3 20020715 US 2002-196010 A2 20020715 WO 2002-US22451 W 20020715 US 2003-441141P P 20030116 US 2004-759484 B1 20040116				

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 138:117673  
 AB Methods and compds. for treating a variety of diseases with tetracycline compds. having a target therapeutic activity are described, as is compound preparation  
 OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)  
 REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 4 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2000:683633 CAPLUS  
 DOCUMENT NUMBER: 134:129378  
 TITLE: Coenzyme Q10 concentrations and antioxidant status in tissues of breast cancer patients  
 AUTHOR(S): Portakal, Oytun; Ozkaya, Ozay; Inal, Mine Erden; Bozan, Berrin; Kosan, Muberra; Sayek, Iskender  
 CORPORATE SOURCE: Department of Biochemistry, The Medical School of



SOURCE: Osmangazi University, Eskisehir, Turk.  
Clinical Biochemistry (2000), 33(4), 279-284  
CODEN: CLBIAS; ISSN: 0009-9120

PUBLISHER: Elsevier Science Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB An increasing amount of exptl. and epidemiol. evidence implicates the involvement of oxygen derived radicals in the pathogenesis of cancer development. Oxygen derived radicals are able to cause damage to membranes, mitochondria, and macromols. including proteins, lipids and DNA. Accumulation of DNA damages has been suggested to contribute to carcinogenesis. It would, therefore, be advantageous to pinpoint the effects of oxygen derived radicals in cancer development. In the present study, we investigated the relationship between oxidative stress and breast cancer development in tissue level. Breast cancer is the most common malignant disease in Western women. Twenty-one breast cancer patients, who underwent radical mastectomy and diagnosed with infiltrative ductal carcinoma, were used in the study. We determined coenzyme Q10 (Q) concns., antioxidant enzyme activities (mitochondrial and total superoxide dismutase (SOD), glutathione peroxidase (GSH-Px), catalase), and malondialdehyde (MDA) levels in tumor and surrounding tumor-free tissues. Q concns. in tumor tissues significantly decreased as compared to the surrounding normal tissues ( $p < 0.001$ ). Higher MDA levels were observed in tumor tissues than noncancerous tissues ( $p < 0.001$ ). The activities of MnSOD, total SOD, GSH-Px and catalase in tumor tissues significantly increased ( $p < 0.001$ ) compared to the controls. These findings may support that reactive oxygen species increased in malignant cells, and may cause overexpression of antioxidant enzymes and the consumption of coenzyme Q10. Increased antioxidant enzyme activities may be related with the susceptibility of cells to carcinogenic agents and the response of tumor cells to the chemotherapeutic agents. Administration of coenzyme Q10 by nutrition may induce the protective effect of coenzyme Q10 on breast tissue.

OS.CITING REF COUNT: 55 THERE ARE 55 CAPLUS RECORDS THAT CITE THIS RECORD (55 CITINGS)

REFERENCE COUNT: 40 THERE ARE 40 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 5 OF 16 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 1998:594275 CAPLUS

DOCUMENT NUMBER: 129:287378

ORIGINAL REFERENCE NO.: 129:58489a,58492a

TITLE: Effect of radiation therapy on small-cell lung cancer is reduced by ubiquinone intake

AUTHOR(S): Lund, E. L.; Quistorff, B.; Spang-Thomsen, M.; Kristjansen, P. E. G.

CORPORATE SOURCE: Institute of Molecular Pathology and NMR-Center, University of Copenhagen, 2100, Den.

SOURCE: Folia Microbiologica (Prague) (1998), 43(5), 505-506  
CODEN: FOMIAZ; ISSN: 0015-5632

PUBLISHER: Institute of Microbiology, Academy of Sciences of the Czech Republic

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effect of oral ubiquinone (Q10) intake on the in vivo response of tumors to single dose radiotherapy was examined. The human small-cell lung cancer (SCLC) line CPH 054A, which is sensitive to relatively low doses of X-radiation, was grown as s.c. transplants in the flanks of nude nu/nu mice. When macroscopical growth was established, groups of mice received either 10, 20 or 40 mg/kg Q10 in 30 mL soy oil intragastrically daily on 4

consecutive days. Controls received either 30 mL of pure soy oil or nothing. Three h after the last dose half of the tumors in each group received a single radiation dose of 5 Gy, using a 300 kV therapeutic unit. The macroscopic growth pre- and posttreatment was analyzed according to a transformed Gompertz algorithm using the software program GROWTH. Treatment with Q10 or soy oil alone had no effect on tumor growth compared with untreated controls. Groups of tumors that received Q10 and radiotherapy had a significantly lower specific growth delay (SGD) than the radiotherapy-only groups. This effect was significant at 40 mg/kg and borderline at 20 mg/kg, whereas at 10 mg/kg no radioprotection was seen. We conclude that systemic Q10 reduces the response to single dose tumor irradiation in xenotransplanted human SCLC tumors.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)  
REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 6 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1998:591187 CAPLUS  
DOCUMENT NUMBER: 129:215028  
ORIGINAL REFERENCE NO.: 129:43683a,43686a  
TITLE: Plasma coenzyme Q10 concentrations in breast cancer.  
Prognosis and therapeutic consequences  
AUTHOR(S): Jolliet, P.; Simon, N.; Barre, J.; Pons, J.-Y.;  
Boukef, M.; Paniel, B.-J.; Tillement, J.-P.  
CORPORATE SOURCE: Service Hospitalo-Universitaire Pharmacologie,  
Creteil, F-94010, Fr.  
SOURCE: International Journal of Clinical Pharmacology and  
Therapeutics (1998), 36(9), 506-509  
CODEN: ICTHEK; ISSN: 0946-1965  
PUBLISHER: Dustri-Verlag Dr. Karl Feistle  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB To understand the role of coenzyme Q10 or ubiquinone in the pathogenesis of breast cancer, a clin. trial was conducted, including women hospitalized for the biopsy and/or the ablation of a breast tumor. Ubiquinone blood plasma concns. were determined simultaneously with vitamin E blood plasma concns. by HPLC. A coenzyme Q10 deficiency was noted both in the carcinoma (80 patients) and non-malignant lesions (120 patients), while vitamin E concns. were within the normal range. A correlation was shown between the intensity of the deficiency and the bad prognosis of the breast disease based on high TNM and SBR values or the lack of estrogen receptors. It is concluded that ubiquinone supplementation could be relevant in breast cancer.

OS.CITING REF COUNT: 26 THERE ARE 26 CAPLUS RECORDS THAT CITE THIS  
RECORD (26 CITINGS)

L7 ANSWER 7 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1997:60453 CAPLUS  
DOCUMENT NUMBER: 126:87756  
ORIGINAL REFERENCE NO.: 126:16925a,16928a  
TITLE: Coenzyme Q10 tissular levels in colonic and gastric carcinomas  
AUTHOR(S): Romagnoli, A.; Oradei, A.; Destito, C.; Marin, A.  
Wiel; Littarru, G. P.  
CORPORATE SOURCE: Istituto di Clinica chirurgica generale e Terapia  
chirurgica, Universita Cattolica del Sacro Cuore,  
Rome, Italy  
SOURCE: Acta Medica Romana (1994), 32(4), 561-565  
CODEN: AMROBA; ISSN: 0001-6098  
PUBLISHER: Vita e Pensiero

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The authors evaluated Coenzyme Q10 levels in colorectal and gastric carcinomas in comparison with resp. normal mucosas in the same patients. In bowel carcinomas Coenzyme Q10 levels are statistically higher than in normal mucosa levels, whereas gastric carcinomas Coenzyme Q10 levels are similar to the levels seen in normal mucosa. Further studies are necessary to inquire the different antioxidant behavior of these neoplasms and to correlate Coenzyme Q10 tissue content to radiosensitivity of these neoplasms.

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L7 ANSWER 8 OF 16 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 1995:673575 CAPLUS

DOCUMENT NUMBER: 123:47542

ORIGINAL REFERENCE NO.: 123:8295a,8298a

TITLE: Progress on therapy of breast cancer with vitamin Q10 and the regression of metastases

AUTHOR(S): Lockwood, Knud; Moesgaard, Sven; Yamamoto, Tatsuo; Folkers, Karl

CORPORATE SOURCE: Malmoegeade 5, Copenhagen, Den.

SOURCE: Biochemical and Biophysical Research Communications (1995), 212(1), 172-7

CODEN: BBRCA9; ISSN: 0006-291X

PUBLISHER: Academic

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Over 35 yr, data and knowledge have internationally evolved from biochem., biomedical and clin. research on vitamin Q10 (coenzyme Q10; CoQ10) and cancer, which led in 1993 to overt complete regression of the tumors in two cases of breast cancer. Continuing this research, three addnl. breast cancer patients also underwent a conventional protocol of therapy which included a daily oral dosage of 390 mg of vitamin Q10 (Bio-Quinone of Pharma Nord) during the complete trials over 3-5 yr. The numerous metastases in the liver of a 44-yr-old patient "disappeared," and no signs of metastases were found elsewhere. A 49-yr-old patient, on a dosage of 390 mg of vitamin Q10, revealed no signs of tumor in the pleural cavity after six months, and her condition was excellent. A 75-yr-old patient with carcinoma in one breast, after lumpectomy and 390 mg of CoQ10, showed no cancer in the tumor bed or metastases. Control blood levels of CoQ10 of 0.83-0.97 and of 0.62 µg/mL increased to 3.34-3.64 and to 3.77 µg/mL, resp., on therapy with CoQ10 for patients A-MRH and EEL.

OS.CITING REF COUNT: 32 THERE ARE 32 CAPLUS RECORDS THAT CITE THIS RECORD (32 CITINGS)

L7 ANSWER 9 OF 16 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 1993:662505 CAPLUS

DOCUMENT NUMBER: 119:262505

ORIGINAL REFERENCE NO.: 119:46681a,46684a

TITLE: Method of inhibiting carcinogenesis by treatment with dehydroepiandrosterone and analogs thereof

INVENTOR(S): Nyce, Jonathan W.

PATENT ASSIGNEE(S): East Carolina University, USA

SOURCE: PCT Int. Appl., 34 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9316704	A1	19930902	WO 1993-US1637	19930223 <--
W: AU, CA, JP RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9337317	A	19930913	AU 1993-37317	19930223 <--
AU 676470	B2	19970313		
EP 627921	A1	19941214	EP 1993-906193	19930223 <--
EP 627921	B1	20000531		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
JP 07504198	T	19950511	JP 1993-515052	19930223 <--
AT 193447	T	20000615	AT 1993-906193	19930223 <--
ES 2146228	T3	20000801	ES 1993-906193	19930223 <--
CA 2117532	C	20010410	CA 1993-2117532	19930223 <--
US 5527789	A	19960618	US 1994-284307	19940802 <--
PRIORITY APPLN. INFO.:			US 1992-840510	A 19920224
			WO 1993-US1637	A 19930223
OTHER SOURCE(S): MARPAT 119:262505				
AB Dehydroepiandrosterone (DHEA) or a DHEA analog is used to combat cancer. An Ubiquinone is used to combat heart failure induced by the DHEA or analog. DHEA inhibited the growth of HT-29 SF cells; DHEA produced a G1 block in the cells in a time- and dose-dependent manner. Anal. of reversal of DHEA-mediated growth inhibition and reversal of DHEA-induced cell-cycle arrest is also described.				
OS.CITING REF COUNT:		8	THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)	
REFERENCE COUNT:		2	THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT	
L7 ANSWER 10 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN				
ACCESSION NUMBER:		1993:210313	CAPLUS	
DOCUMENT NUMBER:		118:210313		
ORIGINAL REFERENCE NO.:		118:36173a,36176a		
TITLE: Vitamin E and coenzyme Q10 in normal human skin and in basal cell epitheliomas				
AUTHOR(S): Rusciani, Luigi; Petrelli, Giuseppina; Lipa, Silvio				
CORPORATE SOURCE: Cathol. Univ. Sacred Heart, Rome, Italy				
SOURCE: Vitam. E Health Dis. (1993), 765-73.				
Editor(s): Packer, Lester; Fuchs, Juergen. Dekker: New York, N. Y.				
CODEN: 58VAAM				
DOCUMENT TYPE:		Conference; General Review		
LANGUAGE:		English		
AB A review with 31 refs.				
L7 ANSWER 11 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN				
ACCESSION NUMBER:		1991:426885	CAPLUS	
DOCUMENT NUMBER:		115:26885		
ORIGINAL REFERENCE NO.:		115:4701a,4704a		
TITLE: Studies on the polyisoprenoid composition in hepatocellular carcinomas and its correlation with their differentiation				
AUTHOR(S): Eggens, I.; Elmberger, P. G.				
CORPORATE SOURCE: Dep. Cell. Neuropathol., Huddinge Hosp., Huddinge, Swed.				
SOURCE: APMIS (1990), 98(6), 535-42				
CODEN: APMSEL; ISSN: 0903-4641				
DOCUMENT TYPE:		Journal		
LANGUAGE:		English		
AB The levels of cholesterol, ubiquinone and dolichol and the polyprenol				

composition of dolichol in human hepatocellular carcinomas (hepatomas) with different degrees of differentiation were analyzed and compared with healthy liver tissue. Dolichols were also analyzed in liver metastases. The total level of cholesterol was increased, while the levels of dolichol and ubiquinone were decreased in all hepatomas, but no correlation between these levels and the degree of differentiation of the hepatomas could be observed. The level of dolichol decreased more in the hepatomas than in the liver metastases. The dolichol fraction from hepatomas with a low degree of differentiation contained higher relative amts. of short polyisoprenols (D17) and slightly lower relative amts. of D21 compared with healthy liver tissue, metastatic liver tumors or hepatomas with a high degree of differentiation. The significance of the lipid values found in the different groups is discussed.

L7 ANSWER 12 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1985:453034 CAPLUS

DOCUMENT NUMBER: 103:53034

ORIGINAL REFERENCE NO.: 103:8535a,8538a

TITLE: Effects of vitamins on lipid peroxidation and suppression of DNA synthesis induced by adriamycin in Ehrlich cells

AUTHOR(S): Okamoto, Kouji; Ogura, Ryohei

CORPORATE SOURCE: Sch. Med., Kurume Univ., Kurume, 830, Japan

SOURCE: Journal of Nutritional Science and Vitaminology (

1985), 31(2), 129-37

CODEN: JNSVA5; ISSN: 0301-4800

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The effects of various vitamins on lipid peroxidn. and the suppression of DNA synthesis induced by adriamycin (ADR) [23214-92-8] in vitro using Ehrlich ascites carcinoma (EAC) cells were studied. ADR produced a concentration-dependent stimulation of lipid peroxidn. in EAC cells.  $\alpha$ -Tocopherol [59-02-9] and coenzyme Q10 [303-98-0] inhibited ADR-induced lipid peroxidn. to about the same extent and these effects were the greatest for all antioxidants added. The inhibitory effect of riboflavin 2',3',4',5'-tetrabutyrate [752-56-7] was greater than that of riboflavin 5'-phosphate [146-17-8]. On measuring incorporation of [3H]thymidine into EAC cells, these vitamins did not alter appreciably the magnitude of the ADR-induced suppression of DNA synthesis in EAC cells.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L7 ANSWER 13 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1979:604479 CAPLUS

DOCUMENT NUMBER: 91:204479

ORIGINAL REFERENCE NO.: 91:32811a,32814a

TITLE: Effect of BCG, coenzyme Q10, or their combination on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats

AUTHOR(S): Niitani, Hisanobu; Kawase, Ichiro; Jaijo, Nagahiro;

Taniguchi, Takeshi

CORPORATE SOURCE: Natl. Cancer Cent. Hosp., Tokyo, Japan

SOURCE: Gan to Kagaku Ryoho (1979), 6(Rinji Zokan

2), 213-18

CODEN: GTKRDX; ISSN: 0385-0684

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB The content of both coenzyme Q9 [303-97-9] and coenzyme Q10 [303-98-0] in spleen lymphocytes decreased in rats bearing Sato lung carcinoma. Oligomycin-sensitive ATPase [9000-83-3]

activity in spleen lymphocytes was also depressed. The depressed, oligomycin-sensitive ATPase activity was recovered by i.m. administration of coenzyme Q10 emulsified with EtOH and saline, and the decreased content of coenzyme Q9 and Q10 was slightly restored by this treatment. This enzyme activity was also significantly recovered by an i.v. administration of BCG, and was elevated more by the combined treatment with BCG and the emulsified coenzyme Q10.

L7 ANSWER 14 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1979:517384 CAPLUS

DOCUMENT NUMBER: 91:117384

ORIGINAL REFERENCE NO.: 91:18841a,18844a

TITLE: Combined effect of BCG and coenzyme Q10 on ATPase activity and coenzyme Q content in spleen lymphocytes of tumor-bearing rats

AUTHOR(S): Niotani, Hisanobu; Kawase, Ichiro; Taniguchi, Takeshi; Saijo, Nagahiro; Irimajiri, Nobuhiro

CORPORATE SOURCE: Natl. Cancer Cent. Hosp., Tokyo, 104, Japan


SOURCE: Gann (1979), 70(3), 315-22

CODEN: GANNA2; ISSN: 0016-450X

DOCUMENT TYPE: Journal

LANGUAGE: English

GI



I

AB Effect of Mycobacterium BCG, coenzyme Q10 (I) [303-98-0], or their combination on ATPase activity in spleen lymphocytes of tumor-bearing rats was investigated in relation to changes in the content of individual coenzyme Q homologs in these cells. Contents of both coenzyme Q9 [303-97-9] and I in spleen lymphocytes significantly decreased in the late stage of Donryu rats bearing Sato lung carcinoma. Oligomycin-sensitive ATPase [9000-83-3] activity in spleen lymphocytes was also significantly depressed in this stage. The depressed, oligomycin-sensitive ATPase activity was significantly recovered by a 3-time i.m. administration of I emulsified with EtOH and saline, and the decreased contents of coenzymes Q9 and I were slightly restored by this treatment. This enzyme activity was also significantly recovered by an i.v. administration of BCG, and was elevated more by the combined treatment with BCG and the emulsified I. Apparently, the combined treatment with BCG and emulsified I can contribute to the improvement of the depressed bioenergetics in lymphocytes of tumor-bearing animals, and this combined effect of BCG and emulsified I might be based on the combination of their individual activating effects on lymphocytes.

L7 ANSWER 15 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1968:10731 CAPLUS

DOCUMENT NUMBER: 68:10731

ORIGINAL REFERENCE NO.: 68:2027a,2030a

TITLE: Biosynthetic changes of vitamin K3 and ubiquinone 0 in man

AUTHOR(S): Ritzl, Friedrich

SOURCE: Berichte der Kernforschungsanlage Juelich ( 1966), No. 420-ME, 57 pp.  
CODEN: BKEJAS; ISSN: 0366-0885

DOCUMENT TYPE: Journal  
LANGUAGE: German

AB A number of K-vitamins are known to alleviate vitamin K deficiencies in man; however, the K vitamin specific for human function is still unknown. On the other hand the effective ubiquinone (I) in humans has been shown to be I(10). Nevertheless after injections of I(0) a series of ubiquinones in addition to I(10) were identified in humans. In order to clarify some of these points, 3H-labeled menadione (II) (vitamin K3), as either the Na bisulfite or tetrasodium diphosphate compds., and 3H-labeled I(0) were injected into human test subjects. These subjects were 25 patients with normal livers and 18 patients with inoperable bronchial carcinoma, without liver metastases. The radioactivity in the serum, urine, and bile were determined as a function of time. Specific 3H-labeled K vitamins and ubiquinones were separated and identified by countercurrent distribution in the system n-heptane-methylglycol-water (10:7:3) or by column chromatog. on kieselgel with CHCl3. Twenty min. after the injection of II, vitamin K2 (20) was detected in the blood stream, indicating that vitamin K2 (20) is the specific K vitamin for humans. Small quantities of vitamins K2 (45) and (50) were also detected, but only after the injection of high specific activity II. More radioactivity was detected in the bile after injection of II than after injection of I(0). Also most of the radioactivity was excreted in the urine following injection of I(0), while only 50% of the radioactivity was excreted after injection of II. The main radioactive components in blood and urine following the injection of I(0) were I(10), I(9), and occasionally I(6). A discussion of the distribution of the K vitamins and ubiquinones in microorganisms, and the functional efficiency of the various K vitamins and ubiquinone homologs is presented.

L7 ANSWER 16 OF 16 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1962:34409 CAPLUS

DOCUMENT NUMBER: 56:34409

ORIGINAL REFERENCE NO.: 56:6539c-d

TITLE: Histochemical studies of the effects of coenzyme Q10 and menadione on oxidative enzymes in normal and neoplastic cells

AUTHOR(S): Wattenberg, Lee W.

CORPORATE SOURCE: Univ. of Minnesota, Minneapolis

SOURCE: Ciba Foundation Symposium Quinones Electron Transport (1961), 1960, 367-84

DOCUMENT TYPE: Journal

LANGUAGE: Unavailable

AB Two flavoprotein-tetrazolium salt reduction systems, succinate-2-(p-iodophenyl)-3-(p-nitrophenyl)-5-phenyltetrazolium chloride reductase and  $\alpha$ -glycerophosphate-2-(p-iodophenyl)-3-(p-nitrophenyl)-5-phenyltetrazolium chloride reductase, in which coenzyme Q10 serves as an electron-transport agent are unsatd. with respect to quinone in rat liver. Normal liver shows less unsatn. than regenerating tissue and hepatoma. Coenzyme Q10 is very effective in hepatoma but not in regenerating liver, where menadione is effective. Carcinoma of the large bowel has shown a marked reductase enhancement in response to added coenzyme Q10.

=> s 15 and carcinoma and topical

5610 L5

222424 CARCINOMA

39195 CARCINOMAS

179 CARCINOMATA

1 CARCINOMATAS  
 231654 CARCINOMA  
 (CARCINOMA OR CARCINOMAS OR CARCINOMATA OR CARCINOMATAS)  
 60702 TOPICAL  
 49 TOPICALS  
 60723 TOPICAL  
 (TOPICAL OR TOPICALS)  
 L8 7 L5 AND CARCINOMA AND TOPICAL

=> d ibib abs total

L8 ANSWER 1 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2008:1156621 CAPLUS  
 DOCUMENT NUMBER: 149:409737  
 TITLE: Topical formulations comprising lipophilic  
 bioactive agents having enhanced bioavailability  
 INVENTOR(S): McCook, John Patrick; Narain, Niven Rajin; Persaud,  
 Indushekhkar  
 PATENT ASSIGNEE(S): Pathfinder Management, Inc., USA  
 SOURCE: PCT Int. Appl., 68pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008116135	A2	20080925	WO 2008-US57786	20080321
WO 2008116135	A3	20081224		
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
AU 2008228764	A1	20080925	AU 2008-228764	20080321
CA 2680825	A1	20080925	CA 2008-2680825	20080321
US 20080233183	A1	20080925	US 2008-52825	20080321
EP 2136787	A2	20091230	EP 2008-732635	20080321
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR			
NO 2009003032	A	20091022	NO 2009-3032	20090921
MX 2009010170	A	20091126	MX 2009-10170	20090922
PRIORITY APPLN. INFO.:			US 2007-919554P	P 20070322
			WO 2008-US57786	W 20080321

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present disclosure provides compns. suitable for delivering lipophilic bioactive agents. The compns. may be utilized to treat numerous diseases and conditions that would benefit from the application of a lipophilic bioactive agent. Thus, a cream contained Polysorbate-80 25.000, ubidecarenone 21.000, propylene glycol 10.000, phenoxyethanol 0.500, water 35.500, and lecithin 8.000%.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD



## (1 CITINGS)

L8 ANSWER 2 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2008:978432 CAPLUS  
 DOCUMENT NUMBER: 149:259457  
 TITLE: Method of cancer screening; method of cancer treatment; and method of auto-immune disease treatment  
 INVENTOR(S): Woodward, John R.  
 PATENT ASSIGNEE(S): Les Medecins L.P., USA  
 SOURCE: U.S. Pat. Appl. Publ., 14 pp., Cont.-in-part of U.S. Ser. No. 533,805.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 7  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080193482	A1	20080814	US 2008-100089	20080409
US 20060063211	A1	20060323	US 2004-946213	20040921
US 20060063212	A1	20060323	US 2004-3293	20041203
US 20060062755	A1	20060323	US 2005-32399	20050110
US 20060062757	A1	20060323	US 2005-133838	20050519
US 7125836	B2	20061024		
US 20070014821	A1	20070118	US 2006-533805	20060921
US 7507703	B2	20090324		
PRIORITY APPLN. INFO.:			US 2004-946213	B2 20040921
			US 2004-3293	B2 20041203
			US 2005-32399	B3 20050110
			US 2005-133838	A1 20050519
			US 2006-533805	A2 20060921

## ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A method of cancer screening comprising the steps of administering the Blood CA 27,29 testing procedure; if the result is pos. administering a mammogram; if the result is pos. administering a needle biopsy; if the result is pos. administering a PET scan; if the result is pos. administering a blood tumor cell count. If all of the foregoing steps are pos., the cancer is treated by selecting one or more treatments from a group of provided treatment according to the patient's body and condition. A method of treating auto-immune diseases comprises selecting one or more treatments from another group of provided treatments, the one or more treatments selected and administered according to the patient's body and condition.

L8 ANSWER 3 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2008:556974 CAPLUS  
 DOCUMENT NUMBER: 148:529432  
 TITLE: Composition, its use for treating systemic diseases a conditions, and product containing said composition  
 INVENTOR(S): Lindblom, Ragnvald Erik; Lindblom, Jonas Erik; De Faire, Johan; Janchanakit, Jirawat  
 PATENT ASSIGNEE(S): Salutory Care Limited, Cyprus  
 SOURCE: PCT Int. Appl., 49pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

WO 2008054293 A1 20080508 WO 2007-SE970 20071101  
 W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA,  
 CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI,  
 GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG,  
 KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME,  
 MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL,  
 PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN,  
 TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW  
 RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF,  
 BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG, BW,  
 GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ,  
 BY, KG, KZ, MD, RU, TJ, TM  
 PRIORITY APPLN. INFO.: SE 2006-2333 A 20061103  
 SE 2006-2334 A 20061103  
 AB The invention relates to a pharmaceutical composition useful for treating or  
 preventing cancerous diseases, HIV/AIDS, chronic inflammations, certain  
 autoimmune diseases, and secondary conditions of such primary diseases,  
 such as bacterial, viral and fungous infections, inflammations, diarrhea,  
 dehydration, and pain, said composition comprising a catalytic product (based  
 on a serine protease extracted and isolated from fish, molluscs and crustacean  
 species), named Mecosome, and a microbial agent (comprising the strains  
 Pedicoccus pentosaceus, Pichia farinosa, Dekkera bruxellensis), named  
 M-powder, to the use of said composition, and to a product containing said  
 composition  
 The composition may further comprise a chemical agent, named Mesodine, and a  
 herbal component, named Phumpat. The invention also relates to the use of  
 the last mentioned composition, and to a product containing said last mentioned  
 composition  
 REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT  
 L8 ANSWER 4 OF 7 CAPLUS COPYRIGHT 2010 ACS on SIN  
 ACCESSION NUMBER: 2008:349028 CAPLUS  
 DOCUMENT NUMBER: 148:338999  
 TITLE: Foamable vehicle and vitamin and flavonoid  
 pharmaceutical compositions thereof for treatment of  
 skin and other disorders  
 INVENTOR(S): Tamarkin, Dov; Friedman, Doron; Eini, Meir; Berman,  
 Tal; Schuz, David  
 PATENT ASSIGNEE(S): Foamix Ltd., Israel  
 SOURCE: U.S. Pat. Appl. Publ., 57pp., Cont.-in-part of U.S.  
 Ser. No. 430,599.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 35  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080069779	A1	20080320	US 2007-900072	20070910
US 20050031547	A1	20050210	US 2004-835505	20040428
AU 2004313285	A1	20050929	AU 2004-313285	20041216
ZA 2005007018	A	20080227	ZA 2005-7018	20041216
US 20060275218	A1	20061207	US 2006-430599	20060509
AU 2006298442	A1	20070412	AU 2006-298442	20060509
CA 2609953	A1	20070412	CA 2006-2609953	20060509
WO 2007039825	A2	20070412	WO 2006-1B3628	20060509
WO 2007039825	A3	20080306		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA

AU 2006313443 A1 20070518 AU 2006-313443 20060509  
CA 2610662 A1 20070518 CA 2006-2610662 20060509  
WO 2007054818 A2 20070518 WO 2006-IB3519 20060509  
WO 2007054818 A3 20081023

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA

EP 1888032 A2 20080220 EP 2006-831721 20060509  
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU

EP 1893396 A2 20080305 EP 2006-809259 20060509  
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU

JP 2008540508 T 20081120 JP 2008-510676 20060509  
JP 2008540511 T 20081120 JP 2008-510679 20060509  
US 20070280891 A1 20071206 US 2006-645444 20061226  
ZA 2007010621 A 20090325 ZA 2007-10621 20070101  
US 20080050317 A1 20080228 US 2007-894668 20070820  
MX 2007014106 A 20080829 MX 2007-14106 20071109  
MX 2007014101 A 20090213 MX 2007-14101 20071109  
IN 2007KN04432 A 20080125 IN 2007-KN4432 20071203  
IN 2007KN04590 A 20080704 IN 2007-KN4590 20071203  
ZA 2007010619 A 20090826 ZA 2007-10619 20071204

PRIORITY APPLN. INFO.:

US 2003-492385P P 20030804  
US 2003-530015P P 20031216  
US 2004-835505 A2 20040428  
US 2005-679020P P 20050509  
US 2006-784793P P 20060321  
US 2006-430599 A2 20060509  
US 2006-843140P P 20060908  
WO 2006-IB3519 W 20060509  
WO 2006-IB3628 W 20060509

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Vitamin and flavonoid containing compns. are provided that are stable to degradation. Stabilized compns. include one or more features including a hygroscopic solvent at a sufficient concentration to provide an Aw value of the hygroscopic vitamin and or flavonoid containing composition of less than 0.9, antioxidant flavonoids that are preferentially oxidized before the

vitamin, preservatives, and hydrocarbon propellants selected to reduce the oxidation potential of the composition. Thus, a foamable carrier was prepared containing

propylene glycol 88.00, stearyl alc. 2.00, hydroxypropyl cellulose 2.00, Laureth-4 2.00, GMS NE 2.00, macrogol cetostearyl ether 1.00, and PPG-15 stearyl ether 3.00%, resp. Ascorbic acid and niacinamide were concurrently added to the carrier at 5.00% and 2.00%, resp. Following addition of a propellant, the foamable composition was obtained, which upon release from an aerosol pressurized container afforded foam of good quality. The foam was easily spread and immediately absorbed into the facial skin with no extensive rubbing.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L8 ANSWER 5 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:1215784 CAPLUS

DOCUMENT NUMBER: 147:491621

TITLE: Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer

INVENTOR(S): Mazzio, Elizabeth; Soliman, Karam

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S.

Ser. No. 233,279.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070248693	A1	20071025	US 2007-711883	20070227
US 20060035981	A1	20060216	US 2005-233279	20050920
PRIORITY APPLN. INFO.:			US 2003-491841P	P 20030802
			US 2004-540525P	P 20040129
			US 2004-909590	B2 20040802
			US 2005-233279	A2 20050920

AB The invention describes a pharmaceutical composition and method for treating cancer comprising (a) 2,3-dimethoxy-5-methyl-1,4-benzoquinone, and/or (b) at least one of wild yam root, teasel root, balm of gilead bud, bakuchi seed, dichroa root, kochia seed, kanta kari, bushy knotweed rhizome, arjun, babul chall bark, opopanax and bhumi amalaki; optionally one or more of frankincense, garcinia fruit, vitex, dragons blood, mace, sage and red sandalwood with at least (c) one compound capable of maximizing oxidative mitochondrial function, preferably riboflavin or vitamin B2 derivs., FAD, FMN, 5-amino-6-(5'-phosphoribitylamino)uracil, 6,7-dimethyl-8-(1-D-ribityl)lumazine, ribitol, 5,6-dimethylbenzimidazole, tetrahydrobiopterin, vitamin B1, lipoic acid, biotin, vitamin B6, vitamin B12, folate, niacin, vitamin C and pantothenate, and/or (d) at least one lactic acid dehydrogenase inhibitor, preferably 2',3,4',5',7-pentahydroxyflavone and optionally (f) an alkalinizing agent (Aloe vera, chlorella, wheat grass, sodium or potassium bicarbonate, potassium), (g) an antiproliferative herb (speranskia or goldenseal), and (h) a pharmaceutically acceptable carrier. A method for inhibiting cancer optionally comprises one or more chemotherapy drug(s), selected, among others, from acetogenins, actinomycin D, adriamycin, aminoglutethimide, asparaginase, bleomycin, bullatacin, busulfan, carmustine, carboplatin, chlorambucil, cisplatin, etc. Thus, a composition comprised rosemary (Rosmarinus officinalis) .apprx.1000, myrrh gum (Commiphora molmol) .apprx.500, 2,3-dimethoxy-5-methyl-1,4 benzoquinone .apprx.800, and

riboflavin .apprx.300 mg/day, resp.

L8 ANSWER 6 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:458900 CAPLUS

DOCUMENT NUMBER: 146:427847

TITLE: Topically applied glucosamine sulfate and all its related, precursor, and derivative compounds significantly increases the skin's natural production of hyaluronic acid for the rejuvenation of healthier younger-looking skin; while phosphatidylcholine is required to replace its deficiency caused by topical dimethylaminoethanol (DMAE)

INVENTOR(S): Jacobs, Eric

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 13pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
US 20070092469	A1	20070426	US 2006-527334	20060927
PRIORITY APPLN. INFO.:			US 2005-729947P	P 20051026

AB A topical skin rejuvenation preparation comprises (i) about 0.001 to 50% of glucosamine (2-amino-2-deoxy- $\alpha$ -D-glucose), a hexosamine (6 carbon amino sugar), including its derivative and precursor compds., glucosamine sulfate, glucosamine hydrochloride, glucose-6-phosphate, acetyl glucosamine, fructose-6-phosphate, and glucosamine-6-phosphate to increase production of hyaluronic acid and collagen and to relieve wrinkles, increase the skin's natural production of hyaluronic acid, reverse the lack of suppleness, hydrate from within, erase spider veins, reduce varicose veins, lighten aging dark blotches ("liver spots"/lentigos, senile lentigines), decrease acne, and reduce under eye puffiness, (ii) 0.0001 to 50% of dimethylaminoethanol (DMAE) to increase skin muscle tone, and (iii) 0.01 to 30% of phosphatidylcholine to overcome deficiency created by application of DMAE in each cell's production of phosphatidylcholine, whose deficiency damages cell membranes, as well as mitochondrial and lysosome membranes.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L8 ANSWER 7 OF 7 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2006:365124 CAPLUS

DOCUMENT NUMBER: 144:398343

TITLE: Methods and compositions for the treatment of diseases characterized by calcification and/or plaque formation  
INVENTOR(S): Kajander, E. Olavi; Aho, K.; Ciftcioglu, Neva; Millican, H. B.; Maniscalco, B.

PATENT ASSIGNEE(S): Nanobac Pharmaceuticals, Inc., USA

SOURCE: U.S. Pat. Appl. Publ., 14 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
US 20060083727	A1	20060420	US 2005-182076	20050715



TITLE: Topical formulations comprising lipophilic bioactive agents having enhanced bioavailability  
 INVENTOR(S): McCook, John Patrick; Narain, Niven Rajin; Persaud, Indushekhar  
 PATENT ASSIGNEE(S): Pathfinder Management, Inc., USA  
 SOURCE: PCT Int. Appl., 68pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008116135	A2	20080925	WO 2008-US57786	20080321
WO 2008116135	A3	20081224		
W: AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
AU 2008228764	A1	20080925	AU 2008-228764	20080321
CA 2680825	A1	20080925	CA 2008-2680825	20080321
US 20080233183	A1	20080925	US 2008-52825	20080321
EP 2136787	A2	20091230	EP 2008-732635	20080321
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR				
NO 2009003032	A	20091022	NO 2009-3032	20090921
MX 2009010170	A	20091126	MX 2009-10170	20090922
PRIORITY APPLN. INFO.:			US 2007-919554P	P 20070322
			WO 2008-US57786	W 20080321

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present disclosure provides compns. suitable for delivering lipophilic bioactive agents. The compns. may be utilized to treat numerous diseases and conditions that would benefit from the application of a lipophilic bioactive agent. Thus, a cream contained Polysorbate-80 25.000, ubidecarenone 21.000, propylene glycol 10.000, phenoxyethanol 0.500, water 35.500, and lecithin 8.000%.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

=> d ibib abs 110 total

L10 ANSWER 1 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:1156621 CAPLUS  
 DOCUMENT NUMBER: 149:409737  
 TITLE: Topical formulations comprising lipophilic bioactive agents having enhanced bioavailability  
 INVENTOR(S): McCook, John Patrick; Narain, Niven Rajin; Persaud, Indushekhar  
 PATENT ASSIGNEE(S): Pathfinder Management, Inc., USA  
 SOURCE: PCT Int. Appl., 68pp.

DOCUMENT TYPE: CODEN: PIXXD2  
 LANGUAGE: Patent  
 FAMILY ACC. NUM. COUNT: English  
 PATENT INFORMATION: 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008116135	A2	20080925	WO 2008-US57786	20080321
WO 2008116135	A3	20081224		
W:				
AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW				
RW:				
AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
AU 2008228764	A1	20080925	AU 2008-228764	20080321
CA 2680825	A1	20080925	CA 2008-2680825	20080321
US 20080233183	A1	20080925	US 2008-52825	20080321
EP 2136787	A2	20091230	EP 2008-732635	20080321
R:				
AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LI, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR				
NO 2009003032	A	20091022	NO 2009-3032	20090921
MX 2009010170	A	20091126	MX 2009-10170	20090922

PRIORITY APPLN. INFO.:

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present disclosure provides compns. suitable for delivering lipophilic bioactive agents. The compns. may be utilized to treat numerous diseases and conditions that would benefit from the application of a lipophilic bioactive agent. Thus, a cream contained Polysorbate-80 25.000, ubidecarenone 21.000, propylene glycol 10.000, phenoxyethanol 0.500, water 35.500, and lecithin 8.000%.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L10 ANSWER 2 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:978432 CAPLUS

DOCUMENT NUMBER: 149:259457

TITLE: Method of cancer screening; method of cancer treatment; and method of auto-immune disease treatment  
 Woodward, John R.

INVENTOR(S): Les Medecins L.P., USA

PATENT ASSIGNEE(S): U.S. Pat. Appl. Publ., 14 pp., Cont.-in-part of U.S.

SOURCE: Ser. No. 533,805.  
 CODEN: USXXCO

DOCUMENT TYPE: Patent  
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 7  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080193482	A1	20080814	US 2008-100089	20080409



US 20060063211	A1	20060323	US 2004-946213	20040921
US 20060063212	A1	20060323	US 2004-3293	20041203
US 20060062755	A1	20060323	US 2005-32399	20050110
US 20060062757	A1	20060323	US 2005-133838	20050519
US 7125836	B2	20061024		
US 20070014821	A1	20070118	US 2006-533805	20060921
US 7507703	B2	20090324		

PRIORITY APPLN. INFO.:

US 2004-946213	B2	20040921
US 2004-3293	B2	20041203
US 2005-32399	B3	20050110
US 2005-133838	A1	20050519
US 2006-533805	A2	20060921

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A method of cancer screening comprising the steps of administering the Blood CA 27,29 testing procedure; if the result is pos. administering a mammogram; if the result is pos. administering a needle biopsy; if the result is pos. administering a PET scan; if the result is pos. administering a blood tumor cell count. If all of the foregoing steps are pos., the cancer is treated by selecting one or more treatments from a group of provided treatment according to the patient's body and condition. A method of treating auto-immune diseases comprises selecting one or more treatments from another group of provided treatments, the one or more treatments selected and administered according to the patient's body and condition.

L10 ANSWER 3 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:556974 CAPLUS  
 DOCUMENT NUMBER: 148:529432  
 TITLE: Composition, its use for treating systemic diseases a conditions, and product containing said composition  
 INVENTOR(S): Lindblom, Ragnvald Erik; Lindblom, Jonas Erik; De Faire, Johan; Janchanakit, Jirawat  
 PATENT ASSIGNEE(S): Salutory Care Limited, Cyprus  
 SOURCE: PCT Int. Appl., 49pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008054293	A1	20080508	WO 2007-SE970	20071101
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

PRIORITY APPLN. INFO.:

SE 2006-2333	A	20061103
SE 2006-2334	A	20061103

AB The invention relates to a pharmaceutical composition useful for treating or preventing cancerous diseases, HIV/AIDS, chronic inflammations, certain autoimmune diseases, and secondary conditions of such primary diseases, such as bacterial, viral and fungous infections, inflammations, diarrhea,

dehydration, and pain, said composition comprising a catalytic product (based on a serine protease extracted and isolated from fish, molluscs and crustacean species), named Mecosome, and a microbial agent (comprising the strains *Pediococcus pentosaceus*, *Pichia farinosa*, *Dekkera bruxellensis*), named M-powder, to the use of said composition, and to a product containing said composition

The composition may further comprise a chemical agent, named Mesodine, and a herbal component, named Phumpat. The invention also relates to the use of the last mentioned composition, and to a product containing said last mentioned composition

REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 4 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:349028 CAPLUS

DOCUMENT NUMBER: 148:338999

TITLE: Foamable vehicle and vitamin and flavonoid pharmaceutical compositions thereof for treatment of skin and other disorders

INVENTOR(S): Tamarkin, Dov; Friedman, Doron; Eini, Meir; Berman, Tal; Schuz, David

PATENT ASSIGNEE(S): Foamix Ltd., Israel

SOURCE: U.S. Pat. Appl. Publ., 57pp., Cont.-in-part of U.S. Ser. No. 430,599.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 35

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080069779	A1	20080320	US 2007-900072	20070910
US 20050031547	A1	20050210	US 2004-835505	20040428
AU 2004313285	A1	20050929	AU 2004-313285	20041216
ZA 2005007018	A	20080227	ZA 2005-7018	20041216
US 20060275218	A1	20061207	US 2006-430599	20060509
AU 2006298442	A1	20070412	AU 2006-298442	20060509
CA 2609953	A1	20070412	CA 2006-2609953	20060509
WO 2007039825	A2	20070412	WO 2006-IB3628	20060509
WO 2007039825	A3	20080306		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
AU 2006313443	A1	20070518	AU 2006-313443	20060509
CA 2610662	A1	20070518	CA 2006-2610662	20060509
WO 2007054818	A2	20070518	WO 2006-IB3519	20060509
WO 2007054818	A3	20081023		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX,			

MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW

RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA

EP 1888032 A2 20080220 EP 2006-831721 20060509

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU

EP 1893396 A2 20080305 EP 2006-809259 20060509

R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU

JP 2008540508 T 20081120 JP 2008-510676 20060509

JP 2008540511 T 20081120 JP 2008-510679 20060509

US 20070280891 A1 20071206 US 2006-645444 20061226

ZA 2007010621 A 20090325 ZA 2007-10621 20070101

US 20080050317 A1 20080228 US 2007-894668 20070820

MX 2007014106 A 20080829 MX 2007-14106 20071109

MX 2007014101 A 20090213 MX 2007-14101 20071109

IN 2007KN04432 A 20080125 IN 2007-KN4432 20071203

IN 2007KN04590 A 20080704 IN 2007-KN4590 20071203

ZA 2007010619 A 20090826 ZA 2007-10619 20071204

PRIORITY APPLN. INFO.: US 2003-492385P P 20030804

US 2003-530015P P 20031216

US 2004-835505 A2 20040428

US 2005-679020P P 20050509

US 2006-784793P P 20060321

US 2006-430599 A2 20060509

US 2006-843140P P 20060908

WO 2006-IB3519 W 20060509

WO 2006-IB3628 W 20060509

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Vitamin and flavonoid containing compns. are provided that are stable to degradation. Stabilized compns. include one or more features including a hygroscopic solvent at a sufficient concentration to provide an Aw value of the hygroscopic vitamin and/or flavonoid containing composition of less than 0.9, antioxidant flavonoids that are preferentially oxidized before the vitamin, preservatives, and hydrocarbon propellants selected to reduce the oxidation potential of the composition. Thus, a foamable carrier was prepared containing propylene glycol 88.00, stearyl alc. 2.00, hydroxypropyl cellulose 2.00, Laureth-4 2.00, GMS NE 2.00, macrogol cetostearyl ether 1.00, and PPG-15 stearyl ether 3.00%, resp. Ascorbic acid and niacinamide were concurrently added to the carrier at 5.00% and 2.00%, resp. Following addition of a propellant, the foamable composition was obtained, which upon release from an aerosol pressurized container afforded foam of good quality. The foam was easily spread and immediately absorbed into the facial skin with no extensive rubbing.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

L10 ANSWER 5 OF 8 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 2007:1215784 CAPLUS

DOCUMENT NUMBER: 147:491621

TITLE: Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer

INVENTOR(S): Mazzio, Elizabeth; Soliman, Karam  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S.  
 Ser. No. 233,279.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070248693	A1	20071025	US 2007-711883	20070227
US 20060035981	A1	20060216	US 2005-233279	20050920
PRIORITY APPLN. INFO.:			US 2003-491841P	P 20030802
			US 2004-540525P	P 20040129
			US 2004-909590	B2 20040802
			US 2005-233279	A2 20050920

AB The invention describes a pharmaceutical composition and method for treating cancer comprising (a) 2,3-dimethoxy-5-methyl-1,4-benzoquinone, and/or (b) at least one of wild yam root, teasel root, balm of gilead bud, bakuchi seed, dichroa root, kochia seed, kanta kari, bushy knotweed rhizome, arjun, babul chall bark, opopanax and bhumy amalaki; optionally one or more of frankincense, garcinia fruit, vitex, dragons blood, mace, sage and red sandalwood with at least (c) one compound capable of maximizing oxidative mitochondrial function, preferably riboflavin or vitamin B2 derivs., FAD, FMN, 5-amino-6-(5'-phosphoribitylamino)uracil, 6,7-dimethyl-8-(1-D-ribityl)lumazine, ribitol, 5,6-dimethylbenzimidazole, tetrahydrobiopterin, vitamin B1, liponic acid, biotin, vitamin B6, vitamin B12, folate, niacin, vitamin C and pantothenate, and/or (d) at least one lactic acid dehydrogenase inhibitor, preferably 2',3,4',5,7-pentahydroxyflavone and optionally (f) an alkalizing agent (Aloe vera, chlorella, wheat grass, sodium or potassium bicarbonate, potassium), (g) an antiproliferative herb (speranskia or goldenseal), and (h) a pharmaceutically acceptable carrier. A method for inhibiting cancer optionally comprises one or more chemotherapy drug(s), selected, among others, from acetogenins, actinomycin D, adriamycin, aminoglutethimide, asparaginase, bleomycin, bullatacin, busulfan, carmustine, carboplatin, chlorambucil, cisplatin, etc. Thus, a composition comprised rosemary (Rosmarinus officinalis) .apprx.1000, myrrh gum (Commiphora molmol) .apprx.500, 2,3-dimethoxy-5-methyl-1,4 benzoquinone .apprx.800, and riboflavin .apprx.300 mg/day, resp.

L10 ANSWER 6 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:458900 CAPLUS

DOCUMENT NUMBER: 146:427847

TITLE: Topically applied glucosamine sulfate and all its related, precursor, and derivative compounds significantly increases the skin's natural production of hyaluronic acid for the rejuvenation of healthier younger-looking skin; while phosphatidylcholine is required to replace its deficiency caused by topical dimethylaminoethanol (DMAE)

INVENTOR(S): Jacobs, Eric  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 13pp.  
 CODEN: USXXCO

DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070092469	A1	20070426	US 2006-527334	20060927
PRIORITY APPLN. INFO.:			US 2005-729947P	P 20051026

AB A topical skin rejuvenation preparation comprises (i) about 0.001 to 50% of glucosamine (2-amino-2-deoxy-alpha-D-glucose), a hexosamine (6 carbon amino sugar), including its derivative and precursor compds., glucosamine sulfate, glucosamine hydrochloride, glucose-6-phosphate, acetyl glucosamine, fructose-6-phosphate, and glucosamine-6-phosphate to increase production of hyaluronic acid and collagen and to relieve wrinkles, increase the skin's natural production of hyaluronic acid, reverse the lack of suppleness, hydrate from within, erase spider veins, reduce varicose veins, lighten aging dark blotches ("liver spots"/lentigos, senile lentigines), decrease acne, and reduce under eye puffiness, (ii) 0.0001 to 50% of dimethylaminoethanol (DMAE) to increase skin muscle tone, and (iii) 0.01 to 30% of phosphatidylcholine to overcome deficiency created by application of DMAE in each cell's production of phosphatidylcholine, whose deficiency damages cell membranes, as well as mitochondrial and lysosome membranes.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L10 ANSWER 7 OF 8 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2006:691731 CAPLUS

DOCUMENT NUMBER: 145:262676

TITLE: NAD+/NADH and/or CoQ/CoQH2 ratios from plasma membrane electron transport may determine ceramide and sphingosine-1-phosphate levels accompanying G1 arrest and apoptosis

AUTHOR(S): De Luca, Thomas; Morre, Dorothy M.; Zhao, Haiyun; Morre, D. James

CORPORATE SOURCE: Department of Foods and Nutrition, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: BioFactors (2005), 25(1-4), 43-60  
CODEN: BIFAEU; ISSN: 0951-6433

PUBLISHER: IOS Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB To elucidate possible biochem. links between growth arrest from antiproliferative chemotherapeutic agents and apoptosis, our work has focused on agents (EGCG, capsaicin, cis platinum, adriamycin, anti-tumor sulfonylureas, phenoxodiol) that target tNOX. tNOX is a cancer-specific cell surface NADH oxidase (ECTO-NOX protein), that functions in cancer cells as the terminal oxidase for plasma membrane electron transport. When tNOX is active, coenzyme Q10 (ubiquinone) of the plasma membrane is oxidized and NADH is oxidized at the cytosolic surface of the plasma membrane. However, when tNOX is inhibited and plasma membrane electron transport is diminished, both reduced coenzyme Q10 (ubiquinol) and NADH would be expected to accumulate. To relate inhibition of plasma membrane redox to increased ceramide levels and arrest of cell proliferation in G1 and apoptosis, we show that neutral sphingomyelinase, a major contributor to plasma membrane ceramide, is inhibited by reduced glutathione and ubiquinone. Ubiquinol is without effect or stimulates. In contrast, sphingosine kinase, which generates anti-apoptotic sphingosine-1-phosphate, is stimulated by ubiquinone but inhibited by ubiquinol and NADH. Thus, the quinone and pyridine nucleotide products of plasma membrane redox, ubiquinone and ubiquinol, as well as NAD+ and NADH, may directly modulate in a reciprocal manner two key plasma membrane enzymes, sphingomyelinase and sphingosine kinase, potentially leading to G1 arrest (increase in ceramide) and apoptosis

(loss of sphingosine-1-phosphate). As such, the findings provide potential links between coenzyme Q10-mediated plasma membrane electron transport and the anticancer action of several clin.-relevant anticancer agents.

OS.CITING REF COUNT: 14 THERE ARE 14 CAPLUS RECORDS THAT CITE THIS RECORD (14 CITINGS)  
REFERENCE COUNT: 61 THERE ARE 61 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L10 ANSWER 8 OF 8 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2006:365124 CAPLUS  
DOCUMENT NUMBER: 144:398343  
TITLE: Methods and compositions for the treatment of diseases characterized by calcification and/or plaque formation  
INVENTOR(S): Kajander, E. Olavi; Aho, K.; Ciftcioglu, Neva; Millican, H. B.; Maniscalco, B.  
PATENT ASSIGNEE(S): Nanobac Pharmaceuticals, Inc., USA  
SOURCE: U.S. Pat. Appl. Publ., 14 pp.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20060083727	A1	20060420	US 2005-182076	20050715
US 20070048296	A1	20070301	US 2006-544048	20061006
PRIORITY APPLN. INFO.:			US 2004-587871P	P 20040715
			US 2005-182076	A1 20050715

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention provides methods and compns. that include a nutraceutical supplement, antibiotic, and metal chelating agent that is administered to a patient to treat or prevent pathol. calcification and or plaque formation as associated with Nanobacteria Calcifying Nano-Particles and/or diseases caused there-from, The method includes the administration of a therapeutically effective nutraceutical supplement, tetracycline HCL, and EDTA calcium di-sodium salt to a patient in order to prevent and treat calcific disease.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

=> d his

(FILE 'HOME' ENTERED AT 13:27:42 ON 22 APR 2010)

FILE 'CAPLUS' ENTERED AT 13:29:20 ON 22 APR 2010  
S UBIQUINONE/CN

L1 FILE 'REGISTRY' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S UBIQUINONE/CN

L2 FILE 'CAPLUS' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S L1  
S COENZYME Q10/CN

L3 FILE 'REGISTRY' ENTERED AT 13:30:10 ON 22 APR 2010  
1 S COENZYME Q10/CN

FILE 'CAPLUS' ENTERED AT 13:30:10 ON 22 APR 2010

L4 5610 S L3

FILE 'REGISTRY' ENTERED AT 13:30:20 ON 22 APR 2010

L5 1 S COENZYME Q10/CN

FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010

L6 41 S L5 AND CARCINOMA  
L7 16 S L6 AND PY<=2004  
L8 7 S L5 AND CARCINOMA AND TOPICAL  
L9 38 S L5 AND ?CARCINOMA  
L10 8 S L9 AND (TOPICAL OR SURFACE)

=> s l5 and cancer and (topical or surface)

5610 L5  
452589 CANCER  
66393 CANCERS  
468904 CANCER  
(CANCER OR CANCERS)  
60702 TOPICAL  
49 TOPICALS  
60723 TOPICAL  
(TOPICAL OR TOPICALS)  
2995599 SURFACE  
540221 SURFACES  
3213094 SURFACE  
(SURFACE OR SURFACES)

L11 13 L5 AND CANCER AND (TOPICAL OR SURFACE)

=> l11 and py<=2004

L11 IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.

For a list of commands available to you in the current file, enter  
"HELP COMMANDS" at an arrow prompt (=>).

=> s l11 and py<=2004

25157969 PY<=2004

L12 5 L11 AND PY<=2004

=> d ibib abs

L12 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:832565 CAPLUS

DOCUMENT NUMBER: 137:329452

TITLE: Compositions with a non-glucocorticoid steroid and/or  
a ubiquinone and kit for treatment of respiratory and  
lung disease

INVENTOR(S): Nyce, Jonathan W.

PATENT ASSIGNEE(S): Epigenesis Pharmaceuticals, Inc., USA

SOURCE: PCT Int. Appl., 51 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
WO 2002085297	A2	20021031	WO 2002-US12555	20020422 <--
WO 2002085297	A3	20030403		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				

GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,  
 LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,  
 PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,  
 UA, UG, US, UZ, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,  
 KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,  
 GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA,  
 GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 AU 2002303427 A1 20021105 AU 2002-303427 20020422 <--  
 US 20040082522 A1 20040429 US 2003-454061 20030603 <--  
 US 7456161 B2 20081125  
 US 20080292709 A1 20081127 US 2008-172033 20080711  
 PRIORITY APPLN. INFO.: US 2001-286124P P 20010424  
 WO 2002-US12555 W 20020422  
 US 2003-454061 A3 20030603

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 137:329452

AB A pharmaceutical or veterinary composition comprises as the active agent (i) a non-glucocorticoid steroid or its analog, and (ii) a ubiquinone or their salts, in an amount effective for reducing levels of, or hypersensitivity to, adenosine, increasing levels of lung surfactant or ubiquinone, or for preventing or treating respiratory, lung and cancer diseases. The present treatment is useful for treating asthma, rhinitis, COPD, CF, RDS, pulmonary fibrosis, cancer and other diseases. For example, a metered dose inhaler contained ubiquinone 200 mg, dehydroepiandrosterone (DHEA) 200 mg, a stabilizer 5.0 µg, trichlorofluoromethane 23.70 mg, and dichlorodifluoromethane 61.25 mg.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib abs total

L12 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:832565 CAPLUS

DOCUMENT NUMBER: 137:329452

TITLE: Compositions with a non-glucocorticoid steroid and/or a ubiquinone and kit for treatment of respiratory and lung disease

INVENTOR(S): Nyce, Jonathan W.

PATENT ASSIGNEE(S): Epigenesis Pharmaceuticals, Inc., USA

SOURCE: PCT Int. Appl., 51 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002085297	A2	20021031	WO 2002-US12555	20020422 <--
WO 2002085297	A3	20030403		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				



KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,  
GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA,  
GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2002303427 A1 20021105 AU 2002-303427 20020422 <--  
US 20040082522 A1 20040429 US 2003-454061 20030603 <--  
US 7456161 B2 20081125  
US 20080292709 A1 20081127 US 2008-172033 20080711  
PRIORITY APPLN. INFO.: US 2001-286124P P 20010424  
WO 2002-US12555 W 20020422  
US 2003-454061 A3 20030603

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 137:329452

AB A pharmaceutical or veterinary composition comprises as the active agent (i) a non-glucocorticoid steroid or its analog, and (ii) a ubiquinone or their salts, in an amount effective for reducing levels of, or hypersensitivity to, adenosine, increasing levels of lung surfactant or ubiquinone, or for preventing or treating respiratory, lung and cancer diseases. The present treatment is useful for treating asthma, rhinitis, COPD, CF, RDS, pulmonary fibrosis, cancer and other diseases. For example, a metered dose inhaler contained ubiquinone 200 mg, dehydroepiandrosterone (DHEA) 200 mg, a stabilizer 5.0 µg, trichlorofluoromethane 23.70 mg, and dichlorodifluoromethane 61.25 mg.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:185691 CAPLUS

DOCUMENT NUMBER: 136:236872

TITLE: Epiandrosterones or ubiquinones for treatment of asthma and reduction of adenosine/adenosine receptor levels

INVENTOR(S): Nyce, Jonathan W.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 10 pp., Cont.-in-part of U.S. Ser. No. 488,236.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20020032160	A1	20020314	US 2001-841426	20010424 <--
US 5660835	A	19970826	US 1995-393863	19950224 <--
EP 1555025	A2	20050720	EP 2005-4694	19960215
EP 1555025	A3	20050803		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE				
US 6087351	A	20000711	US 1997-861962	19970522 <--
AU 9911317	A	19990304	AU 1999-11317	19990114 <--
AU 730453	B2	20010308		
US 6670349	B1	20031230	US 2000-488236	20000120 <--
US 20020119936	A1	20020829	US 2001-72010	20011025 <--
WO 2002085373	A1	20021031	WO 2002-US12489	20020422 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,				

UA, UG, UZ, VN, YU, ZA, ZM, ZW  
 RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH,  
 CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR,  
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2002254682	A1	20021105	AU 2002-254682	20020422 <--
JP 2005306880	A	20051104	JP 2005-162494	20050602
US 20060111306	A1	20060525	US 2005-275327	20051222
US 20090053143	A1	20090226	US 2008-196223	20080821
US 20090054385	A1	20090226	US 2008-196233	20080821

PRIORITY APPLN. INFO.:

US 1995-393863	A3	19950224
US 1997-861962	A1	19970522
US 2000-488236	A2	20000120
AU 1996-48677	A3	19960215
EP 1996-904622	A3	19960215
JP 1996-525728	A3	19960215
US 2001-841426	A3	20010424
US 2001-72010	B1	20011025
WO 2002-US12489	W	20020422

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 136:236872

AB A composition and various formulations comprise preventative or therapeutic amts. of an epiandrosterone, analog thereof or salt thereof, and/or a ubiquinone or salt thereof, and a pharmaceutically or veterinarily acceptable carrier or diluent. The composition and formulations are useful for treating bronchoconstriction, respiratory tract inflammation and allergies, asthma, and cancer. A method of treating diseases associated with low adenosine levels or adenosine depletion comprises administering folicinic acid or a pharmaceutically acceptable salt hereof in a preventative or therapeutic amount, or an amount effective to treat adenosine depletion. For example, rats administered DHEA or methyltestosterone daily for two weeks showed multi-organ depletion of adenosine. Depletion was dramatic in brain (60% depletion for DHEA, 34% for high dose methyltestosterone) and heart (37% depletion for DHEA, 22% depletion for high dose methyltestosterone). Coadministration of folicinic acid completely abrogated steroid-mediated adenosine depletion. Folicinic acid administered alone induce increase in adenosine levels for all organs studied. Also, both DHEA and ubiquinones inhibited NADPH levels in vitro by inhibiting the activity of glucose-6-phosphate dehydrogenase, an enzyme involved in the conversion of NADP to NADPH.

L12 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:573651 CAPLUS  
 DOCUMENT NUMBER: 133:159948  
 TITLE: Ubiquinone Qn for pain treatment  
 INVENTOR(S): Enzmann, Franz  
 PATENT ASSIGNEE(S): MSE Pharmazeutika G.m.b.H., Germany  
 SOURCE: PCT Int. Appl., 7 pp.  
 CODEN: PIXXD2

DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
WO 2000047192	A2	20000817	WO 2000-EP1011	20000209 <--
WO 2000047192	A3	20010412		
W: CA, JP, US				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
DE 19905879	A1	20000817	DE 1999-19905879	19990211 <--

CA 2362577	A1	20000817	CA 2000-2362577	20000209 <--
EP 1150682	A2	20011107	EP 2000-914075	20000209 <--
EP 1150682	B1	20050817		

R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI

AT 302009	T	20050915	AT 2000-914075	20000209
ES 2243243	T3	20051201	ES 2000-914075	20000209
US 20040034107	A1	20040219	US 2003-424987	20030429 <--

PRIORITY APPLN. INFO.: DE 1999-19905879 A 19990211  
WO 2000-EP1011 W 20000209  
US 2001-890276 B1 20010810

AB Ubiquinone Qn and its precursors can be used in the oral, parenteral, local, inhalative, or intranasal treatment of neurogenic pain, migraine, or pain resulting from dialysis, herpes zoster, cancer, etc. (no data).

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1999:359962 CAPLUS

DOCUMENT NUMBER: 131:181506

TITLE: The plasma membrane NADH oxidase of HeLa cells has hydroquinone oxidase activity

AUTHOR(S): Kishi, Takeo; Morre, Dorothy M.; Morre, D. James

CORPORATE SOURCE: Department of Medicinal Chemistry and Molecular Pharmacology, Purdue University, West Lafayette, IN, 47907, USA

SOURCE: Biochimica et Biophysica Acta, Bioenergetics (1999), 1412(1), 66-77

CODEN: BBBB4; ISSN: 0005-2728

PUBLISHER: Elsevier B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The plasma membrane NADH oxidase activity partially purified from the surface of HeLa cells exhibited hydroquinone oxidase activity. The preps. completely lacked NADH:ubiquinone reductase activity. However, in the absence of NADH, reduced coenzyme Q10 (Q10H2=ubiquinol) was oxidized at a rate of  $15 \pm 6$  nmol min<sup>-1</sup> mg protein<sup>-1</sup> depending on degree of purification. The apparent Km for Q10H2 oxidation was 33  $\mu$ M. Activities were inhibited competitively by the cancer cell-specific NADH oxidase inhibitors, capsaicin and the antitumor sulfonylurea N-(4-methylphenylsulfonyl)-N'-(4-chlorophenyl)urea (LY181984). With coenzyme Q0, where the preps. were unable to carry out either NADH:quinone reduction or reduced quinone oxidation, quinol oxidation

was observed with an equal mixture of the Q0 and Q0H2 forms. With the mixture, a rate of Q0H2 oxidation of  $8-17$  nmol min<sup>-1</sup> mg protein<sup>-1</sup> was observed with an apparent Km of 0.22 mM. The rate of Q10H2 oxidation was not stimulated by addition of equal amts. of Q10 and Q10H2. However, addition of Q0 to the Q10H2 did stimulate. The oxidation of Q10H2 proceeded with what appeared to be a two-electron transfer. The oxidation of Q0H2 may involve Q0, but the mechanism was not clear. The findings suggest the potential participation of the plasma membrane NADH oxidase as a terminal oxidase of plasma membrane electron transport from cytosolic NAD(P)H via naturally occurring hydroquinones to acceptors at the cell surface.

OS.CITING REF COUNT: 67 THERE ARE 67 CAPLUS RECORDS THAT CITE THIS RECORD (67 CITINGS)

REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L12 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1985:566163 CAPLUS  
 DOCUMENT NUMBER: 103:166163  
 ORIGINAL REFERENCE NO.: 103:26599a,26602a  
 TITLE: Treatment of radiation-induced ulcers by ubidecarenone  
 PATENT ASSIGNEE(S): Eisai Co., Ltd., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 5 pp.  
 CODEN: JKXXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 60100517	A	19850604	JP 1983-208395	19831108 <--
US 4617187	A	19861014	US 1984-666099	19841029 <--
EP 146742	A1	19850703	EP 1984-113349	19841106 <--
EP 146742	B1	19910130		

R: BE, CH, DE, FR, GB, IT, LI, NL, SE  
 PRIORITY APPLN. INFO.: JP 1983-208395 A 19831108  
 ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
 AB Topical formulations containing ubidecarenone [303-98-0]  
 are prepared for treatment of ulcers induced during the radiation therapy of  
 cancer and other diseases. Thus, a formulation consists of  
 stearyl alc. 5, stearic acid 2, lanolin 2, squalane 6, iso-Pr myristate 4,  
 polyoxyethylene cetyl alc. ether 3, glyceryl monostearate 2, ubidecarenone  
 0.3, propylene glycol 5, butylparaben 0.2, q.s. antioxidant, perfume, and  
 H2O 70.5% by weight  
 OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD  
 (10 CITINGS)

=> file reg		
COST IN U.S. DOLLARS	SINCE FILE ENTRY	TOTAL SESSION
FULL ESTIMATED COST	180.94	201.07
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	-32.30	-32.30

FILE 'REGISTRY' ENTERED AT 13:48:56 ON 22 APR 2010  
 USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
 PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
 COPYRIGHT (C) 2010 American Chemical Society (ACS)

Property values tagged with IC are from the ZIC/VINITI data file  
 provided by InfoChem.

STRUCTURE FILE UPDATES: 21 APR 2010 HIGHEST RN 1219909-65-5  
 DICTIONARY FILE UPDATES: 21 APR 2010 HIGHEST RN 1219909-65-5

New CAS Information Use Policies, enter HELP USAGETERMS for details.

TSCA INFORMATION NOW CURRENT THROUGH January 8, 2010.

Please note that search-term pricing does apply when  
 conducting SmartSELECT searches.

REGISTRY includes numerically searchable data for experimental and predicted properties as well as tags indicating availability of experimental property data in the original document. For information on property searching in REGISTRY, refer to:

<http://www.cas.org/support/stngen/stdoc/properties.html>

```
=> s fluorouracil/cn
L13      1 FLUOROURACIL/CN

=> d ibib
'IBIB' IS NOT A VALID FORMAT FOR FILE 'REGISTRY'
```

The following are valid formats:

Substance information can be displayed by requesting individual fields or predefined formats. The predefined substance formats are: (RN = CAS Registry Number)

```
REG      - RN
SAM      - Index Name, MF, and structure - no RN
FIDE     - All substance data, except sequence data
IDE      - FIDE, but only 50 names
SQIDE    - IDE, plus sequence data
SQIDE3   - Same as SQIDE, but 3-letter amino acid codes are used
SQD      - Protein sequence data, includes RN
SQD3     - Same as SQD, but 3-letter amino acid codes are used
SQN      - Protein sequence name information, includes RN

EPROP    - Table of experimental properties
PPROP    - Table of predicted properties
PROP     - EPROP, ETAG, PPROP
```

Any CA File format may be combined with any substance format to obtain CA references citing the substance. The substance formats must be cited first. The CA File predefined formats are:

```
ABS -- Abstract
APPS -- Application and Priority Information
BIB -- CA Accession Number, plus Bibliographic Data
CAN -- CA Accession Number
CBIB -- CA Accession Number, plus Bibliographic Data (compressed)
IND -- Index Data
IPC -- International Patent Classification
PATS -- PI, SO
STD -- BIB, IPC, and NCL

IABS -- ABS, indented, with text labels
IBIB -- BIB, indented, with text labels
ISTD -- STD format, indented

OBIB ----- AN, plus Bibliographic Data (original)
OIBIB ----- OBIB, indented with text labels

SBIB ----- BIB, no citations
SIBIB ----- IBIB, no citations
```

The ALL format gives FIDE BIB ABS IND RE, plus sequence data when it is available.

The MAX format is the same as ALL plus SPEC.

The IALL format is the same as ALL with BIB ABS and IND indented,

with text labels.

For additional information, please consult the following help messages:

HELP DFIELDS -- To see a complete list of individual display fields.  
HELP FORMATS -- To see detailed descriptions of the predefined formats.  
ENTER DISPLAY FORMAT (IDE):SAM

L13 ANSWER 1 OF 1 REGISTRY COPYRIGHT 2010 ACS on STN  
IN 2,4(1H,3H)-Pyrimidinedione, 5-fluoro-  
MF C4 H3 F N2 O2  
CI COM



\*\*PROPERTY DATA AVAILABLE IN THE 'PROP' FORMAT\*\*

=> d his

(FILE 'HOME' ENTERED AT 13:27:42 ON 22 APR 2010)

FILE 'CAPLUS' ENTERED AT 13:29:20 ON 22 APR 2010  
S UBIQUINONE/CN

L1 FILE 'REGISTRY' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S UBIQUINONE/CN

L2 FILE 'CAPLUS' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S L1  
S COENZYME Q10/CN

L3 FILE 'REGISTRY' ENTERED AT 13:30:10 ON 22 APR 2010  
1 S COENZYME Q10/CN

L4 FILE 'CAPLUS' ENTERED AT 13:30:10 ON 22 APR 2010  
5610 S L3

L5 FILE 'REGISTRY' ENTERED AT 13:30:20 ON 22 APR 2010  
1 S COENZYME Q10/CN

L6 FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010  
41 S L5 AND CARCINOMA  
L7 16 S L6 AND PY<=2004  
L8 7 S L5 AND CARCINOMA AND TOPICAL  
L9 38 S L5 AND ?CARCINOMA  
L10 8 S L9 AND (TOPICAL OR SURFACE)  
L11 13 S L5 AND CANCER AND (TOPICAL OR SURFACE)  
L12 5 S L11 AND PY<=2004

FILE 'REGISTRY' ENTERED AT 13:48:56 ON 22 APR 2010  
L13 1 S FLUOROURACIL/CN

=> s l13 and l5  
L14 0 L13 AND L5

	SINCE FILE ENTRY	TOTAL SESSION
=> file caplus		
COST IN U.S. DOLLARS	7.59	208.66
FULL ESTIMATED COST		
DISCOUNT AMOUNTS (FOR QUALIFYING ACCOUNTS)	SINCE FILE ENTRY	TOTAL SESSION
CA SUBSCRIBER PRICE	0.00	-32.30

FILE 'CAPLUS' ENTERED AT 13:50:02 ON 22 APR 2010  
USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.  
PLEASE SEE "HELP USAGETERMS" FOR DETAILS.  
COPYRIGHT (C) 2010 AMERICAN CHEMICAL SOCIETY (ACS)

Copyright of the articles to which records in this database refer is held by the publishers listed in the PUBLISHER (PB) field (available for records published or updated in Chemical Abstracts after December 26, 1996), unless otherwise indicated in the original publications. The CA Lexicon is the copyrighted intellectual property of the American Chemical Society and is provided to assist you in searching databases on STN. Any dissemination, distribution, copying, or storing of this information, without the prior written consent of CAS, is strictly prohibited.

FILE COVERS 1907 - 22 Apr 2010 VOL 152 ISS 17  
FILE LAST UPDATED: 21 Apr 2010 (20100421/ED)  
REVISED CLASS FIELDS (/NCL) LAST RELOADED: Feb 2010  
USPTO MANUAL OF CLASSIFICATIONS THESAURUS ISSUE DATE: Feb 2010

CAPLUS now includes complete International Patent Classification (IPC) reclassification data for the first quarter of 2010.

CAS Information Use Policies apply and are available at:

<http://www.cas.org/legal/infopolicy.html>

This file contains CAS Registry Numbers for easy and accurate substance identification.

=> s l13 and l5  
22369 L13  
5610 L5  
L15 27 L13 AND L5

=> d ibib abs total

L15 ANSWER 1 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2009:1557971 CAPLUS  
DOCUMENT NUMBER: 152:135746  
TITLE: Cheminformatics Analysis of Assertions Mined from Literature that Describe Drug-Induced Liver Injury in Different Species  
AUTHOR(S): Fourches, Denis; Barnes, Julie C.; Day, Nicola C.; Bradley, Paul; Reed, Jane Z.; Tropsha, Alexander

CORPORATE SOURCE: Laboratory of Molecular Modeling, Eshelman School of  
Pharmacy, University of North Carolina at Chapel Hill,  
Chapel Hill, NC, 27599, USA  
SOURCE: Chemical Research in Toxicology (2010), 23(1), 171-183  
CODEN: CRTOC; ISSN: 0893-228X  
PUBLISHER: American Chemical Society  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Drug-induced liver injury is one of the main causes of drug attrition. The ability to predict the liver effects of drug candidates from their chemical structures is critical to help guide exptl. drug discovery projects toward safer medicines. In this study, the authors have compiled a data set of 951 compds. reported to produce a wide range of effects in the liver in different species, comprising humans, rodents, and nonrodents. The liver effects for this data set were obtained as assertional metadata, generated from MEDLINE abstrs. using a unique combination of lexical and linguistic methods and ontol. rules. The authors have analyzed this data set using conventional cheminformatics approaches and addressed several questions pertaining to cross-species concordance of liver effects, chemical determinants of liver effects in humans, and the prediction of whether a given compound is likely to cause a liver effect in humans. The authors found that the concordance of liver effects was relatively low (.apprx.39-44%) between different species, raising the possibility that species specificity could depend on specific features of chemical structure. Compds. were clustered by their chemical similarity, and similar compds. were examined for the expected similarity of their species-dependent liver effect profiles. In most cases, similar profiles were observed for members of the same cluster, but some compds. appeared as outliers. The outliers were the subject of focused assertion regeneration from MEDLINE as well as other data sources. In some cases, addnl. biol. assertions were identified, which were in line with expectations based on compds.' chemical similarities. The assertions were further converted to binary annotations of underlying chems. (i.e., liver effect vs. no liver effect), and binary quant. structure-activity relationship (QSAR) models were generated to predict whether a compound would be expected to produce liver effects in humans. Despite the apparent heterogeneity of data, models have shown good predictive power assessed by external 5-fold cross-validation procedures. The external predictive power of binary QSAR models was further confirmed by their application to compds. that were retrieved or studied after the model was developed. To the best of the authors' knowledge, this is the first study for chemical toxicity prediction that applied QSAR modeling and other cheminformatics techniques to observational data generated by the means of automated text mining with limited manual curation, opening up new opportunities for generating and modeling chemical toxicol. data.

REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 2 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2009:1368423 CAPLUS  
DOCUMENT NUMBER: 152:51216  
TITLE: Drug Effects Viewed from a Signal Transduction Network  
Perspective  
AUTHOR(S): Fliri, Anton F.; Logging, William T.; Volkmann, Robert  
A.  
CORPORATE SOURCE: Pfizer Global Research and Development, Groton, CT,  
06340, USA  
SOURCE: Journal of Medicinal Chemistry (2009), 52(24),  
8038-8046  
CODEN: JMCMAR; ISSN: 0022-2623  
PUBLISHER: American Chemical Society



DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Understanding how drugs affect cellular network structures and how resulting signals are translated into drug effects holds the key to the discovery of medicines. Herein we examine this cause-effect relationship by determining protein network structures associated with the generation of specific in vivo drug-effect patterns. Medicines having similar in vivo pharmacol. have been identified by a comparison of drug-effect profiles of 1320 medicines. Protein network positions reached by these medicines were ascertained by examining the coinvestigation frequency of these medicines and 1179 protein network constituents in millions of scientific investigations. Interestingly, medicine assocns. obtained by comparing by drug-effect profiles mirror those obtained by comparing drug-protein coinvestigation frequency profiles, demonstrating that these drug-protein reachability profiles are relevant to in vivo pharmacol. By using protein assocns. obtained in these investigations and independent, curated protein interaction information, drug-mediated protein network topol. models can be constructed. These protein network topol. models reveal that drugs having similar pharmacol. profiles reach similar discrete positions in cellular protein network systems and provide a network view of medicine cause-effect relationships.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 3 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:1398483 CAPLUS

DOCUMENT NUMBER: 149:570734

TITLE: Ghrelin modulating compounds and combinations thereof

INVENTOR(S): Watson, Alan; Distefano, Peter; Geesaman, Bard J.

PATENT ASSIGNEE(S): Elixir Pharmaceuticals, Inc., USA

SOURCE: PCT Int. Appl., 182pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2008141189	A1	20081120	WO 2008-US63257	20080509
W:	AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, NO, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			

PRIORITY APPLN. INFO.: US 2007-917054P P 20070509

OTHER SOURCE(S): MARPAT 149:570734

AB Comps. that modulate GHS-R are disclosed here.

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 4 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2008:526300 CAPLUS

DOCUMENT NUMBER: 148:456694

TITLE: Hybrid lipid-polymer nanoparticulate drug delivery composition  
 INVENTOR(S): Gao, Hai Yan; Schwarz, Joseph; Weisspapir, Michael  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 14pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080102127	A1	20080501	US 2007-848484	20070831
PRIORITY APPLN. INFO.:			US 2006-854458P	P 20061026

AB The invention relates to a nanoparticulate colloidal delivery vehicle comprising a biodegradable polymer in combination with a hydrophobic lipid component. Variation of the lipid and polymer types and variation in the ratio between the polymer and lipid components allows regulation of drug loading and release rate. Thus, drug-loaded hybrid lipid-polymer nanoparticles comprised: streptomycin sulfate 100 mg, PLGA 750 mg, cholesterol 100 mg, cholesteryl sulfate sodium salt 25 mg, Cremophor EL 2%, prepared in 24 mL Et acetate by emulsification; yield 86%; drug binding 69.3%; size 223 nm.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L15 ANSWER 5 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:1215784 CAPLUS

DOCUMENT NUMBER: 147:491621

TITLE: Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer

INVENTOR(S): Mazzio, Elizabeth; Soliman, Karam

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S. Ser. No. 233,279.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070248693	A1	20071025	US 2007-711883	20070227
US 20060035981	A1	20060216	US 2005-233279	20050920
PRIORITY APPLN. INFO.:			US 2003-491841P	P 20030802
			US 2004-540525P	P 20040129
			US 2004-909590	B2 20040802
			US 2005-233279	A2 20050920

AB The invention describes a pharmaceutical composition and method for treating cancer comprising (a) 2,3-dimethoxy-5-methyl-1,4-benzoquinone, and/or (b) at least one of wild yam root, teasel root, balm of gilead bud, bakuchi seed, dichroa root, kochia seed, kanta kari, bushy knotweed rhizome, arjun, babul chall bark, opopanax and bhumi amalaki; optionally one or more of frankincense, garcinia fruit, vitex, dragons blood, mace, sage and red sandalwood with at least (c) one compound capable of maximizing oxidative mitochondrial function, preferably riboflavin or vitamin B2 derivs., FAD, FMN, 5-amino-6-(5'-phosphoribitylamino)uracil, 6,7-dimethyl-8-(1-D-ribityl)lumazine, ribitol, 5,6-dimethylbenzimidazole,

tetrahydrobiopterin, vitamin B1, lipoic acid, biotin, vitamin B6, vitamin B12, folate, niacin, vitamin C and pantothenate, and/or (d) at least one lactic acid dehydrogenase inhibitor, preferably 2',3,4',5,7-pentahydroxyflavone and optionally (f) an alkalinizing agent (Aloe vera, chlorella, wheat grass, sodium or potassium bicarbonate, potassium), (g) an antiproliferative herb (speranskia or goldenseal), and (h) a pharmaceutically acceptable carrier. A method for inhibiting cancer optionally comprises one or more chemotherapy drug(s), selected, among others, from acetogenins, actinomycin D, adriamycin, aminoglutethimide, asparaginase, bleomycin, bullatacin, busulfan, carmustine, carboplatin, chlorambucil, cisplatin, etc. Thus, a composition comprised rosemary (Rosmarinus officinalis) .apprx.1000, myrrh gum (Commiphora molmol) .apprx.500, 2,3-dimethoxy-5-methyl-1,4 benzoquinone .apprx.800, and riboflavin .apprx.300 mg/day, resp.

L15 ANSWER 6 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:113642 CAPLUS

DOCUMENT NUMBER: 146:198725

TITLE: Proteasome inhibitors and other small compounds that correct protein misfolding and uses thereof

INVENTOR(S): Kaushal, Shalesh; Noorwez, Syed Mohammed

PATENT ASSIGNEE(S): University of Florida Research Foundation, Inc., USA

SOURCE: PCT Int. Appl., 60 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2007014327	A2	20070201	WO 2006-US29402	20060727
WO 2007014327	A3	20090430		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
AU 2006272497	A1	20070201	AU 2006-272497	20060727
CA 2616537	A1	20070201	CA 2006-2616537	20060727
EP 1909812	A2	20080416	EP 2006-800453	20060727
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, RS			
JP 2009502954	T	20090129	JP 2008-524192	20060727
MX 2008001130	A	20080529	MX 2008-1130	20080124
IN 2008DN01290	A	20090320	IN 2008-DN1290	20080214
KR 2008033463	A	20080416	KR 2008-704687	20080227
CN 101600475	A	20091209	CN 2006-80035439	20080326
US 20100004156	A1	20100107	US 2009-989356	20090805
PRIORITY APPLN. INFO.:			US 2005-703068P	P 20050727
			WO 2006-US29402	W 20060727

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention discloses compns. and methods that are useful for treating

or preventing a protein conformation disease in a subject by correcting misfolded proteins in vivo. In addition, the invention provides compns. and methods that are useful for expressing a recombinant protein in a biochem. functional conformation.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L15 ANSWER 7 OF 27 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 2006:1155706 CAPLUS  
DOCUMENT NUMBER: 145:465766  
TITLE: Materials and methods for enhanced degradation of mutant proteins associated with human disease  
INVENTOR(S): Kaushal, Shalesh; Malhotra, Ritu; Dunn, William A.  
PATENT ASSIGNEE(S): University of Florida, USA  
SOURCE: PCT Int. Appl., 87pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006116716	A2	20061102	WO 2006-US16368	20060427
WO 2006116716	A3	20070510		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA				
AU 2006239219	A1	20061102	AU 2006-239219	20060427
AU 2006239219	A2	20061102		
CA 2606226	A1	20061102	CA 2006-2606226	20060427
EP 1874118	A2	20080109	EP 2006-751856	20060427
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR				
JP 2008539276	T	20081113	JP 2008-509194	20060427
MX 2007013324	A	20080529	MX 2007-13324	20071025
IN 2007DN08513	A	20080704	IN 2007-DN8513	20071105
KR 2008018874	A	20080228	KR 2007-727659	20071127
CN 101287370	A	20081015	CN 2006-80023254	20071227
US 20100087474	A1	20100408	US 2009-919371	20091209
PRIORITY APPLN. INFO.:				
			US 2005-675143P	P 20050427
			US 2005-723288P	P 20051003
			WO 2006-US16368	W 20060427

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention features compns. and methods that are useful for treating or preventing a protein conformation disease in a subject by enhancing the degradation of misfolded proteins in vivo.

L15 ANSWER 8 OF 27 CAPLUS COPYRIGHT 2010 ACS ON STN

ACCESSION NUMBER: 2006:825000 CAPLUS  
DOCUMENT NUMBER: 145:412897  
TITLE: Biochemical characterization of recombinant

dihydroorotate dehydrogenase from the opportunistic pathogenic yeast *Candida albicans*  
 AUTHOR(S): Zameitat, Elke; Gojkovic, Zoran; Knecht, Wolfgang; Piskur, Jure; Loeffler, Monika  
 CORPORATE SOURCE: Institute for Physiological Chemistry, Philipps-University, Marburg, Germany  
 SOURCE: FEBS Journal (2006), 273(14), 3183-3191  
 CODEN: FJEOAC; ISSN: 1742-464X  
 PUBLISHER: Blackwell Publishing Ltd.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB *Candida albicans* is the most prevalent yeast pathogen in humans, and recently it has become increasingly resistant to the current antifungal agents. In this study we investigated *C. albicans* dihydroorotate dehydrogenase (DHODH, EC 1.3.99.11), which catalyzes the fourth step of de novo pyrimidine synthesis, as a new target for controlling infection. We propose that the enzyme is a member of the DHODH family 2, which comprises mitochondrially bound enzymes, with quinone as the direct electron acceptor and oxygen as the final electron acceptor. Full-length DHODH and N-terminally truncated DHODH, which lacks the targeting sequence and the transmembrane domain, were subcloned from *C. albicans*, recombinantly expressed in *Escherichia coli*, purified, and characterized for their kinetics and substrate specificity. An inhibitor screening with 28 selected compounds was performed. Only the dianisidine derivative, redoxal, and the biphenyl quinoline-carboxylic acid derivative, brequnar sodium, which are known to be potent inhibitors of mammalian DHODH, markedly reduced *C. albicans* DHODH activity. This study provides a background for the development of antiprimidines with high efficacy for decreasing *in situ* pyrimidine nucleotide pools in *C. albicans*.  
 OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
 REFERENCE COUNT: 38 THERE ARE 38 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 9 OF 27 CAPLUS COPYRIGHT 2010 ACS ON STN  
 ACCESSION NUMBER: 2006:666025 CAPLUS  
 DOCUMENT NUMBER: 145:152690  
 TITLE: Method for inducing crystalline state transition in pharmaceuticals  
 INVENTOR(S): Nakamichi, Kouichi; Izumi, Shougo; Oka, Masaaki  
 PATENT ASSIGNEE(S): Nippon Shinyaju Company, Ltd., Japan  
 SOURCE: U.S., 18 pp., Cont.-in-part of U. S. 5,456,923.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5811547	A	19980922	US 1995-416815	19950609
CA 2147279	A1	19940428	CA 1993-2147279	19931013
WO 9408561	A1	19940428	WO 1993-JP1469	19931013
W: AU, BR, CA, FI, HU, JP, KR, NO, NZ, RU, US				
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9351607	A	19940509	AU 1993-51607	19931013
EP 665009	A1	19950802	EP 1993-922625	19931013
EP 665009	B1	20000216		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE				
AT 189770	T	20000315	AT 1993-922625	19931013
ES 2145063	T3	20000701	ES 1993-922625	19931013

US 5456923 A 19951010 US 1993-129133 19931115  
 PRIORITY APPLN. INFO.: JP 1992-303085 A 19921014  
 WO 1993-JP1469 W 19931013  
 US 1993-129133 A2 19931115  
 JP 1991-112554 A 19910416  
 WO 1992-JP470 W 19920414

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB This invention has for its object to provide a method of inducing a transition in crystalline state of a crystallizable pharmaceutical with great ease and improved efficiency and uniformity on a high production scale. An extruder is used for inducing a transition from one crystalline state ( $\Delta$ ) to another crystalline state in a crystallizable pharmaceutical. An extruded indomethacin (form  $\alpha$ ) was converted to an amorphous form.

OS.CITING REF COUNT: 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 10 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2006:437475 CAPLUS

DOCUMENT NUMBER: 144:460856

TITLE: Methods and compositions using a bile acid and a carbohydrate for reducing neurodegeneration in amyotrophic lateral sclerosis or other neurodegenerative disease

INVENTOR(S): Yoo, Seo Hong

PATENT ASSIGNEE(S): USA

SOURCE: PCT Int. Appl., 64 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2006050165	A2	20060511	WO 2005-US39089	20051031
WO 2006050165	A3	20060706		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
AU 2005302452	A1	20060511	AU 2005-302452	20051031
CA 2585471	A1	20060511	CA 2005-2585471	20051031
US 20060142241	A1	20060629	US 2005-263087	20051031
EP 1814558	A2	20070808	EP 2005-820886	20051031
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR				
CN 101048164	A	20071003	CN 2005-80037307	20051031
JP 2008518935	T	20080605	JP 2007-539200	20051031
KR 2007089926	A	20070904	KR 2007-712360	20070531
IN 2007KN01990	A	20070810	IN 2007-KN1990	20070604
PRIORITY APPLN. INFO.:			US 2004-624100P	P 20041101

US 2004-628421P P 20041116  
 WO 2005-US39089 W 20051031

AB The invention discloses clear aqueous solns. of one or more bile acids and either an aqueous soluble starch conversion product or a non-starch polysaccharide. The solns. may be administered to a subject in conjunction with a pharmaceutical compound having a therapeutic effect in subjects with a neurodegenerative disease and/or a motor neuron disease. In some embodiments, the disease is amyotrophic lateral sclerosis.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 11 OF 2/ CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2006:234115 CAPLUS

DOCUMENT NUMBER: 144:298855

TITLE: Compositions for treatment of skin discoloration

INVENTOR(S): Boxrud, Cynthia A.

PATENT ASSIGNEE(S): Evera Laboratories, LLC, USA

SOURCE: U.S. Pat. Appl. Publ., 12 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20060057081	A1	20060316	US 2005-37589	20050118
US 7288263	B2	20071030		
WO 2006031555	A2	20060323	WO 2005-US31822	20050907
WO 2006031555	A3	20071011		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA			
EP 1804761	A2	20070711	EP 2005-795962	20050907
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU			
JP 2008512469	T	20080424	JP 2007-531296	20050907
PRIORITY APPLN. INFO.:			US 2004-609543P	P 20040913
			US 2005-37589	A 20050118
			WO 2005-US31822	W 20050907



ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A cosmetically acceptable product for application to human skin is disclosed. The novel comps. are particularly suited for skin lightening and for diminishing the appearance of dark circles under the eyes. The comps. include any of several vasoconstrictors in a carrier with optionally added skin compatible ingredients. Thus, a formulation contained tetrahydrozoline-HCl 2.00, dimethicone 1.00, triethanolamine 1.00, phenoxyethanol 0.50, Carbomer 0.50, methylparaben 0.20, isopropylparaben 0.10, propylparaben 0.10, isobutylparaben 0.02,

butylparaben, and water 94.56%.  
 REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 12 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2005:904349 CAPLUS  
 DOCUMENT NUMBER: 143:248278  
 TITLE: Preparation of sulfonylpyrrolidines as modulators of  
 androgen receptor  
 INVENTOR(S): Hamann, Lawrence G.; Bi, Yingzhi; Manfredi, Mark C.;  
 Nirschl, Alexandra A.; Sutton, James C.  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 35 pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20050187267	A1	20050825	US 2005-48439	20050201
PRIORITY APPLN. INFO.:			US 2004-541869P	P 20040204
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT				
OTHER SOURCE(S):			CASREACT 143:248278; MARPAT 143:248278	
GI				



AB Title compds. I or II [R1 = H, (un)substituted alkyl, alkenyl, etc.; R2 = H, halo, SR6, etc.; R3 and R4 independently = H, (un)substituted alkynyl, cycloalkyl, etc.; R5 = H, (un)substituted aryl, arylalkyl, etc.; R6 = H, CHF2, CF3, etc.; X = (CH2)n; G = (un)substituted aryl, heterocycle or heteroaryl; Z = O or NR7; R7 = H, (un)substituted alkyl, alkenyl, etc.; n and m independently = 1-2] and their pharmaceutically acceptable salts,



are prepared and disclosed as modulators of androgen receptor. Thus, e.g., III was prepared by hydrolysis of (2S,3R)-1-(3-chloro-4-cyano-2-methyl-phenylsulfamoyl)-3-hydroxy-pyrrolidine-2-carboxylic acid Me ester (preparation given) followed by cyclization. The activity of I was evaluated in transactivation assays of a transfected reporter construct and using the endogenous androgen receptor of the host cells (no data). I as modulator of androgen receptor should prove useful in the treatment of neoplasm, Alzheimer's disease and obesity. Pharmaceutical compns. comprising I are disclosed.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L15 ANSWER 13 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:902874 CAPLUS

DOCUMENT NUMBER: 143:248277

TITLE: Preparation of sulfonylpyrrolidines as modulators of androgen receptor

INVENTOR(S): Hamann, Lawrence H.; Bi, Yingzhi; Manfredi, Mark C.;

Nirschl, Alexandra A.; Sutton, James C.

PATENT ASSIGNEE(S): Bristol-Myers Squibb Company, USA

SOURCE: PCT Int. Appl., 91 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2




PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005077925	A1	20050825	WO 2005-US2834	20050202
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1718626	A1	20061108	EP 2005-712320	20050202
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, TR, BG, CZ, EE, HU, PL, SK, HR, IS, YU			

PRIORITY APPLN. INFO.: US 2004-541869P P 20040204  
WO 2005-US2834 W 20050202

OTHER SOURCE(S): CASREACT 143:248277; MARPAT 143:248277

GI



AB Title compds. I or II [R1 = H, (un)substituted alkyl, alkenyl, etc.; R2 = H, halo, SR6, etc.; R3 and R4 independently = H, (un)substituted alkynyl, cycloalkyl, etc.; R5 = H, (un)substituted aryl, arylalkyl, etc.; R6 = H, CHF2, CF3, etc.; X = (CH2)n; G = (un)substituted aryl, heterocycle or heteroaryl; Z = O or NR7; R7 = H, (un)substituted alkyl, alkenyl, etc.; n and m independently = 1-2] and their pharmaceutically acceptable salts, are prepared and disclosed as modulators of androgen receptor. Thus, e.g., III was prepared by hydrolysis of (2S,3R)-1-(3-chloro-4-cyano-2-methyl-phenylsulfamoyl)-3-hydroxy-pyrrolidine-2-carboxylic acid Me ester (preparation given) followed by cyclization. The activity of I was evaluated in transactivation assays of a transfected reporter construct and using the endogenous androgen receptor of the host cells (no data). I as modulator of androgen receptor should prove useful in the treatment of neoplasm, Alzheimer's disease and obesity. Pharmaceutical compns. comprising I are disclosed.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 14 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:824492 CAPLUS

DOCUMENT NUMBER: 143:222525

TITLE: Method of using 3-cyano-4-arylpyridine derivatives as modulators of androgen receptor function, preparation thereof, and use with other agents

INVENTOR(S): Nirschl, Alexandra A.; Hamann, Lawrence G.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 25 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
------------	------	------	-----------------	------

-----  
 US 20050182105 A1 20050818 US 2005-48437 20050201  
 PRIORITY APPLN. INFO.: US 2004-541780P P 20040204  
 ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
 OTHER SOURCE(S): MARPAT 143:222525  
 GI



AB A method is provided for treating androgen receptor-associated conditions, such as age-related diseases, e.g. sarcopenia, employing a compound I [R1 = CN, H; X = O, S; R2 = (substituted) alkyl, (substituted) cycloalkyl, etc; R3, R4 = H, (substituted) alkyl, etc.; G = (substituted) aryl, (substituted) heteroaryl], or a pharmaceutically acceptable salt or prodrug ester thereof. Preparation of selected I is described. I may be used in combination with other agents.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L15 ANSWER 15 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:334805 CAPLUS

DOCUMENT NUMBER: 138:348669

TITLE: Lymphocyte assay-based methods for determining toxicity-reversing agents

INVENTOR(S): Shive, William; Pettit, Flora H.

PATENT ASSIGNEE(S): Board of Regents, The University of Texas System, USA; Shive, Gwyn

SOURCE: PCT Int. Appl., 46 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003034803	A2	20030501	WO 2002-IB5160	20021018
WO 2003034803	A3	20031002		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG				
US 20030087331	A1	20030508	US 2001-10716	20011026
US 6723527	B2	20040420		
AU 2002363105	A1	20030506	AU 2002-363105	20021018

PRIORITY APPLN. INFO.: US 2001-10716 A1 20011026  
WO 2002-IB5160 W 20021018

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Methods are disclosed for assessment of the ability of substances to ameliorate the toxic effects of compds. based on a lymphocyte culture assay. The lymphocyte assay is a repeatable and quant. assay for lymphocyte growth in a chemical defined media in which specific compds. with potential toxicity and substances with potential abilities to ameliorate the toxicity can be added to determine specific and individualized requirements for such substances. Also disclosed are methods for ameliorating side-effects by administering a substance identified by the methods of the invention to a patient undergoing therapy with a drug that has a toxic effect. Further provided is a composition that ameliorates the toxic-effect of the statin family of drugs. Methods and processes for partially purifying and/or isolating this composition from plasma are also provided. Thus, the methods of the invention not only provide substances for reversal of compound toxicity but also provide methods for pre-approving compds. for human use.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 16 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:261643 CAPLUS

DOCUMENT NUMBER: 138:260506

TITLE: Granules having improved dosing properties

INVENTOR(S): Murai, Kouji; Uchida, Akihiro; Aimoto, Masaharu; Kato, Yasuki

PATENT ASSIGNEE(S): Kyowa Hakko Kogyo Co., Ltd., Japan

SOURCE: PCT Int. Appl., 23 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: Japanese

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003026619	A1	20030403	WO 2002-JP9910	20020926
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2002338097	A1	20030407	AU 2002-338097	20020926
PRIORITY APPLN. INFO.:			JP 2001-295143	A 20010926
			WO 2002-JP9910	W 20020926

AB It is intended to provide granules having relieved coarseness in the oral cavity in dosing, characterized by containing an active ingredient which is hardly soluble in water or saliva and a component which is converted into a viscous liquid upon the addition of water. Oxatamide 2, hydroxypropyl Me cellulose 0.5, hydroxypropyl starch 5.5, and mannitol 91.5 g were mixed and kneaded in 15 mL water. The mixture was granulated and dried.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 17 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2001:730533 CAPLUS  
 DOCUMENT NUMBER: 135:262281  
 TITLE: Water-soluble additives for the manufacture of easy-to-take granules  
 INVENTOR(S): Murai, Kouji; Narita, Shoichi; Ogasa, Takehiro  
 PATENT ASSIGNEE(S): Kyowa Hakko Kogyo Co., Ltd., Japan  
 SOURCE: PCT Int. Appl., 20 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001072285	A1	20011004	WO 2001-JP2406	20010326
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
AU 2001042783	A	20011008	AU 2001-42783	20010326
CA 2403594	A1	20020918	CA 2001-2403594	20010326
EP 1269995	A1	20030102	EP 2001-915776	20010326
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 20030104066	A1	20030605	US 2002-239751	20021029
PRIORITY APPLN. INFO.:			JP 2000-86516	A 20000327
			WO 2001-JP2406	W 20010326

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Disclosed are easy-to-take granules which comprise an active ingredient, at least one soluble additive having an average particle diameter smaller than 50

µm, and at least one disintegrator. The granules are easily dissolved or disintegrated in the buccal cavity. D-Mannitol 90 g was pulverized and mixed with croscopovidone 5.5, hydroxypropyl cellulose 2, and oxatamide 2 g. Water was added to the mixture for kneading and granulation.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 18 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1998:489534 CAPLUS  
 DOCUMENT NUMBER: 129:293760  
 ORIGINAL REFERENCE NO.: 129:59843a, 59846a  
 TITLE: Percutaneous absorption of one hundred drugs and the derivation of an experimental regression equation  
 AUTHOR(S): Xu, Jingfeng; Zhao, Weijuan; Zhang, Mei; Liu, Mei; Wang, Jinping; Jin, Yinghua; Wang, Yurong  
 CORPORATE SOURCE: Beijing Military Command Clinical Pharmaceutical Institute, Beijing, 100700, Peop. Rep. China  
 SOURCE: Zhongguo Yaoxue Zazhi (Beijing) (1998), 33(2), 86-91

PUBLISHER: Zhongguo Yaoxuehui  
DOCUMENT TYPE: Journal  
LANGUAGE: Chinese

AB The pharmaceutical regularity of percutaneous absorption was studied. The percutaneous absorption speed of 100 drugs and the comparison with the permeation enhancer of 2% and 5% Azone were studied in mouse with an improved Fick's diffusion installation by computing accumulative permeation quantity (Q), steady percutaneous speed (J), and permeation coefficient (Kp). The rules in pharmaceuticals of drug's phys. and chemical characteristics and percutaneous absorption were discussed, and the exptl. regression equation of drug percutaneous absorption were calculated and the influence of different concns. of azone on drug percutaneous permeation and equation were studied.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L15 ANSWER 19 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1994:200438 CAPLUS

DOCUMENT NUMBER: 120:200438

ORIGINAL REFERENCE NO.: 120:35325a,35328a

TITLE: Controlled-release transdermal pharmaceuticals containing cryogels

INVENTOR(S): Wood, Louis L.; Calton, Gary J.

PATENT ASSIGNEE(S): SRCHEM Inc., USA

SOURCE: U.S., 15 pp.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 5260066	A	19931109	US 1992-821627	19920116
US 5288503	A	19940222	US 1992-899369	19920616
			US 1992-821627	A3 19920116

PRIORITY APPLN. INFO.:

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A controlled-release transdermal pharmaceutical containing therapeutic agents in a poly(vinyl alc.) (I) cryogel is disclosed. A slurry of 11.0 mg ciprofloxacin.HCl (II) and 200 mg 10% I was warmed to 50-60° to obtain a clear homogeneous solution. The solution was then placed in a mold and subjected to 6 freeze-thaw cycles to give a white opaque elastomeric cryogel having 15mm diameter and 0.5mm thickness. The release of II from the gel in 0.9% NaCl was 74% in the 1st 4 hs and it was constant in the subsequent 5-24 hs.

OS.CITING REF COUNT: 40 THERE ARE 40 CAPLUS RECORDS THAT CITE THIS RECORD (43 CITINGS)

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L15 ANSWER 20 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1985:119628 CAPLUS

Correction of: 1984:478825

DOCUMENT NUMBER: 102:119628

Correction of: 101:78825

ORIGINAL REFERENCE NO.: 102:18731a,18734a

TITLE: Drug solubilization by tri-O-methyl- $\beta$ -cyclodextrin

PATENT ASSIGNEE(S): Zeria Pharmaceutical Co., Ltd., Japan

SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.

DOCUMENT TYPE: CODEN: JKXXAF  
 LANGUAGE: Patent  
 FAMILY ACC. NUM. COUNT: Japanese  
 PATENT INFORMATION: 1

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59046228	A	19840315	JP 1982-156424	19820908
PRIORITY APPLN. INFO.:			JP 1982-156424	19820908

AB Insol. drugs are solubilized by tri-O-methyl- $\beta$ -cyclodextrin [74948-17-7]. These drugs include neoplasm inhibitors, inflammation inhibitors, etc. Thus, 71.3 g tri-O-methyl- $\beta$ -cyclodextrin was dissolved in 100 mL H<sub>2</sub>O, and flurbiprofen (I) [5104-49-4] was added. The maximum concentration of I was 0.896 g/100 mL, whereas that of I dissolved in H<sub>2</sub>O alone was 0.059 g/100 mL.

L15 ANSWER 21 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1984:478825 CAPLUS  
 DOCUMENT NUMBER: 101:78825  
 ORIGINAL REFERENCE NO.: 101:12051a,12054a  
 TITLE: Drug solubilization by tri-O-methyl- $\beta$ -cyclodextrin  
 PATENT ASSIGNEE(S): Zeria Shinyaku Kogyo K. K., Japan  
 SOURCE: Jpn. Kokai Tokkyo Koho, 9 pp.  
 CODEN: JKXXAF  
 DOCUMENT TYPE: Patent  
 LANGUAGE: Japanese  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
JP 59046228 A		19840315	JP 1982-156424	19820908

AB Insol. drugs are solubilized by tri-O-methyl- $\beta$ -cyclodextrin [55216-11-0]. These drugs include neoplasm inhibitors, inflammation inhibitors, etc. Thus, 71.3 g tri-O-methyl- $\beta$ -cyclodextrin was dissolved in 100 mL H<sub>2</sub>O, and flurbiprofen (I) [5104-49-4] was added. The maximum concentration of I was 0.896 g/100 mL, whereas that of I dissolved in H<sub>2</sub>O alone was 0.059 g/100 mL.

L15 ANSWER 22 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1983:587238 CAPLUS  
 DOCUMENT NUMBER: 99:187238  
 ORIGINAL REFERENCE NO.: 99:28595a,28598a  
 TITLE: Cardiotoxicogenetic mechanism of 5-fluorouracil (5-FU)  
 AUTHOR(S): Saso, Fumio  
 CORPORATE SOURCE: Sch. Med., Jikei Univ., Tokyo, 105, Japan  
 SOURCE: Tokyo Jikeikai Ika Daigaku Zasshi (1983), 98(2), 186-204  
 CODEN: TJIDAH; ISSN: 0375-9172  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Japanese  
 GI



I

AB The effects of 5-fluorouracil (I) [51-21-8] and N1-(2-tetrahydrofuryl)-5-fluorouracil (II) [17902-23-7] on cardiac function and myocardial metabolism were compared in isolated and perfused rat hearts. I (10-4 g/mL) suppressed heart rate, peak systolic pressure, and coronary flow. I lowered the tissue levels of ATP [56-65-5]. Coenzyme Q10 [303-98-0] significantly alleviated I-induced suppression of cardiac function. The actions of II (10-3 g/mL) were similar but weaker than to those of I.

L15 ANSWER 23 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1981:454780 CAPLUS

DOCUMENT NUMBER: 95:54780

ORIGINAL REFERENCE NO.: 95:9141a,9144a

TITLE: Liver cell injury by antineoplastic agents and the influence of coenzyme Q10 on the cellular K+ and membrane potential difference (PD) in the rat

AUTHOR(S): Okada, Katsuhiko; Kitade, Fumio; Yamada, Shinichi; Kawashima, Yasuo; Okajima, Kunio; Fujimoto, Mamoru

CORPORATE SOURCE: Dep. Surg., Osaka Med. Coll., Osaka, Japan


SOURCE: Nippon Shokakibyo Gakkai Zasshi (1979), 76(4), 896-904

CODEN: NIPAA4; ISSN: 0369-4259

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

GI



I



II

AB Both the membrane PD and intracellular K+ concentration were decreased by administration of antineoplastic agents (mitomycin C (I) [50-07-7] or 5-fluorouracil (II) [51-21-8]). Both I and II seem to elicit a hypofunction of hepatic cells as a result of their side effects, being characterized by a decrease of cellular energy metabolism, cellular K+ accumulation, and Na+ transport out of the cell. Administration of coenzyme Q10 [303-98-0] was recognized to partially reverse these side effects.

L15 ANSWER 24 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN


ACCESSION NUMBER: 1981:150213 CAPLUS

DOCUMENT NUMBER: 94:150213

ORIGINAL REFERENCE NO.: 94:24431a,24434a




TITLE: Coenzyme Q10 treatment for the liver damages induced by antineoplastic agents  
 AUTHOR(S): Yamada, Shinichi  
 CORPORATE SOURCE: Dep. Surg., Osaka Med. Coll., Osaka, Japan  
 SOURCE: Nippon Gan Chiryo Gakkaishi (1981), 15(6), 1003-15  
 CODEN: NGCJAK; ISSN: 0021-4671  
 DOCUMENT TYPE: Journal  
 LANGUAGE: Japanese  
 GI



AB Simultaneous administration of coenzyme Q10 (I) [303-98-0] with either mitomycin C (II) [50-07-7] or 5-fluorouracil (III) [51-21-8] to rats bearing AH-130 tumor decreased the side effects of these antitumor agents without decreasing the antitumor activity. I prevented the swelling of the liver cells and the decrease of cellular energy metabolism

L15 ANSWER 25 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1980:597636 CAPLUS  
 DOCUMENT NUMBER: 93:197636  
 ORIGINAL REFERENCE NO.: 93:31355a,31358a  
 TITLE: Injury of rat liver cells by antineoplastic agents and preventive effects of coenzyme Q10  
 AUTHOR(S): Okada, Katsuhiko; Kitade, Fumio; Yamada, Shinichi; Kawashima, Yasuo; Okajima, Kunio; Fujimoto, Mamoru  
 CORPORATE SOURCE: Dep. Surg., Osaka Med. Coll., Osaka, Japan  
 SOURCE: Biomed. Clin. Aspects Coenzyme Q, Proc. Int. Symp., 2nd (1980), Meeting Date 1979, 159-77. Editor(s): Yamamura, Yuichi; Folkers, Karl August; Ito, Y.  
 CODEN: 44IYAO  
 DOCUMENT TYPE: Conference  
 LANGUAGE: English  
 GI



I

AB Membrane p.d. and intracellular activity of the K ion (aK) in rat liver cells were measured simultaneously using double-barreled potassium ion-selective microelectrodes. Both PD and aK in liver cells were depressed after treatment with antineoplastic agents (5-fluorouracil [51-21-8] and mitomycin C [50-07-7]), suggesting that these drugs would induce disturbances of cellular energy metabolism in liver cell. When the antineoplastic agents were used in combination with coenzyme Q10 (I) [303-98-0], the depression of PD and aK and the hypofunction of liver cells in energy metabolism were significantly prevented.

L15 ANSWER 26 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1980:437238 CAPLUS

DOCUMENT NUMBER: 93:37238


ORIGINAL REFERENCE NO.: 93:6021a,6024a

TITLE: Cell injury by antineoplastic agents and influence of coenzyme Q10 on cellular potassium activity and potential difference across the membrane in rat liver cells

AUTHOR(S): Okada, Katsuhiko; Yamada, Shinichi; Kawashima, Yasuo; Kitade, Fumio; Okajima, Kunio; Fujimoto, Mamoru  
CORPORATE SOURCE: Dep. Surg., Osaka Med. Coll., Osaka, 569, Japan  
SOURCE: Cancer Research (1980), 40(5), 1663-7  
CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE: Journal  
LANGUAGE: English

GI




I

AB The p.d. across the cell membrane and the intracellular activity of the K ion in rat liver cells were measured simultaneously using double-barreled K selective microelectrodes. Both the p.d. across the membrane and K activity in liver cells were depressed after treatment with the antineoplastic agents mitomycin C [50-07-7] and 5-Fluorouracil Dry Syrup [51-21-8], suggesting that these drugs would induce disturbances of cellular energy metabolism in liver cells. When the antineoplastic agents were used in combination with coenzyme Q10 (I) [303-98-0], the depression of p.d. across the membrane and K activity and the hypofunction of liver cells in energy metabolism were prevented.

L15 ANSWER 27 OF 27 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1979:48372 CAPLUS

DOCUMENT NUMBER: 90:48372  
 ORIGINAL REFERENCE NO.: 90:7637a,7640a  
 TITLE: Experimental results with the combination of bleomycin plus mitomycin C  
 AUTHOR(S): Yamanaka, N.; Fukushima, M.; Kato, T.; Koizumi, K.; Ota, K.  
 CORPORATE SOURCE: Lab. Chemother., Aichi Cancer Cent. Res. Inst., Nagoya, Japan  
 SOURCE: Recent Results in Cancer Research (1978), 63(Antitumor Antibiot.), 211-18  
 CODEN: RRCRBU; ISSN: 0080-0015  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



II

AB Bleomycin (I) [11056-06-7] was more effective in killing KB cancer cells in culture when combined with mitomycin C (II) [50-07-7] or other quinone-containing anticancer agents or oxidizing and reducing vitamins such as vitamin C [50-81-7] and vitamin K2 [11032-49-8]. In AH66 tumor-bearing rats, the simultaneous treatments of I extended the lifespan. The I-induced DNA chain breakage was enhanced by the NADPH-dependent microsomal electron transport system. The enhancement was also observed at the level of isolated nuclei and cells. Vitamin K2 and II increased breakage at the cellular level by I and NADPH. I-Cu<sup>2+</sup> had the tendency to increase lipid peroxidn. by the microsomes. However, the reaction was effectively inhibited by antioxidants. I induced aldehyde formation from DNA breakage. The formation was effectively inhibited by scavenging reactions with hydralazine-HCl or isoniazid. The possibility of suppressing the side effects of I is discussed in relation to 2-thiobarbiturate-reactive compds.

=> l15 and cancer  
 L15 IS NOT A RECOGNIZED COMMAND  
 The previous command name entered was not recognized by the system.  
 For a list of commands available to you in the current file, enter  
 "HELP COMMANDS" at an arrow prompt (=>).

=> s l15 and cancer  
 452589 CANCER  
 66393 CANCERS  
 468904 CANCER  
 (CANCER OR CANCERS)

L16 4 L15 AND CANCER

=> d ibib abs total

L16 ANSWER 1 OF 4 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2009:1368423 CAPLUS

DOCUMENT NUMBER: 152:51216  
 TITLE: Drug Effects Viewed from a Signal Transduction Network Perspective  
 AUTHOR(S): Fliri, Anton F.; Logging, William T.; Volkmann, Robert A.  
 CORPORATE SOURCE: Pfizer Global Research and Development, Groton, CT, 06340, USA  
 SOURCE: Journal of Medicinal Chemistry (2009), 52(24), 8038-8046  
 CODEN: JMCMAR; ISSN: 0022-2623  
 PUBLISHER: American Chemical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Understanding how drugs affect cellular network structures and how resulting signals are translated into drug effects holds the key to the discovery of medicines. Herein we examine this cause-effect relationship by determining protein network structures associated with the generation of specific in vivo drug-effect patterns. Medicines having similar in vivo pharmacol. have been identified by a comparison of drug-effect profiles of 1320 medicines. Protein network positions reached by these medicines were ascertained by examining the coinvestigation frequency of these medicines and 1179 protein network constituents in millions of scientific investigations. Interestingly, medicine assocns. obtained by comparing by drug-effect profiles mirror those obtained by comparing drug-protein coinvestigation frequency profiles, demonstrating that these drug-protein reachability profiles are relevant to in vivo pharmacol. By using protein assocns. obtained in these investigations and independent, curated protein interaction information, drug-mediated protein network topol. models can be constructed. These protein network topol. models reveal that drugs having similar pharmacol. profiles reach similar discrete positions in cellular protein network systems and provide a network view of medicine cause-effect relationships.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L16 ANSWER 2 OF 4 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:1215784 CAPLUS  
 DOCUMENT NUMBER: 147:491621  
 TITLE: Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer  
 INVENTOR(S): Mazzio, Elizabeth; Soliman, Karam  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S. Ser. No. 233,279.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070248693	A1	20071025	US 2007-711883	20070227
US 20060035981	A1	20060216	US 2005-233279	20050920
PRIORITY APPLN. INFO.:			US 2003-491841P	P 20030802
			US 2004-540525P	P 20040129
			US 2004-909590	B2 20040802
			US 2005-233279	A2 20050920

AB The invention describes a pharmaceutical composition and method for treating cancer comprising (a) 2,3-dimethoxy-5-methyl-1,4-benzoquinone,

and/or (b) at least one of wild yam root, teasel root, balm of gilead bud, bakuchi seed, dichroa root, kochia seed, kanta kari, bushy knotweed rhizome, arjun, babul chall bark, opopanax and bhumi amalaki; optionally one or more of frankincense, garcinia fruit, vitex, dragons blood, mace, sage and red sandalwood with at least (c) one compound capable of maximizing oxidative mitochondrial function, preferably riboflavin or vitamin B2 derivs., FAD, FMN, 5-amino-6-(5'-phosphoribitylamino)uracil, 6,7-dimethyl-8-(1-D-ribityl)lumazine, ribitol, 5,6-dimethylbenzimidazole, tetrahydrobiopterin, vitamin B1, lipoic acid, biotin, vitamin B6, vitamin B12, folate, niacin, vitamin C and pantothenate, and/or (d) at least one lactic acid dehydrogenase inhibitor, preferably 2',3,4',5,7-pentahydroxyflavone and optionally (f) an alkalizing agent (Aloe vera, chlorella, wheat grass, sodium or potassium bicarbonate, potassium), (g) an antiproliferative herb (speranskia or goldenseal), and (h) a pharmaceutically acceptable carrier. A method for inhibiting cancer optionally comprises one or more chemotherapy drug(s), selected, among others, from acetogenins, actinomycin D, adriamycin, aminoglutethimide, asparaginase, bleomycin, bullatacin, busulfan, carmustine, carboplatin, chlorambucil, cisplatin, etc. Thus, a composition comprised rosemary (Rosmarinus officinalis) .apprx.1000, myrrh gum (Commiphora molmol) .apprx.500, 2,3-dimethoxy-5-methyl-1,4 benzoquinone .apprx.800, and riboflavin .apprx.300 mg/day, resp.

L16 ANSWER 3 OF 4 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:824492 CAPLUS

DOCUMENT NUMBER: 143:222525

TITLE: Method of using 3-cyano-4-arylpyridine derivatives as modulators of androgen receptor function, preparation thereof, and use with other agents

INVENTOR(S): Nirschl, Alexandra A.; Hamann, Lawrence G.

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 25 pp.

CODEN: USXXCO

DOCUMENT TYPE: Patent


LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	---	-----	-----	-----
US 20050182105	A1	20050818	US 2005-48437	20050201
PRIORITY APPLN. INFO.:			US 2004-541780P	P 20040204
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE			IN LSUS DISPLAY FORMAT	
OTHER SOURCE(S):	MARPAT 143:222525			

GI



AB A method is provided for treating androgen receptor-associated conditions, such as age-related diseases, e.g. sarcopenia, employing a compound I [R1 = CN, H; X = O, S; R2 = (substituted) alkyl, (substituted) cycloalkyl, etc; R3, R4 = H, (substituted) alkyl, etc.; G = (substituted) aryl, (substituted) heteroaryl], or a pharmaceutically acceptable salt or



APPS ----- AI, PRAI  
 BIB ----- AN, plus Bibliographic Data and PI table (default)  
 CAN ----- List of CA abstract numbers without answer numbers  
 CBIB ----- AN, plus Compressed Bibliographic Data  
 CLASS ----- IPC, NCL, ECLA, FTERM  
 DALL ----- ALL, delimited (end of each field identified)  
 DMAX ----- MAX, delimited for post-processing  
 FAM ----- AN, PI and PRAI in table, plus Patent Family data  
 FBIB ----- AN, BIB, plus Patent FAM  
 IND ----- Indexing data  
 IPC ----- International Patent Classifications  
 MAX ----- ALL, plus Patent FAM, RE  
 PATS ----- PI, SO  
 SAM ----- CC, SX, TI, ST, IT  
 SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;  
                   SCAN must be entered on the same line as the DISPLAY,  
                   e.g., D SCAN or DISPLAY SCAN)  
 STD ----- BIB, CLASS  
  
 IABS ----- ABS, indented with text labels  
 IALL ----- ALL, indented with text labels  
 IBIB ----- BIB, indented with text labels  
 IMAX ----- MAX, indented with text labels  
 ISTD ----- STD, indented with text labels  
  
 OBIB ----- AN, plus Bibliographic Data (original)  
 OIBIB ----- OBIB, indented with text labels  
  
 SBIB ----- BIB, no citations  
 SIBIB ----- IBIB, no citations  
  
 HIT ----- Fields containing hit terms  
 HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)  
                   containing hit terms  
 HITRN ----- HIT RN and its text modification  
 HITSTR ----- HIT RN, its text modification, its CA index name, and  
                   its structure diagram  
 HITSEQ ----- HIT RN, its text modification, its CA index name, its  
                   structure diagram, plus NTE and SEQ fields  
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and  
                   its structure diagram  
 FHITSEQ ----- First HIT RN, its text modification, its CA index name, its  
                   structure diagram, plus NTE and SEQ fields  
 KWIC ----- Hit term plus 20 words on either side  
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI; TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.

ENTER DISPLAY FORMAT (BIB):d l15 and ?carcinoma  
 'D' IS NOT A VALID FORMAT FOR FILE 'CAPLUS'

The following are valid formats:

ABS ----- GI and AB  
 ALL ----- BIB, AB, IND, RE  
 APPS ----- AI, PRAI  
 BIB ----- AN, plus Bibliographic Data and PI table (default)  
 CAN ----- List of CA abstract numbers without answer numbers  
 CBIB ----- AN, plus Compressed Bibliographic Data  
 CLASS ----- IPC, NCL, ECLA, FTERM  
 DALL ----- ALL, delimited (end of each field identified)  
 DMAX ----- MAX, delimited for post-processing  
 FAM ----- AN, PI and PRAI in table, plus Patent Family data  
 FBIB ----- AN, BIB, plus Patent FAM  
 IND ----- Indexing data  
 IPC ----- International Patent Classifications  
 MAX ----- ALL, plus Patent FAM, RE  
 PATS ----- PI, SO  
 SAM ----- CC, SX, TI, ST, IT  
 SCAN ----- CC, SX, TI, ST, IT (random display, no answer numbers;  
               SCAN must be entered on the same line as the DISPLAY,  
               e.g., D SCAN or DISPLAY SCAN)  
 STD ----- BIB, CLASS  
  
 IABS ----- ABS, indented with text labels  
 IALL ----- ALL, indented with text labels  
 IBIB ----- BIB, indented with text labels  
 IMAX ----- MAX, indented with text labels  
 ISTD ----- STD, indented with text labels  
  
 OBIB ----- AN, plus Bibliographic Data (original)  
 OIBIB ----- OBIB, indented with text labels  
  
 SBIB ----- BIB, no citations  
 SIBIB ----- IBIB, no citations  
  
 HIT ----- Fields containing hit terms  
 HITIND ----- IC, ICA, ICI, NCL, CC and index field (ST and IT)  
               containing hit terms  
 HITRN ----- HIT RN and its text modification  
 HITSTR ----- HIT RN, its text modification, its CA index name, and  
               its structure diagram  
 HITSEQ ----- HIT RN, its text modification, its CA index name, its  
               structure diagram, plus NTE and SEQ fields  
 FHITSTR ----- First HIT RN, its text modification, its CA index name, and  
               its structure diagram  
 FHITSEQ ----- First HIT RN, its text modification, its CA index name, its  
               structure diagram, plus NTE and SEQ fields  
 KWIC ----- Hit term plus 20 words on either side  
 OCC ----- Number of occurrence of hit term and field in which it occurs

To display a particular field or fields, enter the display field codes. For a list of the display field codes, enter HELP DFIELDS at an arrow prompt (=>). Examples of formats include: TI, TI,AU; BIB,ST; TI,IND; TI,SO. You may specify the format fields in any order and the information will be displayed in the same order as the format specification.

All of the formats (except for SAM, SCAN, HIT, HITIND, HITRN, HITSTR, FHITSTR, HITSEQ, FHITSEQ, KWIC, and OCC) may be used with DISPLAY ACC to view a specified Accession Number.  
 ENTER DISPLAY FORMAT (BIB):bib



AN 2009:1557971 CAPLUS  
 DN 152:135746  
 TI Cheminformatics Analysis of Assertions Mined from Literature that Describe  
 Drug-Induced Liver Injury in Different Species  
 AU Fourches, Denis; Barnes, Julie C.; Day, Nicola C.; Bradley, Paul; Reed,  
 Jane Z.; Tropsha, Alexander  
 CS Laboratory of Molecular Modeling, Eshelman School of Pharmacy, University  
 of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA  
 SO Chemical Research in Toxicology (2010), 23(1), 171-183  
 CODEN: CRTOC; ISSN: 0893-228X  
 PB American Chemical Society  
 DT Journal  
 LA English  
 RE.CNT 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d his

(FILE 'HOME' ENTERED AT 13:27:42 ON 22 APR 2010)  
 FILE 'CAPLUS' ENTERED AT 13:29:20 ON 22 APR 2010  
 S UBIQUINONE/CN  
 FILE 'REGISTRY' ENTERED AT 13:29:44 ON 22 APR 2010  
 L1 0 S UBIQUINONE/CN  
 FILE 'CAPLUS' ENTERED AT 13:29:44 ON 22 APR 2010  
 L2 0 S L1  
 S COENZYME Q10/CN  
 FILE 'REGISTRY' ENTERED AT 13:30:10 ON 22 APR 2010  
 L3 1 S COENZYME Q10/CN  
 FILE 'CAPLUS' ENTERED AT 13:30:10 ON 22 APR 2010  
 L4 5610 S L3  
 FILE 'REGISTRY' ENTERED AT 13:30:20 ON 22 APR 2010  
 L5 1 S COENZYME Q10/CN  
 FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010  
 L6 41 S L5 AND CARCINOMA  
 L7 16 S L6 AND PY<=2004  
 L8 7 S L5 AND CARCINOMA AND TOPICAL  
 L9 38 S L5 AND ?CARCINOMA  
 L10 8 S L9 AND (TOPICAL OR SURFACE)  
 L11 13 S L5 AND CANCER AND (TOPICAL OR SURFACE)  
 L12 5 S L11 AND PY<=2004  
 FILE 'REGISTRY' ENTERED AT 13:48:56 ON 22 APR 2010  
 L13 1 S FLUOROURACIL/CN  
 L14 0 S L13 AND L5  
 FILE 'CAPLUS' ENTERED AT 13:50:02 ON 22 APR 2010  
 L15 27 S L13 AND L5  
 L16 4 S L15 AND CANCER  
 => s l15 and ?carcinoma  
 243285 ?CARCINOMA  
 L17 2 L15 AND ?CARCINOMA

=> d ibib abs

L17 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2009:1368423 CAPLUS

DOCUMENT NUMBER: 152:51216

TITLE: Drug Effects Viewed from a Signal Transduction Network Perspective

AUTHOR(S): Fliri, Anton F.; Logging, William T.; Volkmann, Robert A.

CORPORATE SOURCE: Pfizer Global Research and Development, Groton, CT, 06340, USA

SOURCE: Journal of Medicinal Chemistry (2009), 52(24), 8038-8046

CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Understanding how drugs affect cellular network structures and how resulting signals are translated into drug effects holds the key to the discovery of medicines. Herein we examine this cause-effect relationship by determining protein network structures associated with the generation of specific in vivo drug-effect patterns. Medicines having similar in vivo pharmacol. have been identified by a comparison of drug-effect profiles of 1320 medicines. Protein network positions reached by these medicines were ascertained by examining the coinvestigation frequency of these medicines and 1179 protein network constituents in millions of scientific investigations. Interestingly, medicine assocns. obtained by comparing by drug-effect profiles mirror those obtained by comparing drug-protein coinvestigation frequency profiles, demonstrating that these drug-protein reachability profiles are relevant to in vivo pharmacol. By using protein assocns. obtained in these investigations and independent, curated protein interaction information, drug-mediated protein network topol. models can be constructed. These protein network topol. models reveal that drugs having similar pharmacol. profiles reach similar discrete positions in cellular protein network systems and provide a network view of medicine cause-effect relationships.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> d ibib abs total

L17 ANSWER 1 OF 2 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2009:1368423 CAPLUS

DOCUMENT NUMBER: 152:51216

TITLE: Drug Effects Viewed from a Signal Transduction Network Perspective

AUTHOR(S): Fliri, Anton F.; Logging, William T.; Volkmann, Robert A.

CORPORATE SOURCE: Pfizer Global Research and Development, Groton, CT, 06340, USA

SOURCE: Journal of Medicinal Chemistry (2009), 52(24), 8038-8046

CODEN: JMCMAR; ISSN: 0022-2623

PUBLISHER: American Chemical Society

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Understanding how drugs affect cellular network structures and how resulting signals are translated into drug effects holds the key to the discovery of medicines. Herein we examine this cause-effect relationship by determining protein network structures associated with the generation of

specific in vivo drug-effect patterns. Medicines having similar in vivo pharmacol. have been identified by a comparison of drug-effect profiles of 1320 medicines. Protein network positions reached by these medicines were ascertained by examining the coinvestigation frequency of these medicines and 1179 protein network constituents in millions of scientific investigations. Interestingly, medicine assocns. obtained by comparing by drug-effect profiles mirror those obtained by comparing drug-protein coinvestigation frequency profiles, demonstrating that these drug-protein reachability profiles are relevant to in vivo pharmacol. By using protein assocns. obtained in these investigations and independent, curated protein interaction information, drug-mediated protein network topol. models can be constructed. These protein network topol. models reveal that drugs having similar pharmacol. profiles reach similar discrete positions in cellular protein network systems and provide a network view of medicine cause-effect relationships.

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L17 ANSWER 2 OF 2 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2007:1215784 CAPLUS

DOCUMENT NUMBER: 147:491621

TITLE: Nutraceutical composition comprising 2,3-dimethoxy-5-methyl-1,4-benzoquinone and method of use for treatment/prevention of cancer

INVENTOR(S): Mazzio, Elizabeth; Soliman, Karam

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S.

Ser. No. 233,279.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20070248693	A1	20071025	US 2007-711883	20070227
US 20060035981	A1	20060216	US 2005-233279	20050920
PRIORITY APPLN. INFO.:				
			US 2003-491841P	P 20030802
			US 2004-540525P	P 20040129
			US 2004-909590	B2 20040802
			US 2005-233279	A2 20050920

AB The invention describes a pharmaceutical composition and method for treating cancer comprising (a) 2,3-dimethoxy-5-methyl-1,4-benzoquinone, and/or (b) at least one of wild yam root, teasel root, balm of gilead bud, bakuchi seed, dichroa root, kochia seed, kanta kari, bushy knotweed rhizome, arjun, babul chall bark, opopanax and bhumi amalaki; optionally one or more of frankincense, garcinia fruit, vitex, dragons blood, mace, sage and red sandalwood with at least (c) one compound capable of maximizing oxidative mitochondrial function, preferably riboflavin or vitamin B2 derivs., FAD, FMN, 5-amino-6-(5'-phosphoribitylamino)uracil, 6,7-dimethyl-8-(1-D-ribityl)lumazine, ribitol, 5,6-dimethylbenzimidazole, tetrahydrobiopterin, vitamin B1, lipioic acid, biotin, vitamin B6, vitamin B12, folate, niacin, vitamin C and pantothenate, and/or (d) at least one lactic acid dehydrogenase inhibitor, preferably 2',3,4',5,7-pentahydroxyflavone and optionally (f) an alkalinizing agent (Aloe vera, chlorella, wheat grass, sodium or potassium bicarbonate, potassium), (g) an antiproliferative herb (speranskia or goldenseal), and (h) a pharmaceutically acceptable carrier. A method for inhibiting cancer optionally comprises one or more chemotherapy drug(s), selected, among others, from acetogenins, actinomycin D, adriamycin, aminoglutethimide,

asparaginase, bleomycin, bullatacin, busulfan, carmustine, carboplatin, chlorambucil, cisplatin, etc. Thus, a composition comprised rosemary (Rosmarinus officinalis) .apprx.1000, myrrh gum (Commiphora molmol) .apprx.500, 2,3-dimethoxy-5-methyl-4 benzoquinone .apprx.800, and riboflavin .apprx.300 mg/day, resp.

=> d his

(FILE 'HOME' ENTERED AT 13:27:42 ON 22 APR 2010)

FILE 'CAPLUS' ENTERED AT 13:29:20 ON 22 APR 2010  
S UBIQUINONE/CN

L1 FILE 'REGISTRY' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S UBIQUINONE/CN

L2 FILE 'CAPLUS' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S L1  
S COENZYME Q10/CN

L3 FILE 'REGISTRY' ENTERED AT 13:30:10 ON 22 APR 2010  
1 S COENZYME Q10/CN

L4 FILE 'CAPLUS' ENTERED AT 13:30:10 ON 22 APR 2010  
5610 S L3

L5 FILE 'REGISTRY' ENTERED AT 13:30:20 ON 22 APR 2010  
1 S COENZYME Q10/CN

L6 FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010  
41 S L5 AND CARCINOMA  
L7 16 S L6 AND PY<=2004  
L8 7 S L5 AND CARCINOMA AND TOPICAL  
L9 38 S L5 AND ?CARCINOMA  
L10 8 S L9 AND (TOPICAL OR SURFACE)  
L11 13 S L5 AND CANCER AND (TOPICAL OR SURFACE)  
L12 5 S L11 AND PY<=2004

L13 FILE 'REGISTRY' ENTERED AT 13:48:56 ON 22 APR 2010  
1 S FLUOROURACIL/CN  
L14 0 S L13 AND L5

L15 FILE 'CAPLUS' ENTERED AT 13:50:02 ON 22 APR 2010  
27 S L13 AND L5  
L16 4 S L15 AND CANCER  
L17 2 S L15 AND ?CARCINOMA

=> l13 and (topical or surface)

L13 IS NOT A RECOGNIZED COMMAND

The previous command name entered was not recognized by the system.  
For a list of commands available to you in the current file, enter  
"HELP COMMANDS" at an arrow prompt (=>).

=> s l13 and (topical or surface)

22369 L13  
60702 TOPICAL  
49 TOPICALS  
60723 TOPICAL  
(TOPICAL OR TOPICALS)  
2995599 SURFACE

540221 SURFACES  
 3213094 SURFACE  
 (SURFACE OR SURFACES)  
 L18 1153 L13 AND (TOPICAL OR SURFACE)  
 => s l18 and cancer  
 452589 CANCER  
 66393 CANCERS  
 468904 CANCER  
 (CANCER OR CANCERS)  
 L19 388 L18 AND CANCER  
 => s l18 and ?carcinoma  
 243285 ?CARCINOMA  
 L20 260 L18 AND ?CARCINOMA  
 => l20 and py<=2004  
 L20 IS NOT A RECOGNIZED COMMAND  
 The previous command name entered was not recognized by the system.  
 For a list of commands available to you in the current file, enter  
 "HELP COMMANDS" at an arrow prompt (=>).

=> s l20 and py<=2004  
 25157969 PY<=2004  
 L21 124 L20 AND PY<=2004

=> d ibib abs total

L21 ANSWER 1 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2009:855533 CAPLUS  
 DOCUMENT NUMBER: 151:142694  
 TITLE: Real time electronic cell sensing system and  
 applications for cytotoxicity profiling and compound  
 assays  
 INVENTOR(S): Wang, Xiaobo; Xu, Xiao; Abassi, Yama  
 PATENT ASSIGNEE(S): Acea Biosciences, Inc., USA  
 SOURCE: U.S., 73pp., Cont.-in-part of U.S. Ser. No. 987,732.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 12  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 7560269	B2	20090714	US 2005-55639	20050209
US 20040152067	A1	20040805	US 2003-705615	20031110 <--
US 7459303	B2	20081202		
US 20050112544	A1	20050526	US 2003-705447	20031110
US 7470533	B2	20081230		
US 20050153425	A1	20050714	US 2004-987732	20041112
US 7192752	B2	20070320		
CA 2575297	A1	20060209	CA 2005-2575297	20050804
WO 2006015387	A2	20060209	WO 2005-US27891	20050804
WO 2006015387	A3	20070412		

W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH,  
 CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD,  
 GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ,  
 LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA,  
 NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK,  
 SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU,

	ZA, ZM, ZW		
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA		
CA 2575573	A1	20060216	CA 2005-2575573
WO 2006017762	A2	20060216	WO 2005-US27943
WO 2006017762	A3	20070518	20050804
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW		
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AP, EA, EP, OA		
EP 1773979	A2	20070418	EP 2005-786773
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU		20050804
EP 1773980	A2	20070418	EP 2005-804409
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU		20050804
US 7468255	B2	20081223	US 2005-197994
CA 2580548	A1	20060406	CA 2005-2580548
WO 2006036952	A2	20060406	WO 2005-US34561
WO 2006036952	A3	20061221	20050927
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW		
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM		
EP 1800312	A2	20070627	EP 2005-812268
R:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, BA, HR, MK, YU		20050927
WO 2006058185	A2	20060601	WO 2005-US42684
WO 2006058185	A3	20070301	20051123
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, LY, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW		
RW:	AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, BW, GH,		

GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,			
KG, KZ, MD, RU, TJ, TM			
US 20060121446	A1	20060608	US 2005-286882
EP 1815025	A2	20070808	20051123
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,			20051123
IS, IT, LI, LT, LU, LV, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL,			
BA, HR, MK, YU			
PRIORITY APPLN. INFO.:			
		US 2002-397749P	P 20020720
		US 2002-435400P	P 20021220
		US 2003-469572P	P 20030509
		US 2003-705447	A2 20031110
		US 2003-705615	A2 20031110
		US 2003-519567P	P 20031112
		US 2004-542927P	P 20040209
		US 2004-548713P	P 20040227
		US 2004-614601P	P 20040929
		US 2004-987732	A2 20041112
		WO 2003-US22537	A1 20030718
		WO 2003-US22557	A 20030718
		US 2004-598608P	P 20040804
		US 2004-598609P	P 20040804
		US 2004-613749P	P 20040927
		US 2004-613872P	P 20040927
		WO 2004-US37696	A2 20041112
		US 2004-630071P	P 20041122
		US 2004-630131P	P 20041122
		US 2004-630809P	P 20041124
		US 2004-633019P	P 20041203
		US 2005-647075P	P 20050126
		US 2005-647159P	P 20050126
		US 2005-647189P	P 20050126
		US 2005-55639	A 20050209
		WO 2005-US4481	A2 20050209
		US 2005-653904P	P 20050217
		US 2005-660829P	P 20050310
		US 2005-660898P	P 20050310
		US 2005-673678P	P 20050421
		US 2005-689422P	P 20050610
		US 2005-197994	A 20050804
		US 2005-198831	A 20050804
		WO 2005-US27891	W 20050804
		WO 2005-US27943	W 20050804
		US 2005-235938	A 20050927
		WO 2005-US34561	W 20050927
		WO 2005-US42684	W 20051123

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention includes devices, systems, and methods for assaying cells using cell-substrate impedance monitoring. In one aspect, the invention provides cell-substrate impedance monitoring devices that comprise electrode arrays on a nonconducting substrate, in which each of the arrays has an approx. uniform electrode resistance across the entire array. In another aspect, the invention provides cell-substrate monitoring systems comprising one or more cell-substrate monitoring devices comprising multiple wells each having an electrode array, an impedance analyzer, a device station that connects arrays of individual wells to the impedance analyzer, and software for controlling the device station and impedance analyzer. In another aspect, the invention provides cellular assays that use impedance monitoring to detect changes in cell behavior or state. The methods can be used to test the effects of compds. on cells, such as in cytotoxicity assays. Methods of cytotoxicity profiling of compds. are also provided. The RT-CES system was used to

dynamically monitor cancer cell responses to chemotherapeutic compds. with characterized mechanisms, and to profile the specific cell response patterns.

OS.CITING REF COUNT: 4 THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)  
REFERENCE COUNT: 156 THERE ARE 156 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 2 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2009:4693 CAPLUS  
DOCUMENT NUMBER: 150:90589  
TITLE: Compositions and methods for treating and preventing dermatoses  
INVENTOR(S): Ford, John P.  
PATENT ASSIGNEE(S): USA  
SOURCE: U.S. Pat. Appl. Publ., 31pp., Cont.-in-part of U.S. Ser. No. 73,424.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 4  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20090005405	A1	20090101	US 2008-114602	20080502
US 20030158128	A1	20030821	US 2003-364383	20030212 <--
US 20040077589	A1	20040422	US 2003-684203	20031010 <--
US 6979688	B2	20051227		
US 20050059573	A1	20050317	US 2004-918199	20040813
US 6995165	B2	20060207		
US 20050272689	A1	20051208	US 2005-196921	20050803
US 7368456	B2	20080506		
US 20090012106	A1	20090108	US 2008-73424	20080305
US 7662829	B2	20100216		

PRIORITY APPLN. INFO.:  
US 2002-355764P P 20020212  
US 2003-364383 A2 20030212  
US 2003-684203 A2 20031010  
US 2004-918199 A1 20040813  
US 2005-196921 A1 20050803  
US 2008-73424 A2 20080305

AB The invention encompasses protectant agents including uracil or a metabolite thereof that effectively prevent and/or treat the cutaneous toxicities and dermatol. side-effects associated with chemotherapeutic agents. Addnl., and surprisingly compns. including uracil or a metabolite thereof are effective for treating or preventing various dermatoses.

L21 ANSWER 3 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2008:1537154 CAPLUS  
DOCUMENT NUMBER: 150:71117  
TITLE: Antisense oligonucleotides against thymidylate synthase  
INVENTOR(S): Koropatnick, Donald James; Dean, Nicholas Mark; Vincent, Mark D.  
PATENT ASSIGNEE(S): Can.  
SOURCE: U.S. Pat. Appl. Publ., 70pp., Cont.-in-part of U.S. Ser. No. 597,409.  
CODEN: USXXCO  
DOCUMENT TYPE: Patent  
LANGUAGE: English



FAMILY ACC. NUM. COUNT: 6  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080318891	A1	20081225	US 2007-987568	20071130
WO 9915648	A1	19990401	WO 1998-GB2820	19980917 <--
W:	AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW			
RW:	GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
WO 2005070469	A1	20050804	WO 2005-CA69	20050124
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 20050272683	A1	20051208	US 2005-171435	20050701
US 20080255066	A1	20081016	US 2008-29297	20080211
PRIORITY APPLN. INFO.:			WO 1998-GB2820	W 19980917
			US 2000-509418	B1 20000711
			WO 2005-CA69	W 20050124
			US 2005-171435	A2 20050701
			US 2006-597409	A2 20060724
			GB 1997-20107	A 19970923
			GB 1997-22012	A 19971017
			GB 1998-12140	A 19980606
			CA 2002-2380970	A 20020408
			WO 2003-CA480	W 20030408
			US 2004-538886P	P 20040123
			US 2004-556368P	P 20040326
			US 2005-661993P	P 20050311
			US 2005-510899	A2 20050622
			US 2006-908389	A2 20060313
			WO 2006-CA350	W 20060313
			US 2007-911759P	P 20070413
			US 2007-987568	A2 20071130
AB	Antisense oligonucleotides targeted to sequences in thymidylate synthase (TS) mRNA are provided. In particular the invention relates to antisense oligonucleotides targeted to sequences in the 3' end of TS mRNA, which are both cytostatic on their own when administered to human tumor cell lines, and which also enhance the toxicity of anticancer drugs. The invention further relates to a combination product comprising an antisense oligonucleotide in combination with an anticancer agent such as Tomudex or pemetrexed and to the use of such a combination product in the treatment of cancer.			
L21 ANSWER 4 OF 124	CAPLUS COPYRIGHT 2010 ACS on STN			
ACCESSION NUMBER:	2008:975257 CAPLUS			
DOCUMENT NUMBER:	149:267804			
TITLE:	Purine compounds as A2b adenosine receptor antagonists			

INVENTOR(S): and their preparation  
Kalla, Rao; Perry, Thao; Elzein, Elfatih; Li, Xiaofen;  
Zablocki, Jeff; Zeng, Dewan; Xiao, Dengming;  
Varkhedkar, Vaibhav; Ibrahim, Prabha; Palle, Venkata;  
Zhong, Hongyan

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 47pp., Cont.-in-part of Ser.  
No. US 2005-189202, filed on 25 Jul 2005, now  
patentedPa  
CODEN: USXXCO


DOCUMENT TYPE: Patent

LANGUAGE: English


FAMILY ACC. NUM. COUNT: 7

PATENT INFORMATION:


PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20080194593	A1	20080814	US 2008-13348	20080111
US 20030139428	A1	20030724	US 2002-290921	20021108 <--
US 6825349	B2	20041130		
US 20030229106	A1	20031211	US 2003-431167	20030506 <--
US 6977300	B2	20051220		
CA 2524778	A1	20041209	CA 2003-2524778	20030506 <--
AU 2003249604	A1	20050121	AU 2003-249604	20030506
EP 1622908	A1	20060208	EP 2003-817096	20030506
EP 1622908	B1	20080806		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
CN 1771250	A	20060510	CN 2003-826411	20030506
JP 2006515316	T	20060525	JP 2005-500296	20030506
RU 2318825	C2	20080310	RU 2005-134232	20030506
AT 403656	T	20080815	AT 2003-817096	20030506
PT 1622908	E	20081114	PT 2003-817096	20030506
NZ 543416	A	20090131	NZ 2003-543416	20030506
ES 2311759	T3	20090216	ES 2003-817096	20030506
ZA 2004003485	A	20050622	ZA 2004-3485	20040507
US 20050261316	A1	20051124	US 2005-189202	20050725
US 7317017	B2	20080108		
ZA 2005008262	A	20070228	ZA 2005-8262	20051012
MX 2005011860	A	20060217	MX 2005-11860	20051104
KR 2006055453	A	20060523	KR 2005-720956	20051104
HK 1092137	A1	20090430	HK 2006-108745	20060807
US 20080318983	A1	20081225	US 2008-147382	20080626
PRIORITY APPLN. INFO.:				
			US 2001-348222P	P 20011109
			US 2002-290921	A2 20021108
			US 2003-431167	A1 20030506
			US 2005-189202	A2 20050725
			US 2002-401408P	P 20020805
			WO 2003-US14085	W 20030506
			US 2008-13348	A2 20080111
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT				
OTHER SOURCE(S): MARPAT 149:267804				
GI				



I



II



III

AB Disclosed are methods for treating asthma, inflammatory gastrointestinal tract disorders, cancer, cardiovascular diseases, neurol. disorders, and diseases related to undesirable angiogenesis using A2B adenosine receptor antagonists having the structure of formula I or formula II. Comps. of formula I and II wherein R1 and R2 are independently H, (un)substituted alkyl and D-E; D is a bond and alkylene; E is (un)substituted alkoxy, (un)substituted cycloalkyl, (un)substituted (hetero)aryl, and (un)substituted alkynyl, with the proviso that when D is a covalent bond, then E cannot be alkoxy; R3 is H, (un)substituted alkyl, and (un)substituted cycloalkyl; C is (un)substituted (hetero)arylene; Y is a covalent bond, (un)substituted alkylene, etc.; Z is H, (un)substituted monocyclic (hetero)aryl; are claimed. Example compound III was prepared by alkylation of 7-benzyl-1,3,7-trihydropurine-2,6-dione with 1-iodopropane. All the invention compds. were evaluated for their A2b adenosine receptor antagonistic activity (some data given).

L21 ANSWER 5 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2006:789739 CAPLUS

DOCUMENT NUMBER: 145:202883

TITLE: Hyaluronan as a cytotoxic agent, drug pre-sensitizer and chemo-sensitizer in the treatment of disease

INVENTOR(S): Brown, Tracey; Fox, Richard

PATENT ASSIGNEE(S): Mediatech Research Ltd., Australia

SOURCE: U.S. Pat. Appl. Publ., 53 pp., Cont.-in-part of U.S. Ser. No. 88,774.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20060178342	A1	20060810	US 2005-198663	20050805
WO 2002005852	A1	20020124	WO 2001-AU849	20010713 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,  
 DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF,  
 BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG  
 US 20030180382 A1 20030925 US 2003-88774 20030313 <--  
 PRIORITY APPLN. INFO.: WO 2001-AU849 W 20010713  
 US 2003-88774 A2 20030313  
 AU 2000-8795 A 20000714

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB This application provides methods and compns. for the treatment of cancer.  
 The application provides compns. comprising hyaluronic acid and a  
 chemotherapeutic agent such as irinotecan that are useful in the treatment  
 of cancer.

L21 ANSWER 6 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:1314363 CAPLUS  
 DOCUMENT NUMBER: 144:57544  
 TITLE: Antibody drug conjugates and uses for cancer therapy  
 INVENTOR(S): Ebens, Allen J., Jr.; Jacobson, Frederic S.; Polakis,  
 Paul; Schwall, Ralph H.; Sliwkowski, Mark X.; Spencer,  
 Susan D.  
 PATENT ASSIGNEE(S): Genentech, Inc., USA  
 SOURCE: PCT Int. Appl., 110 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 167  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2005117986	A2	20051215	WO 2005-US18829	20050531
WO 2005117986	A3	20060615		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, GU, HU, ID, IL, IN, IS, JP, KE, KG, KM, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
NZ 528704	A	20050225	NZ 1999-528704	19990308
EP 1466977	A1	20041013	EP 2004-7618	19991202 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY			
NZ 523206	A	20041224	NZ 2000-523206	20000211 <--
NZ 523207	A	20041224	NZ 2000-523207	20000211 <--
NZ 523208	A	20041224	NZ 2000-523208	20000211 <--
NZ 523209	A	20041224	NZ 2000-523209	20000211 <--
CA 2481685	A1	20010308	CA 2000-2481685	20000824 <--
CA 2481691	A1	20010308	CA 2000-2481691	20000824 <--
CA 2481731	A1	20010308	CA 2000-2481731	20000824 <--
CA 2481756	A1	20010308	CA 2000-2481756	20000824 <--
CA 2481788	A1	20010308	CA 2000-2481788	20000824 <--
EP 1657253	A1	20060517	EP 2005-24024	20010601
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY, AL, TR			
EP 1657254	A2	20060517	EP 2005-24029	20010601

EP 1657254	A3	20060607		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI, CY, TR				
EP 1657251	A2	20060517	EP 2005-24036	20010601
EP 1657251	A3	20060524		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI, CY, AL, TR				
CN 1210696	C	20050713	CN 2001-811390	20010613
AU 758921	B2	20030403	AU 2001-57764	20010801 <--
AU 759004	B2	20030403	AU 2001-57765	20010801 <--
CA 2420193	A1	20020228	CA 2001-2420193	20010823 <--
JP 2004520810	T	20040715	JP 2002-522275	20010823 <--
US 20030073129	A1	20030417	US 2001-946374	20010904 <--
US 20030207803	A1	20031106	US 2001-143026	20011019 <--
US 20030199021	A1	20031023	US 2001-13924	20011025 <--
EP 1397383	A2	20040317	EP 2001-990229	20011213 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
AU 772759	B2	20040506	AU 2002-14767	20020201 <--
AU 772723	B2	20040506	AU 2002-14769	20020201 <--
AU 772734	B2	20040506	AU 2002-14771	20020201 <--
AU 778585	B2	20041209	AU 2002-14753	20020201 <--
CA 2449602	A1	20021219	CA 2002-2449602	20020403 <--
WO 2002101069	A2	20021219	WO 2002-US10513	20020403 <--
WO 2002101069	A3	20030904		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH,				
GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR,				
LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH,				
PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ,				
UA, UG, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,				
KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB,				
GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA,				
GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002303231	A1	20021223	AU 2002-303231	20020403 <--
AU 2002303231	B2	20080306		
EP 1402260	A2	20040331	EP 2002-731246	20020403 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2005500030	T	20050106	JP 2003-503819	20020403
US 20030207351	A1	20031106	US 2002-140473	20020506 <--
US 20030148438	A1	20030807	US 2002-145821	20020514 <--
US 20030170788	A1	20030911	US 2002-145634	20020514 <--
US 20030207371	A1	20031106	US 2002-145749	20020514 <--
US 20030166084	A1	20030904	US 2002-146793	20020515 <--
US 20030134380	A1	20030717	US 2002-147509	20020516 <--
US 20040214269	A1	20041028	US 2002-147518	20020516 <--
US 20030180875	A1	20030925	US 2002-147505	20020517 <--
US 20030199027	A1	20031023	US 2002-152396	20020520 <--
US 20050074837	A1	20050407	US 2002-158788	20020530
US 20030064446	A1	20030403	US 2002-194460	20020711 <--
US 20030153037	A1	20030814	US 2002-194457	20020711 <--
US 20030059879	A1	20030327	US 2002-194456	20020712 <--
US 20030064448	A1	20030403	US 2002-194484	20020712 <--
US 20030049747	A1	20030313	US 2002-195899	20020715 <--
US 20030064449	A1	20030403	US 2002-195884	20020715 <--
US 20030063112	A1	20030403	US 2002-195896	20020715 <--
US 20030068705	A1	20030410	US 2002-195886	20020715 <--
US 20030071834	A1	20030417	US 2002-195898	20020715 <--
US 20030049748	A1	20030313	US 2002-196748	20020716 <--

US	20030059880	A1	20030327	US	2002-196758	20020716	<--
US	20030064451	A1	20030403	US	2002-196755	20020716	<--
US	20030068710	A1	20030410	US	2002-196761	20020716	<--
US	20030162250	A1	20030828	US	2002-196747	20020716	<--
US	20030215910	A1	20031120	US	2002-199463	20020718	<--
US	20030180881	A1	20030925	US	2002-202475	20020723	<--
US	20030049777	A1	20030313	US	2002-202939	20020724	<--
US	20030054482	A1	20030320	US	2002-205902	20020725	<--
US	20030064462	A1	20030403	US	2002-206919	20020726	<--
US	20030064463	A1	20030403	US	2002-206922	20020726	<--
US	20030068754	A1	20030410	US	2002-206910	20020726	<--
US	20030068756	A1	20030410	US	2002-206912	20020726	<--
US	20030068759	A1	20030410	US	2002-206920	20020726	<--
US	20030068760	A1	20030410	US	2002-206921	20020726	<--
US	20030073183	A1	20030417	US	2002-206917	20020726	<--
US	20030096359	A1	20030522	US	2002-205910	20020726	<--
US	20040048334	A1	20040311	US	2002-205890	20020726	<--
US	20030064467	A1	20030403	US	2002-207921	20020729	<--
US	20030064468	A1	20030403	US	2002-207922	20020729	<--
US	20030064469	A1	20030403	US	2002-208027	20020729	<--
US	20030068766	A1	20030410	US	2002-207917	20020729	<--
US	20030068767	A1	20030410	US	2002-207918	20020729	<--
US	20030068768	A1	20030410	US	2002-207919	20020729	<--
US	20030068769	A1	20030410	US	2002-207920	20020729	<--
US	20030068770	A1	20030410	US	2002-207925	20020729	<--
US	20030068771	A1	20030410	US	2002-208021	20020729	<--
US	20030068772	A1	20030410	US	2002-208022	20020729	<--
US	20030068773	A1	20030410	US	2002-208023	20020729	<--
US	20030068774	A1	20030410	US	2002-208026	20020729	<--
US	20030068776	A1	20030410	US	2002-208030	20020729	<--
US	20030073184	A1	20030417	US	2002-207923	20020729	<--
US	20030073185	A1	20030417	US	2002-207924	20020729	<--
US	20030073186	A1	20030417	US	2002-208028	20020729	<--
US	20030215912	A1	20031120	US	2002-207915	20020729	<--
US	20040048335	A1	20040311	US	2002-208024	20020729	<--
AU	2002301608	A1	20030306	AU	2002-301608	20021023	<--
AU	2002301608	B2	20041118				
AU	2002301732	A1	20030306	AU	2002-301732	20021024	<--
AU	2002301732	B2	20050414				
AU	2002301730	A1	20030306	AU	2002-301730	20021025	<--
AU	2002301730	B2	20041014				
AU	2002301731	A1	20030306	AU	2002-301731	20021025	<--
AU	2002301731	B2	20050526				
AU	2002301722	A1	20030306	AU	2002-301722	20021028	<--
AU	2002301722	B2	20040916				
AU	2002301723	A1	20030313	AU	2002-301723	20021028	<--
AU	2002301723	B2	20041118				
US	20030120056	A1	20030626	US	2002-289498	20021105	<--
US	20030144498	A1	20030731	US	2002-289527	20021105	<--
US	20040249141	A1	20041209	US	2002-289490	20021105	<--
US	20030211576	A1	20031113	US	2002-298993	20021118	<--
US	20030224984	A1	20031204	US	2002-305654	20021126	<--
AU	2002313838	A1	20030327	AU	2002-313838	20021202	<--
AU	2002313838	B2	20050526				
AU	2002330288	A1	20030417	AU	2002-330288	20021202	<--
AU	2002330288	B2	20050505				
AU	2002317529	A1	20030410	AU	2002-317529	20021205	<--
AU	2002317529	B2	20050526				
AU	2002318894	A2	20030410	AU	2002-318894	20021213	<--
AU	2002318894	A1	20030410				
AU	2002318894	B2	20050512				

AU 2002363307	A1	20030710	AU 2002-363307	20021216 <--
AU 2002363307	B2	20050526		
AU 2002323928	A1	20030403	AU 2002-323928	20021220 <--
AU 2002323928	B2	20050811		
AU 2003200137	A1	20030417	AU 2003-200137	20030115 <--
AU 2003200204	A1	20030410	AU 2003-200204	20030120 <--
AU 2003200204	B2	20050512		
AU 2003200585	A1	20030501	AU 2003-200585	20030213 <--
AU 2003200585	B2	20060713		
AU 2003200721	A1	20030501	AU 2003-200721	20030225 <--
AU 2003200721	B2	20051208		
AU 2003200722	A1	20030501	AU 2003-200722	20030225 <--
AU 2003200722	B2	20060202		
AU 2003200731	A1	20030501	AU 2003-200731	20030225 <--
AU 2003200731	B2	20060413		
AU 2003200740	A1	20030612	AU 2003-200740	20030226 <--
AU 2003200740	B2	20051215		
AU 2003201315	A1	20030612	AU 2003-201315	20030228 <--
AU 2003203679	A1	20030612	AU 2003-203679	20030411 <--
AU 2003203679	B2	20070405		
AU 2003204018	A1	20030612	AU 2003-204018	20030505 <--
AU 2003204018	B2	20050428		
AU 2003204357	A1	20030619	AU 2003-204357	20030526 <--
AU 2003204357	B2	20060330		
AU 2003257515	A1	20031120	AU 2003-257515	20031023 <--
AU 2003257515	B2	20051201		
AU 2003259607	A1	20031127	AU 2003-259607	20031031 <--
AU 2003259607	B2	20051103		
US 20040258710	A1	20041223	US 2004-791618	20040302 <--
US 20050019823	A1	20050127	US 2004-931886	20040831
US 20050187382	A1	20050825	US 2004-950374	20040923
US 20050153396	A1	20050714	US 2004-955952	20040929
US 20050153348	A1	20050714	US 2004-20604	20041221
US 20050176041	A1	20050811	US 2004-26279	20041230
US 20050214819	A1	20050929	US 2005-30464	20050105
US 20050164266	A1	20050728	US 2005-36582	20050113
US 20050170396	A1	20050804	US 2005-36869	20050114
US 20050202475	A1	20050915	US 2005-38328	20050118
US 20050176104	A1	20050811	US 2005-52503	20050204
US 20050136515	A1	20050623	US 2005-56802	20050211
US 20050136475	A1	20050623	US 2005-60652	20050216
US 20050214846	A1	20050929	US 2005-117757	20050427
AU 2005249490	A1	20051215	AU 2005-249490	20050531
CA 2567520	A1	20051215	CA 2005-2567520	20050531
EP 1753463	A2	20070221	EP 2005-804852	20050531
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,				
IS, IT, LI, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
CN 1993146	A	20070704	CN 2005-80026066	20050531
BR 2005010883	A	20071226	BR 2005-10883	20050531
JP 2008501029	T	20080117	JP 2007-515417	20050531
NZ 551180	A	20091030	NZ 2005-551180	20050531
AU 2005205752	A1	20050922	AU 2005-205752	20050831
AU 2005205752	B2	20070607		
AU 2005205754	A1	20050922	AU 2005-205754	20050831
AU 2005205754	B2	20070920		
AU 2005205755	A1	20050922	AU 2005-205755	20050831
AU 2005205755	B2	20071004		
AU 2005205758	A1	20050922	AU 2005-205758	20050831
AU 2005205758	B2	20070913		
US 20070191270	A1	20070816	US 2005-234694	20050922
US 20080166294	A1	20080710	US 2006-461752	20060801

JP 2007037551	A	20070215	JP 2006-221327	20060814
JP 4166254	B2	20081015		
IN 2006DN07052	A	20070713	IN 2006-DN7052	20061123
US 20070269446	A1	20071122	US 2006-564171	20061128
MX 2006014065	A	20070131	MX 2006-14065	20061130
KR 2007037575	A	20070405	KR 2006-725240	20061130
AU 2006252290	A1	20070125	AU 2006-252290	20061229
AU 2006252290	B2	20091126		
NO 2006006075	A	20070228	NO 2006-6075	20061229
US 20070219350	A1	20070920	US 2007-801111	20070507
AU 2007203155	A1	20070726	AU 2007-203155	20070705
AU 2007203291	A1	20070802	AU 2007-203291	20070717
AU 2007214325	A1	20070920	AU 2007-214325	20070830
US 20090148878	A1	20090611	US 2008-77504	20080318
AU 2008201807	A1	20080515	AU 2008-201807	20080424
AU 2008201995	A1	20080529	AU 2008-201995	20080506
AU 2008201996	A1	20080529	AU 2008-201996	20080506
AU 2008201998	A1	20080529	AU 2008-201998	20080506
AU 2008202000	A1	20080529	AU 2008-202000	20080506
AU 2008202004	A1	20080529	AU 2008-202004	20080506
AU 2008202007	A1	20080529	AU 2008-202007	20080506
US 20090075278	A1	20090319	US 2008-156160	20080529
US 20090075279	A1	20090319	US 2008-156180	20080529
PRIORITY APPLN. INFO.:			US 2004-576517P	P 20040601
			US 2004-616098P	P 20041005
			US 1998-83742P	P 19980430
			US 1998-84366P	P 19980505
			US 1998-85339P	A1 19980513
			US 1998-87106P	P 19980528
			US 1998-88326P	P 19980604
			US 1998-88217P	P 19980605
			US 1998-88655P	P 19980609
			US 1998-89947P	P 19980619
			US 1998-90676P	P 19980625
			US 1998-91982P	P 19980707
			AU 1998-84850	A3 19980714
			US 1998-94651P	A1 19980730
			US 1998-96012P	P 19980810
			US 1998-97954P	P 19980826
			US 1998-97974P	P 19980826
			US 1998-97979P	P 19980826
			AU 1998-93881	A3 19980914
			AU 1998-93178	A3 19981002
			AU 1999-10703	A3 19981007
			US 1998-105169P	P 19981022
			AU 1999-11260	A3 19981029
			AU 1999-12883	A3 19981029
			WO 1998-US22992	W 19981029
			AU 1999-30721	A3 19990308
			US 1999-131293P	P 19990427
			US 1999-133459P	P 19990511
			US 1999-140650P	P 19990622
			US 1999-149395P	P 19990817
			US 1999-151689P	P 19990831
			AU 1999-55908	A3 19990901
			AU 2000-17482	A3 19991130
			AU 2000-17499	A3 19991202
			EP 1999-960644	A3 19991202
			AU 2000-28794	A3 20000211
			US 2000-189320P	P 20000314
			US 2000-189328P	P 20000314



US 2000-190828P	P	20000321
US 2000-191048P	P	20000321
US 2000-191314P	P	20000321
US 2000-193032P	P	20000329
US 2000-193053P	P	20000329
US 2000-194449P	P	20000404
US 2000-194647P	P	20000404
US 2000-195975P	P	20000411
US 2000-196000P	P	20000411
US 2000-196187P	P	20000411
US 2000-196820P	XX	20000411
US 2000-198121P	P	20000418
US 2000-198585P	P	20000418
US 2000-199397P	P	20000425
US 2000-199550P	P	20000425
US 2000-201516P	P	20000503
US 2000-204675P	P	20000517
US 2000-213087P	P	20000622
US 2000-220893P	P	20000726
US 2000-222425P	P	20000801
US 2000-227133P	P	20000822
CA 2000-2380355	A3	20000824
US 2000-232887P	P	20000915
US 2000-236009P	P	20000927
US 2000-690189	A3	20001016
JP 2002-576286	A3	20010322
US 2001-816920	B1	20010322
EP 2001-939834	A3	20010601
EP 2004-5726	A3	20010601
US 2001-880457	A	20010612
US 2001-882636	B1	20010614
AU 2001-271938	A3	20010710
AU 2001-71938	T0	20010710
US 2001-927796	B1	20010809
WO 2001-US26626	W	20010823
US 2001-992521	B1	20011114
WO 2001-US48938	W	20011213
US 2002-52586	A1	20020115
WO 2002-US10513	W	20020403
US 2002-123155	A1	20020415
US 2002-127825	A1	20020422
US 2002-141703	A1	20020508
US 2002-145627	A1	20020514
US 2002-145751	A	20020514
US 2002-146793	A1	20020515
US 2002-197703	B1	20020717
US 2002-197708	A1	20020717
US 2002-197942	B1	20020718
US 2002-199666	A1	20020718
US 2002-199464	B1	20020719
US 2002-211858	A1	20020802
AU 2002-330015	A3	20020911
AU 2002-367318	A3	20021230
AU 2003-200137	A3	20030115
AU 2003-203679	A3	20030411
AU 2003-261484	A	20031106
US 2003-520842P	P	20031117
US 2003-532426P	P	20031224
US 2004-872972	A1	20040621
US 2004-989826	A2	20041116
WO 2004-US38262	A2	20041116

AU 2005-200179	A3 20050114
US 2005-141344	A2 20050531
WO 2005-US18829	W 20050531
US 2005-234694	B1 20050922
US 2006-461752	A2 20060801
US 2007-804045	A1 20070515

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 144:57544

AB The present invention relates to antibody-drug conjugate compds. with a formula of Ab-(L-D)p where 1 to 8 (p) maytansinoid drug moieties (D) are covalently linked by L to an antibody (Ab) which binds to an ErbB receptor, or which binds to one or more tumor-associated antigens or cell-surface receptors. These compds. may be used in methods of diagnosis or treatment of cancer, and other diseases and disorders.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

REFERENCE COUNT: 18 THERE ARE 18 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 7 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:1262462 CAPLUS

DOCUMENT NUMBER: 144:590

TITLE: Hyaluronan as a cytotoxic agent, drug pre-sensitizer and chemo-sensitizer in the treatment of disease

INVENTOR(S): Brown, Tracey; Fox, Richard

PATENT ASSIGNEE(S): Mediatech Research Ltd., Australia

SOURCE: U.S. Pat. Appl. Publ., 49 pp., Cont.-in-part of U.S. Ser. No. 88,774.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20050267069	A1	20051201	US 2005-191407	20050727
WO 2002005852	A1	20020124	WO 2001-AU849	20010713 <--
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 20030180382	A1	20030925	US 2003-88774	20030313 <--
PRIORITY APPLN. INFO.:			AU 2000-8795	A 20000714
			WO 2001-AU849	W 20010713
			US 2003-88774	A2 20030313

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB This application provides methods and compns. for the treatment of cancer. The application provides compns. comprising hyaluronic acid and a chemotherapeutic agent such as irinotecan that are useful in the treatment of cancer.

L21 ANSWER 8 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:963840 CAPLUS

DOCUMENT NUMBER: 143:254038

TITLE: Bile-derived biological response modifier for the

INVENTOR(S): treatment of cancer  
 Young, Aiping H.  
 PATENT ASSIGNEE(S): Lorus Therapeutics Inc., Can.  
 SOURCE: U.S. Pat. Appl. Publ., 44 pp., Cont.-in-part of U.S.  
 Ser. No. 416,259.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20050192443	A1	20050901	US 2004-821649	20040408
WO 2002038164	A1	20020516	WO 2001-CA1558	20011108 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
US 20040101511	A1	20040527	US 2004-416259	20040102 <--
PRIORITY APPLN. INFO.:			CA 2000-2325361	A 20001108
			WO 2001-CA1558	W 20011108
			US 2004-416259	A2 20040102

AB The present invention provides anticancer bile-derived biol. response modifier (BD-BRM) for the treatment of cancer. In accordance with an aspect of the present invention, there is provided a composition for the treatment of breast and prostate cancer in a mammal, comprising (i) small mol. weight components of less than 3000 daltons, and having the following properties: is extracted from bile of animals; is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo; is capable of modulating tumor necrosis factor production and/or release; contains no measurable level of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN- $\gamma$ ; is not cytotoxic to human peripheral blood mononuclear cells; and is not an endotoxin; and optionally (ii) one or more anticancer agent(s), wherein the combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the composition or the anticancer agent(s) alone. Another aspect of the present invention provides the use of this composition or combination in the manufacture of a medicament for the treatment of breast or prostate cancer in a mammal. For example, BD-BRM, as a single agent, against human tumors xenografted in mice resulted in a significant delay of breast tumor growth as compared to saline-treated controls. The mean tumor weight of BD-BRM-treated animals was decreased by 77% at the endpoint of the experiment as compared to that of saline controls. In comparison with the tumor growth inhibition by standard chemotherapeutic drug treatments at an optimal dose, 69.4% of tumor weight reduction by Doxorubicin, or 53.2% of tumor weight reduction by Taxol, the efficacy of BD-BRM was higher than these observed in the treatment with either Doxorubicin or Taxol.

L21 ANSWER 9 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:595607 CAPLUS

DOCUMENT NUMBER: 143:400056

TITLE: Mechanism of octreotide reversing multidrug resistance in hepatoma cells

AUTHOR(S): Li, Wenhuan; Cui, Yi; Qiao, Huimei; Zhu, Juren

CORPORATE SOURCE: Shandong Provincial Hospital, Shandong University, Jinan, Shandong Province, 250021, Peop. Rep. China  
SOURCE: Shandong Daxue Xuebao, Yixueban (2004), 42(3), 290-293  
CODEN: SDXYBZ; ISSN: 1671-7554  
PUBLISHER: Shandong Daxue Xuebao, Yixueban Bianjibu  
DOCUMENT TYPE: Journal  
LANGUAGE: Chinese  
AB The mechanism of octreotide reversing multidrug resistance and the effect of octreotide on chemo-sensitizing hepatoma cells were investigated. The expression of the MDR1, MRP2 mRNA and protein were analyzed by RT-PCR and flow cytometry resp. The cytotoxic effects of epirubicin, carboplatin, hydroxyl-camptothecin and 5-fluorouracil were analyzed by MTT assay. The 50% inhibitory concentration of cytotoxic agents were significantly reduced after octreotide treatment. RT-PCR and flow cytometry showed a significantly reduced expression of P-glycoprotein and MRP2 on the cell surface of hepatoma primary culture cells after octreotide treatment. Thus, octreotide can chemo-sensitize hepatoma cells and chemosensitization can be achieved by inhibiting the expression of the MDR1, MRP2 gene on the surface of hepatoma cells.

L21 ANSWER 10 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2005:222263 CAPLUS  
DOCUMENT NUMBER: 143:103045  
TITLE: Surface modification of drug-loaded PLA nanoparticle and its evaluation in vitro  
AUTHOR(S): Hu, Yunxia; Yuan, Xubo; Zhang, Xiaojin; Guo, Yi; Chang, Jin  
CORPORATE SOURCE: School of Material Science and Engineering, Tianjin University, Tianjin, 300072, Peop. Rep. China  
SOURCE: Zhongguo Shengwu Yixue Gongcheng Xuebao (2004), 23(1), 30-36  
CODEN: ZSYXEI; ISSN: 0258-8021  
PUBLISHER: Zhongguo Yixue Kexueyuan  
DOCUMENT TYPE: Journal  
LANGUAGE: Chinese  
AB Biodegradable O-Carboxymethylated Chitosan (O-CMC) was used to modify 5-fluorouracil (5-FU)-loaded poly(lactic acid) (PLA) nanoparticles (NPs) by multiple emulsion technol. The glomeration ability, appearance, structure, and surface of NPs were characterized by AFM, TEM, and XPS. The results showed that O-CMC could be used to prepare drug-loaded NPs as an emulsifier and modifier and the mean size of NPs obtained was 50 nm. Three kinds of tumor cell lines including 803 gastric carcinoma cells, MDA-MB-231 breast carcinoma cells and HCT-8 colorectal cells were used to investigate the cytotoxicity of NPs. It was demonstrated that the 5-FU-loaded NPs had high cytotoxicity of 72.8%, 77.3%, and 75.6% resp. on the three kinds of cells. In addition the 5-FU-loaded NPs showed a sustained release for 12 days.  
OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L21 ANSWER 11 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:1154310 CAPLUS  
DOCUMENT NUMBER: 142:69220  
TITLE: Topical use of valproic acid, alone or with other agents, for the prevention or treatment of skin disorders  
INVENTOR(S): Pelicci, Pier Giuseppe; Minucci, Saverio; Costanzo, Antonio; Chimenti, Sergio; Nistico, Steven Paul; Paolino, Donatella

PATENT ASSIGNEE(S): G2M Cancer Drugs AG, Germany  
 SOURCE: Eur. Pat. Appl., 40 pp.  
 CODEN: EPXXDW  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1491188	A1	20041229	EP 2003-14278	20030625 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
AU 2004251434	A1	20050106	AU 2004-251434	20040623
AU 2004251435	A1	20050106	AU 2004-251435	20040623
CA 2531101	A1	20050106	CA 2004-2531101	20040623
CA 2531107	A1	20050106	CA 2004-2531107	20040623
WO 2005000289	A1	20050106	WO 2004-EP6789	20040623
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2005000282	A2	20050106	WO 2004-EP6797	20040623
WO 2005000282	A3	20050428		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1635808	A1	20060322	EP 2004-740209	20040623
EP 1635808	B1	20081001		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
EP 1635798	A2	20060322	EP 2004-740216	20040623
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK				
JP 2008529964	T	20080807	JP 2006-516036	20040623
JP 2008529966	T	20080807	JP 2006-516039	20040623
AT 409471	T	20081015	AT 2004-740209	20040623
PT 1635808	E	20081120	PT 2004-740209	20040623
ES 2310734	T3	20090116	ES 2004-740209	20040623
EP 2039355	A2	20090325	EP 2008-13246	20040623
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR				
US 20060160897	A1	20060720	US 2005-275258	20051221
US 20070037738	A1	20070215	US 2005-275263	20051221
PRIORITY APPLN. INFO.:			EP 2003-14278	A 20030625
			EP 2004-740209	A3 20040623

WO 2004-EP6789 W 20040623  
 WO 2004-EP6797 W 20040623

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 142:69220

AB The invention relates to a topically applicable formulation containing valproic acid or a derivative thereof which can be used alone or in combination with topically applicable formulations of retinoids or of nuclear receptor ligands, or of chemotherapeutic agents (e.g. 5-Fluorouracil). The formulation is useful for the topical treatment of cancerous skin disorders, e.g. basal cell carcinoma, squamous cell carcinoma, keratoakantoma, Bowen disease, cutaneous T-Cell lymphoma, and also for the topical treatment of premalignant lesions, and of inflammation of the skin and/or mucosa. The invention also relates to the use of this topically applicable formulation for protection from UV light and for the treatment of sunburn. The invention includes the use of valproic acid for the manufacture of a clin. used medicament for the topical treatment of the above human diseases.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)  
 REFERENCE COUNT: 11 THERE ARE 11 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 12 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2004:1126960 CAPLUS

DOCUMENT NUMBER: 142:69160

TITLE: Vimentin directed diagnostics and therapeutics for multidrug resistant (MDR) neoplastic disease, and a vaccine for treating or preventing MDR neoplasm  
 Georges, Elias; Serfass, Lucile; Bonneau, Anne-Marie; Dallaire, Frederic

INVENTOR(S): Aurelium Biopharma Inc., Can.

PATENT ASSIGNEE(S): U.S. Pat. Appl. Publ., 76 pp.

SOURCE: CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040259112	A1	20041223	US 2003-736889	20031215 <--
US 7550256	B2	20090623		
CA 2509987	A1	20050707	CA 2003-2509987	20031215
WO 2005062058	A1	20050707	WO 2003-IB6427	20031215
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, BG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1695089	A1	20060830	EP 2003-819144	20031215
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, SK				
JP 2007515924	T	20070621	JP 2005-512304	20031215
AU 2003296857	A1	20050714	AU 2003-296857	20031216
US 20060014225	A1	20060119	US 2005-173672	20050701

US 7670604 B2 20100302

PRIORITY APPLN. INFO.: US 2002-433480P P 20021213  
 US 2003-736889 A3 20031215  
 WO 2003-1B6427 W 20031215

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Disclosed are methods for detecting multidrug resistance in neoplastic or damaged cells or multidrug resistant (MDR) neoplastic or damaged cells by detecting an increase in the cell surface expression of vimentin protein in such cells as compared to the level of cell surface expression of vimentin protein in a normal cell or a non-MDR neoplastic cell. The invention is based on the discovery that vimentin, a normally intracellular protein, is expressed in full length on the cell surface of neoplastic cells and damaged cells, and is expressed more abundantly on the cell surfaces of MDR neoplastic cells and MDR damaged cells. Although lower levels of vimentin are expressed on the cell surface of drug-sensitive neoplastic cells, vimentin is expressed in only negligible amts. on the cell surface of normal cells of the body. Thus, the invention allows the use of binding agents, to which are bound toxins or other therapeutic or diagnostic agents, that specifically bind to vimentin without detrimental side effects, since the only non-vimentin cells that are being killed are drug-sensitive neoplastic cells or damaged cells; normal cells remain unharmed. A vaccine for treating or preventing MDR neoplasm, comprising vimentin polypeptide, is also claimed. Provided are protein and cDNA sequences for human vimentin useful in vaccine preparation

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 97 THERE ARE 97 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 13 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:1019654 CAPLUS

DOCUMENT NUMBER: 142:5474

TITLE: Monoclonal antibodies that bind  $\alpha$ -folate receptor (FR- $\alpha$ ) tumor antigen and uses thereof in treatment of cancers expressing FR- $\alpha$

INVENTOR(S): Grasso, Luigi; Nicolaides, Nicholas C.; Sass, Philip M.  
 USA

PATENT ASSIGNEE(S):

SOURCE: U.S. Pat. Appl. Publ., 12 pp.  
 CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----
US 20040235108	A1	20041125	US 2004-851786	20040521 <--
AU 2004249673	A1	20041229	AU 2004-249673	20040521 <--
CA 2526647	A1	20041229	CA 2004-2526647	20040521 <--
WO 2004113388	A2	20041229	WO 2004-US16057	20040521 <--
WO 2004113388	A3	20050224		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK,				

EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE,  
SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE,  
SN, TD, TG

EP 1626991 A2 20060222 EP 2004-752961 20040521  
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
IE, SI, FI, RO, CY, TR, BG, CZ, EE, HU, PL, SK

JP 2007537709 T 20071227 JP 2006-533301 20040521

PRIORITY APPLN. INFO.: US 2003-4/2940P P 20030523

WO 2004-US16057 W 20040521

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Disclosed are monoclonal antibodies that specifically bind to the tetrameric form of the alpha-folate receptor (FR- $\alpha$ ) and not the monomeric form. The antibodies are useful in the treatment of certain cancers, particularly cancers that have increased cell surface expression of FR- $\alpha$ , such as ovarian cancer. Hybridoma cells expressing the monoclonal antibodies, antibody derivs., such as chimeric and humanized monoclonal antibodies, antibody fragments, mammalian cells expressing the monoclonal antibodies, derivs. and fragments, and methods of detecting and treating cancer using the antibodies, derivs., and fragments also are provided.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L21 ANSWER 14 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2004:1000292 CAPLUS

DOCUMENT NUMBER: 142:348265

TITLE: Multicenter pilot study of 5-fluorouracil, folinic acid, interferon alpha-2b and degradable starch microspheres via hepatic arterial infusion in patients with nonresectable colorectal liver metastases

AUTHOR(S): Pohlen, U.; Mansmann, U.; Berger, G.; Germer, C. T.; Gallkowski, U.; Boese-Landgraf, J.; Buhr, H. J.

CORPORATE SOURCE: Department of Surgery, Charite, Universitaetsmedizin Berlin, Germany

SOURCE: Anticancer Research (2004), 24(5B), 3275-3282

CODEN: ANTRD4; ISSN: 0250-7005

PUBLISHER: International Institute of Anticancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Background: It is necessary to establish therapeutic regimens for patients with nonresectable hepatic metastases of colorectal carcinoma.

A new regional chemotherapy regimen was tested in a prospective study in three centers. Patients and Methods: An arterial port system was

implanted in 95 patients. From Jan. 1994 to Mar. 1999, intra-arterial treatment was applied via the hepatic artery using 450 mg starch

microspheres with 5 million IU recombinant interferon- $\alpha$  2B, 500 mg/m<sup>2</sup> folinic acid and 600 mg/m<sup>2</sup> 5-FU body surface for 5 days

with a 14-day interval. Results: The tumor response rate was 70%. Median disease progression was 17 mo, median survival 24 mo. The subgroup anal.

shows a significant advantage (p < 0.0001) for patients with a liver tumor involvement of < 25% and a median survival of 39 mo compared to a

tumor involvement of 25 - 50% (24 mo) and > 50% (14 mo). Major toxicity problems were observed in 11%. However, there was no termination of therapy

on account of these problems. Conclusion: Intra-arterial chemotherapy with our new regimen was useful in patients with colorectal liver

metastases who had only an intrahepatic tumor burden of < 50%.

OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD  
(9 CITINGS)

REFERENCE COUNT: 55 THERE ARE 55 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT



L21 ANSWER 15 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2004:996296 CAPLUS  
 DOCUMENT NUMBER: 141:422031  
 TITLE: Multicellular compositions of pluripotent human embryonic stem cells and cancer cells for use in drug screening and testing  
 INVENTOR(S): Skorecki, Karl L.; Tzukerman, Maty  
 PATENT ASSIGNEE(S): Rappaport Family Institute for Research In the Medical Sciences, Israel  
 SOURCE: PCT Int. Appl., 62 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004099364	A2	20041118	WO 2004-IL375	20040505 <--
WO 2004099364	A3	20050707		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
EP 1627071	A2	20060222	EP 2004-731255	20040505
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR			
US 20070087435	A1	20070419	US 2006-555537	20061208
PRIORITY APPLN. INFO.:			IL 2003-155783	A 20030505
			WO 2004-IL375	W 20040505

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention discloses methods for producing novel multicellular comps. comprising cancer cells together with pluripotent human stem cells, which are capable of proliferating and differentiating into various normal cell lines and tissue structures. The present invention further discloses use of these novel multicellular systems for investigating the properties of cancer cells in a normal human tissue microenvironment, and for studying interventions that will modulate these properties including devising, testing and screening therapeutic drugs.

L21 ANSWER 16 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2004:947992 CAPLUS  
 DOCUMENT NUMBER: 142:232573  
 TITLE: A new superinvasive in vitro phenotype induced by selection of human breast carcinoma cells with the chemotherapeutic drugs paclitaxel and doxorubicin  
 AUTHOR(S): Glynn, S. A.; Gammell, P.; Heenan, M.; O'Connor, R.; Liang, Y.; Keenan, J.; Clynes, M.  
 CORPORATE SOURCE: National Institute for Cellular Biotechnology, Dublin City University, Dublin 9, Ire.  
 SOURCE: British Journal of Cancer (2004), 91(10), 1800-1807

PUBLISHER: Nature Publishing Group  
DOCUMENT TYPE: Journal  
LANGUAGE: English

CODEN: BJCAAI; ISSN: 0007-0920

AB Doxorubicin- and paclitaxel-selected variants of an in vitro invasive clonal population of the human breast cancer cell line, MDA-MB-435S, were established by pulse selection, and exhibited a novel superinvasive phenotype. This phenotype is characterized by an ability to relocate to another surface following invasion through matrigel and membrane pores, by decreased adhesion to extracellular matrix proteins and by increased motility. This may represent an in vitro model of a step in the metastatic process occurring subsequent to invasion. The paclitaxel-resistant variants, MDA-MB-435S-F/Taxol-10p and MDA-MB-435S-F/Taxol-10p4p were resistant to paclitaxel, vincristine and docetaxel, but not to doxorubicin, carboplatin, etoposide or 5-fluorouracil. The doxorubicin-selected variants MDA-MB-435S-F/Adr-10p and MDA-MB-435S-F/Adr-10p10p, in contrast, exhibited only small increases in resistance to doxorubicin, although they were slightly resistant to VP-16 and docetaxel, and exhibited increased sensitivity to paclitaxel, carboplatin and 5-fluorouracil.

OS.CITING REF COUNT: 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (22 CITINGS)  
REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 17 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:802607 CAPLUS

DOCUMENT NUMBER: 141:312949

TITLE: Anti-CD22 antibodies conjugated with cytotoxic drug for treating cancer, carcinoma, sarcoma and B cell lymphoma/leukemia

INVENTOR(S): Kunz, Arthur; Moran, Justin Keith; Rubino, Joseph Thomas; Jain, Neera; Vidunas, Eugene Joseph; Simpson, John Mclean; Merchant, Nishith; Dijoseph, John Francis; Ruppen, Mark Edward; Damle, Nitin Krishnaji; Robbins, Paul David; Popplewell, Andrew George  
PATENT ASSIGNEE(S): Wyeth Holdings Corporation, USA  
SOURCE: U.S. Pat. Appl. Publ., 90 pp., Cont.-in-part of U.S. Ser. No. 428,894.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040192900	A1	20040930	US 2003-699874	20031103 <--
US 20040082764	A1	20040429	US 2003-428894	20030502 <--
AU 2009202609	A1	20090716	AU 2009-202609	20090626
PRIORITY APPLN. INFO.:			US 2002-377440P	P 20020502
			US 2003-428894	A2 20030502
			AU 2003-231293	A3 20030502

AB Methods for preparing monomeric cytotoxic drug/carrier conjugates with a drug loading significantly higher than in previously reported procedures and with decreased aggregation and low conjugate fraction (LCF) are described. Cytotoxic drug derivative/antibody conjugates, compns. comprising the conjugates and uses of the conjugates are also described. Monomeric calicheamicin derivative/anti-CD22 antibody conjugates, compns. comprising the conjugates and uses of the conjugates are also described. The anti-CD22 antibody is a monoclonal antibody, human antibody, chimeric antibody,

humanized antibody or fragment. The cytotoxic drug is a calicheamicin, thiotepa, taxane, vincristine, daunorubicin, doxorubicin, epirubicin, esperamicin, actinomycin, anthramycin, azaserine, bleomycin, tamoxifen, idarubicin, etc.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L21 ANSWER 18 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:759839 CAPLUS

DOCUMENT NUMBER: 141:254551

TITLE: Methods and compositions to determine the chemosensitizing dose of suramin used in combination therapy

INVENTOR(S): Au, Jessie L. S.; Wientjes, M. Guillaume

PATENT ASSIGNEE(S): USA

SOURCE: U.S. Pat. Appl. Publ., 19 pp., Cont.-in-part of Appl. No. PCT/US02/30210.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040180955	A1	20040916	US 2004-807620	20040324 <--
WO 2003026574	A2	20030403	WO 2002-US30210	20020924 <--
WO 2003026574	A3	20040415		

W: AE, AG, AL, AU, BA, BB, BG, BR, BZ, CA, CN, CO, CR, CU, CZ, DM, DZ, EC, EE, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KP, KR, KZ, LC, LK, LR, LS, LT, LV, MA, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, RO, SD, SG, SI, SK, SL, TN, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW

RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

PRIORITY APPLN. INFO.: US 2001-324704P P 20010924  
WO 2002-US30210 A2 20020924

AB A method for determining a therapeutically effective amount of suramin for administering to a patient, who is to receive a cytotoxic agent, which comprises the steps of determining the circulating suramin concentration in the patient; administering suramin, if required, to establish a low circulating concentration of suramin in the patient of below about 200  $\mu$ M; and administering the chemotherapeutic agent to the patient when the low circulating concentration of suramin is present in the patient. Conveniently a nomogram can be constructed for use in clin. settings with the suramin.

L21 ANSWER 19 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:662225 CAPLUS

DOCUMENT NUMBER: 142:126720

TITLE: Inhibition of cell survival and invasive potential of colorectal carcinoma cells by the tyrosine kinase inhibitor STI571

AUTHOR(S): Bellone, Graziella; Ferrero, Dario; Carbone, Anna; De Quadros, Marlene R.; Gramigni, Claudia; Prati, Adriana; Davidson, William; Mioli, Pierroberto; Dughera, Luca; Emanuelli, Giorgio; Rodeck, Ulrich

CORPORATE SOURCE: Department of Clinical Physiopathology, University of Torino, Turin, Italy

SOURCE: Cancer Biology & Therapy (2004), 3(4),

385-392

CODEN: CBTAOO; ISSN: 1538-4047

Landes Bioscience

PUBLISHER:

DOCUMENT TYPE:

LANGUAGE:

Journal

English

AB Inhibiting tyrosine kinases has recently emerged as a therapeutic modality in several forms of neoplasia. The tyrosine kinase inhibitor STI571 (IMATINIB MESYLATE; GLEEVEC; GLIVEC) is a case in point as it has shown promise in the treatment of malignancies expressing the BCR/ABL fusion protein. In addition to BCR/ABL, STI571 inhibits the tyrosine kinase moieties of several cell surface receptors including the platelet-derived growth factor (PDGF) receptors and c-Kit. Previous work demonstrated that c-Kit activation supports migration, invasion and, survival of certain colorectal carcinoma cells including DLD-1. Here we describe that blocking c-Kit with STI571 inhibits these malignant traits not only in DLD-1 cells but also in two early passage colorectal carcinoma cell strains. Specifically, STI571 inhibited anchorage-independent colony formation and cell scattering in semi-solid medium. Furthermore, it enhanced apoptosis susceptibility and abrogated invasion of DLD-1 cells through Matrigel. In addition, STI571 treatment affected the balance of the Bcl-2 family of apoptosis regulators on favor of a pro-apoptotic phenotype. Specifically, STI571 treatment of DLD-1 cells was associated with lower levels of Bcl-2 expression accompanied by de novo expression of Bcl-xS. Finally, STI571 acted as a chemosensitizing agent in DLD-1 cells when used in combination with 5-fluorouracil.

OS.CITING REF COUNT: 16 THERE ARE 16 CAPLUS RECORDS THAT CITE THIS RECORD (16 CITINGS)

REFERENCE COUNT: 57 THERE ARE 57 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 20 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:633479 CAPLUS

DOCUMENT NUMBER: 141:162388

TITLE: Modified polysaccharides combination with anti-cancer drugs for enhanced treatment of cancer

INVENTOR(S): Platt, David

PATENT ASSIGNEE(S): Pro-Pharmaceuticals Inc, USA

SOURCE: PCT Int. Appl., 28 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004064777	A2	20040805	WO 2004-US747	20040114 <--
WO 2004064777	A3	20050909		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI				
EP 1592432	A2	20051109	EP 2004-702115	20040114
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2006515647	T	20060601	JP 2006-500921	20040114
US 20050282773	A1	20051222	US 2005-182096	20050715
PRIORITY APPLN. INFO.:			US 2003-440496P	P 20030116
			WO 2004-US747	W 20040114

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Modified polysaccharide compns. and their use in combination with an

anticancer drug for treating subjects with cancer, reduce toxicity and inhibit metastasis, are described. The modified polysaccharide includes a saccharide backbone being <5% esterified and containing repeating units, wherein each repeating unit has a plurality of uronic acid mols., each repeating unit having at least one neutral monosaccharide attached thereto, at least one side chain of saccharides attached to the backbone further comprising a plurality of neutral saccharides or saccharide derivs.; and having an average mol. weight in the range of 15 to 60 kD. The polysaccharide when combined with the chemotherapeutic drug behaves as a delivery vehicle, which pos. enhance the chemotherapeutic effect while reducing side effects.

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 21 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2004:612470 CAPLUS

DOCUMENT NUMBER: 141:153495

TITLE: Methods of labeling proteins exposed to the luminal surface of a vessel in the identification of proteins for targeting drug delivery

INVENTOR(S): Roben, Paul; Stevens, Anthony C.

PATENT ASSIGNEE(S): Utah Ventures II L.P., USA

SOURCE: U.S. Pat. Appl. Publ., 123 pp., Cont.-in-part of Appl. No. PCT/US03/10195.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040146516	A1	20040729	US 2004-794899	20040305 <--
US 6903196	B1	20050607	US 2000-528742	20000320
US 20030021792	A1	20030130	US 2002-165603	20020607 <--
WO 2003084469	A2	20031016	WO 2003-US10195	20030331 <--
WO 2003084469	A3	20040610		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
WO 2005086775	A2	20050922	WO 2005-US7415	20050307
WO 2005086775	A3	20090409		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG, AP, EA, EP, OA				

PRIORITY APPLN. INFO.: US 1999-139579P P 19990617

US 2000-528742	A2 20000320
US 2001-297021P	P 20010608
US 2001-305117P	P 20010712
US 2002-369452P	P 20020401
US 2002-165603	A2 20020607
WO 2003-US10195	A2 20030331
US 2004-794899	A 20040305

AB Reagents that can be used to label proteins exposed on the luminal surface of an anatomical structure are identified. The proteins identified by these reagents may be used as affinity targets for the cell- or tissue-specific delivery of drugs. The method uses labeling reagents that do not pass through biol. membranes. They have a domain that binds or reacts relatively non-specifically to proteins and that is connected to a reporter group by a linker that is labile to non-denaturing reducing conditions. The labeled proteins can then be identified in homogenates. Use of the method to identify proteins of the lumina of several organs of rat is demonstrated. Use of two of these proteins, dipeptidyl peptidase IV and Thy-1 antigen, to direct transcytosis of antibodies in lung is demonstrated. Antibodies to the proteins were transported from the luminal space of the blood vessels of the lung across the endothelium. Similarly, conjugates of antibodies and antibiotics or antineoplastic drugs could also be transported by transcytosis.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L21 ANSWER 22 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:589740 CAPLUS

DOCUMENT NUMBER: 141:134063

TITLE: HSC70 directed diagnostics and therapeutics for multidrug resistant (MDR) neoplastic disease, using HSC70-binding agents along with detecting other MDR markers

INVENTOR(S): Georges, Elias; Serfass, Lucille; Bonneau, Anne-Marie; Dallaire, Frederic

PATENT ASSIGNEE(S): Aurelium Biopharma Inc., Can.

SOURCE: PCI Int. Appl., 166 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004061458	A2	20040722	WO 2003-IB6416	20031215 <--
WO 2004061458	A3	20041125		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2512513	A1	20040722	CA 2003-8	20031215 <--
AU 2003300676	A1	20040729	AU 2003-300676	20031215 <--
US 20040185511	A1	20040923	US 2003-737350	20031215 <--
US 7226748	B2	20070605		
EP 1588162	A2	20051026	EP 2003-814521	20031215

EP 1588162 B1 20080917  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK  
 JP 2006512579 T 20060413 JP 2004-564385 20031215  
 AT 408838 T 20081015 AT 2003-814521 20031215  
 US 20090004102 A1 20090101 US 2007-801415 20070509  
 PRIORITY APPLN. INFO.: US 2003-438012P P 20030103  
 US 2003-737350 A3 20031215  
 WO 2003-IB6416 W 20031215

AB The invention is based upon the discovery that HSC70 (heat shock cognate protein 70), a normally intracellular protein, is expressed in full length on the cell surface of neoplastic cells and damaged cells, and is expressed more abundantly on the cell surfaces of multidrug resistant (MDR) neoplastic cells and MDR damaged cells. Although lower levels of HSC70 are expressed on the cell surface of drug-sensitive neoplastic cells, HSC70 is expressed in only negligible amts. on the cell surface of normal cells of the body. Disclosed are methods for detecting neoplastic or damaged cells and for detecting multidrug resistance in neoplastic or damaged cells by detecting an increase of HSC70 expression on the surface of such a multidrug resistant neoplastic or damaged cells as compared to the level of expression of the HSC70 on the surface of a normal cell. Thus, the invention allows the use of binding agents, to which are bound toxins or other therapeutic or diagnostic agents, that specifically bind to HSC70 without detrimental side effects, since the only non-HSC70 cells that are being killed are drug-sensitive neoplastic cells or damaged cells; normal cells remain unharmed. A vaccine for treating or preventing MDR neoplasms, comprising HSC70 polypeptide, is also claimed.

OS.CITING REF COUNT: 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 23 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:534403 CAPLUS

DOCUMENT NUMBER: 141:66306

TITLE: Nucleophosmin directed diagnostics and therapeutics for multidrug resistant neoplastic disease  
 INVENTOR(S): Georges, Elias; Serfass, Lucile; Bonneau, Anne-Marie; Dallaire, Frederic

PATENT ASSIGNEE(S): Aurelium Biopharma Inc., Can.

SOURCE: PCT Int. Appl., 165 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004055517	A2	20040701	WO 2003-IB6445	20031215 <--
WO 2004055517	A3	20051027		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK,			

TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG  
 CA 2509902 A1 20040701 CA 2003-2509902 20031215 <--  
 AU 2003302986 A1 20040709 AU 2003-302986 20031215 <--  
 US 20050009119 A1 20050113 US 2003-737712 20031215  
 US 7413851 B2 20080819

PRIORITY APPLN. INFO.: US 2002-433351P P 20021213  
 WO 2003-1B6445 W 20031215

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention provides methods for detecting multidrug resistance in neoplastic or damaged cells by measurement of cell surface expression of a nucleophosmin (NPM) protein. The patent also encloses vaccine or binding agents that specifically bind to nucleophosmin as therapeutic components.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 24 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:493868 CAPLUS

DOCUMENT NUMBER: 141:52866

TITLE: A variant of a single-chain antibody to p97 melanotransferrin with increased stability for use in diagnosis and therapy of melanoma

INVENTOR(S): McDonagh, Charlotte F.; Francisco, Joseph A.

PATENT ASSIGNEE(S): Seattle Genetics, Inc., USA

SOURCE: PCT Int. Appl., 71 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004050867	A1	20040617	WO 2002-US38414	20021202 <--
W: CA, US				
US 20060160174	A1	20060720	US 2005-537143	20051024
PRIORITY APPLN. INFO.:			WO 2002-US38414	W 20021202

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB A variant of the L49 single chain antibody (L49-sFv) to p97 melanotransferrin that shows increased refolding efficiency and greater stability in mouse serum, and substantially maintaining binding affinity for p97 melanotransferrin is described. P97 melanotransferrin is expressed on the surface of a number of types of cancer (carcinoma) cells, e.g., melanoma cells, lung cancer cells, renal cancer cells, colon cancer cells, and so may be useful in diagnosis and therapy. The present invention also relates to a modified L49-sFv fused or conjugated to a therapeutic agent, such as a cytotoxic mol. or a pro-drug converting enzyme. The present invention also relates to methods of using the modified L49-sFv mols. fused or conjugated to a therapeutic agent for treatment and/or prophylaxis of cancer, which cancer cells express p97 melanotransferrin. Unusual amino acids predicted to affect stability of the antibody were identified by sequence alignment. These amino acids were substituted with the most common amino acids at these sites and the ability of the substitution variant to bind the antigen was tested. Substitution of three amino acids in the VH region led to the most complete refolding of the antibody. The substitution variants showed binding affinities for melanotransferrin comparable to those of the original single-chain antibody.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS



## RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 25 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:441038 CAPLUS

DOCUMENT NUMBER: 141:405731

TITLE: Induction of a multifactorial resistance phenotype by high paclitaxel selective pressure in a human ovarian carcinoma cell line

AUTHOR(S): Violini, S.; D'Ascenzo, S.; Bagnoli, M.; Millimaggi, D.; Miotti, S.; Canevari, S.; Pavan, A.; Dolo, V.

CORPORATE SOURCE: Dipartimento di Medicina Sperimentale, Università di L'Aquila, L'Aquila, Italy

SOURCE: Journal of Experimental &amp; Clinical Cancer Research (2004), 23(1), 83-91

CODEN: JECRDN; ISSN: 0392-9078

PUBLISHER: Regina Elena Institute for Cancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Paclitaxel (PTX) is a potent anti-neoplastic agent that is highly effective in treating ovarian cancer. Nevertheless, the emergence of PTX resistance has limited the control of this disease. To gain insight into the mol. alterations accompanying drug resistance in ovarian cancer, we generated a new stable PTX-resistant ovarian carcinoma cell line. CABA I cells, which display an intrinsic PTX resistance (IC50 = 800 ng/mL), were subjected to continuous exposure to PTX. From the residual surviving cells, the highly PTX-resistant line CABA-PTX (IC50 = 256000 ng/mL) was generated and stably maintained in vitro. Anal. of  $\beta$ -tubulin expression indicated that only the HM40 and HB9 isotypes were expressed in both parental and resistant cells. No specific point mutations in the HM40 were detected in either cell line, but expression levels of this isotype were significantly reduced (40%) in CABA-PTX cells. HB9 levels were unchanged. In those cells, PTX resistance was associated with cross-resistance to vinblastine but not to methotrexate or 5-fluorouracil. Verapamil treatment did not reverse the intrinsic drug resistance of parental cells, but partially modulated the sensitivity of CABA-PTX cells to PTX and induced total sensitivity to vinblastine. No changes in the cell surface expression of the drug efflux pumps MRP1, MRP2 and P-glycoprotein were observed. PTX influx, monitored using a fluorescent drug derivative, was significantly reduced and delayed in CABA-PTX cells as compared to the parental cells. Together, these findings suggest that more than one mechanism is involved in PTX resistance, making CABA-PTX cell line a potentially valuable in vitro tool to study multifactorial acquired drug resistance in ovarian cancer.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 26 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:420935 CAPLUS

DOCUMENT NUMBER: 141:420107

TITLE: Randomized Trial of Intraperitoneal and/or Systemic Adjuvant Chemotherapy in Patients With Colon Carcinoma

AUTHOR(S): Labianca, Roberto; Fossati, Roldano; Zaniboni, Alberto; Torri, Valter; Marsoni, Silvia; Nitti, Donato; Boffi, Lamberto; Scatizzi, Marco; Tardio, Bernardino; Mastrodonato, Nicola; Banducci, Stefano; Consani, Giampiero; Pancera, Gianfranco

CORPORATE SOURCE: Unita Operativa di Oncologia Medica, Bergamo, Italy

SOURCE: Journal of the National Cancer Institute (2004)

), 96(10), 750-758  
CODEN: JNCIEQ; ISSN: 0027-8874  
Oxford University Press  
Journal  
English

PUBLISHER:  
DOCUMENT TYPE:  
LANGUAGE:

AB Background: 5-Fluorouracil-based adjuvant chemotherapy after surgical resection of colon cancer is standard treatment. However, the choice of best delivery route-i.e., systemic (i.e., i.v. or oral) or regional (i.e., intraportal, i.p., or hepatic arterial infusion)-has been controversial. In a randomized clin. trial of patients with colon cancer, we compared the benefits of chemotherapy delivered by these routes individually or in combination. Methods: From Apr. 2, 1992, through Apr. 30, 1998, 1084 eligible patients with Dukes' stage B or C colon carcinoma were randomly assigned: 369 patients to the IP regimen (continuous portal vein infusion of 5-fluorouracil at 500 mg/m<sup>2</sup> of body surface daily and heparin at 5000 IU daily for 7 consecutive days, beginning on the day of surgery), 358 patients to the SY regimen (six 28-day courses of systemic leucovorin at 100 mg/m<sup>2</sup> daily on days 1 through 5 followed by systemic bolus 5-fluorouracil at 370 mg/m<sup>2</sup> daily on days 1 through 5, with treatment initiated 15-35 days after surgery), and 357 patients to the IP+SY regimen (the IP regimen followed by the SY regimen, with the same scheduling). Primary survival was analyzed with the log-rank statistic and a Cox multivariable regression model. All statistical tests were two sided. Results: At a median follow-up time of 99 mo, 389 events (recurrences, second malignancies, or deaths) had occurred, and 361 patients died. Sites of first recurrences were similar among the three arms. At 5 yr, overall and event-free survival rates were similar among those on the IP (74% and 68%, resp.), SY (78% and 71%), and IP+SY (73% and 67%) regimens. When compared with the group on the SY regimen, the risk for death associated with the IP regimen (hazard ratio [HR] = 1.05, 95% confidence interval [CI] = 0.82 to 1.36) was similar to that associated with the IP+SY regimen (HR = 1.12, 95% CI = 0.78 to 1.45) (P = .69), as were the risks for first event (HR = 1.07, 95% CI = 0.84 to 1.37 and HR = 1.10, 95% CI = 0.86 to 1.41, resp.) (P = .74). Conclusion: Overall and event-free survival rates were similar in all three arms. The combined regimen was no better than either single regimen alone.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)  
REFERENCE COUNT: 23 THERE ARE 23 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 27 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2004:420314 CAPLUS

DOCUMENT NUMBER: 141:64564

TITLE: Pharmacoproteomic analysis of prechemotherapy and postchemotherapy plasma samples from patients receiving neoadjuvant or adjuvant chemotherapy for breast carcinoma

AUTHOR(S): Pusztai, Lajos; Gregory, Betsy W.; Baggerly, Keith A.; Peng, Bo; Koomen, John; Kuerer, Henry M.; Esteva, Francisco J.; Symmans, W. Fraser; Wagner, Peter; Hortobagyi, Gabriel N.; Laronga, Christine; Semmes, O. John; Wright, George L., Jr.; Drake, Richard R.; Vlahou, Antonia

CORPORATE SOURCE: Department of Breast Medical Oncology, The University of Texas M. D. Anderson Cancer Center, Houston, TX, USA

SOURCE: Cancer (New York, NY, United States) (2004), 100(9), 1814-1822

PUBLISHER: CODEN: CANCAR; ISSN: 0008-543X  
John Wiley & Sons, Inc.

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB In this study, proteomic changes were examined in response to paclitaxel chemotherapy or 5-fluorouracil, doxorubicin, and cyclophosphamide (FAC) chemotherapy in plasma from patients with Stage I-III breast carcinoma. The authors also compared the plasma profiles of patients with cancer with the plasma profiles of healthy women to identify breast carcinoma-associated protein markers. Sixty-nine patients and 15 healthy volunteers participated in the study. Plasma was sampled on Day 0 before chemotherapy and on Day 3 posttreatment in the 69 patients or 3 days apart in the 15 healthy women. Twenty-nine patients received preoperative chemotherapy, and 40 received postoperative chemotherapy. Surface-enhanced laser desorption/ionization mass spectrometry was used to generate protein mass profiles. Few changes were observed in plasma during treatment. Only 1 protein peak was identified (mass/charge ratio [m/z], 2790) that was induced by paclitaxel and, to a lesser extent, by FAC chemotherapy. This proteomic response was detectable in 80% of patients who were treated preoperatively but also was present with lesser intensity in approx. 40% of patients treated postoperatively. There was no clear correlation between induction of m/z 2790 during a single course of treatment and final tumor response to preoperative chemotherapy. Five other peaks also were identified that discriminated between plasma from patients with breast carcinoma and plasma from normal women. These same peaks also were detectable in a subset of patients who already had undergone surgery to remove their tumors. A single chemotherapy-inducible SELDI-MS peak and five other peaks that distinguished plasma obtained from patients with breast carcinoma from plasma obtained from normal, healthy women were identified. The (as yet unsequenced) proteins represented by these peaks are candidate markers of micrometastatic disease after surgery.

OS.CITING REF COUNT: 57 THERE ARE 57 CAPLUS RECORDS THAT CITE THIS RECORD (57 CITINGS)  
REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 28 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:269986 CAPLUS

DOCUMENT NUMBER: 140:301820

TITLE: CD43 as a tumor marker, particularly ovarian tumor marker, and methods for diagnosing and treating tumors and suppressing CD antigen promoters

INVENTOR(S): Shelley, Carl Simon; Farokhzad, Omid C.

PATENT ASSIGNEE(S): The General Hospital Corporation, USA

SOURCE: PCT Int. Appl., 133 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004026120	A2	20040401	WO 2003-US30213	20030923 <--
WO 2004026120	A3	20040819		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY,			

KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,  
 FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR,  
 BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG

AU 2003278918 A1 20040408 AU 2003-278918 20030923 <--  
 US 20060216231 A1 20060928 US 2006-528948 20060421  
 PRIORITY APPLN. INFO.: US 2002-412964P P 20020923  
 WO 2003-US30213 W 20030923

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Methods of treating tumors, reducing white blood cell nos., and inhibiting CD antigen promoters are provided. The present invention is based, in part, on the discovery that CD43 (leukosialin) plays a role in the diagnosis and treatment of tumors. The invention is also based, in part, on the discovery that ovarian tumor cells abnormally express CD43 on their surfaces. The invention is also based, in part, on the finding that CD43 inhibitors repress the CD43 promoter which mediates progression of tumors and promotes survival or proliferation of white blood cells. It was shown that hnRNP-K and Pura protein act together to repress transcriptional activity of CD43 gene promoter. In one embodiment, CD43 monoclonal antibodies, B51, MEM-59, 84-3C 1, Bra7G, DF-T1, 1G10, MT1, L10, L14, T2/53, B1-B6, L60, BL-GCE/G3, 6E5, 6F5, 10G7, G10-2, G19-1, DS 1.C1, L66, CBF.78, 148.1B6, 148.1C3, 148.3D4, 161.46, RDP.AD9, OH.01, HI165, and HI161, suggested for diagnosis and therapy.

L21 ANSWER 29 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2004:100803 CAPLUS  
 DOCUMENT NUMBER: 140:139483  
 TITLE: Method for enhancing the effectiveness of therapies of hyperproliferative diseases  
 INVENTOR(S): Chang, Yan; Sasak, Vodek  
 PATENT ASSIGNEE(S): USA  
 SOURCE: U.S. Pat. Appl. Publ., 14 pp., Cont.-in-part of U.S. Ser. No. 176,235.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 3  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20040023925	A1	20040205	US 2003-408723	20030407 <--
US 20030013681	A1	20030116	US 2002-176235	20020620 <--
US 6680306	B2	20040120		
CN 1543351	A	20041103	CN 2002-816003	20020621 <--
CN 100558367	C	20091111		
US 20040043962	A1	20040304	US 2003-657383	20030908 <--
AU 2004229399	A1	20041028	AU 2004-229399	20040407 <--
CA 2521649	A1	20041028	CA 2004-2521649	20040407 <--
WO 2004091634	A1	20041028	WO 2004-US10675	20040407 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: BW, GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
EP 1617849	A1	20060125	EP 2004-759200	20040407

EP 1617849 B1 20080618  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, PL, SK, HR  
 JP 2006522163 T 20060928 JP 2006-59773 20040407  
 AT 398458 T 20080715 AT 2004-759200 20040407  
 EP 1980257 A1 20081015 EP 2008-10897 20040407  
 R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE,  
 IT, LI, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR, AL, HR, LT, LV, MK  
 PRIORITY APPLN. INFO.:  
 US 2001-299991P P 20010621  
 US 2002-176235 A2 20020620  
 US 2003-408723 A 20030407  
 US 2003-461006P P 20030407  
 US 2003-474562P P 20030530  
 EP 2004-759200 A3 20040407  
 WO 2004-US10675 W 20040407

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The efficacy of conventional cancer therapies such as surgery,  
 chemotherapy and radiation is enhanced by the use of a therapeutic  
 material which binds to and interacts with galectins. The therapeutic  
 material can enhance apoptosis thereby increasing the effectiveness of  
 oncolytic agents. It can also inhibit angiogenesis thereby moderating  
 tumor growth and/or metastasis.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD  
 (2 CITINGS)

L21 ANSWER 30 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:60636 CAPLUS

DOCUMENT NUMBER: 140:105262

TITLE: Ciclopirox and analogs thereof with optional  
 antiproliferative agents for the treatment of  
 neoplasms

INVENTOR(S): Lee, Margaret S.; Keith, Curtis; Auspitz, Benjamin A.;  
 Zimmermann, Grant R.; Nichols, M. James

PATENT ASSIGNEE(S): Combinatorx, Incorporated, USA

SOURCE: PCT Int. Appl., 51 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007676	A2	20040122	WO 2003-US21783	20030714 <--
WO 2004007676	A3	20040408		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, ND, TD, TG			
AU 2003251875	A1	20040202	AU 2003-251875	20030714 <--
IN 2004CN02749	A	20060210	IN 2004-CN2749	20041206
PRIORITY APPLN. INFO.:			US 2002-396120P	P 20020715
			US 2002-400905P	P 20020802
			WO 2003-US21783	W 20030714

OTHER SOURCE(S): MARPAT 140:105262

AB The invention features a method for treating a patient having a cancer or other neoplasm, by administering to the patient (i) ciclopirox or a structural or functional analog thereof; and optionally (ii) an antiproliferative agent simultaneously or within 14 days of each other in amts. sufficient to inhibit the growth of the neoplasm.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 31 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:60538 CAPLUS

DOCUMENT NUMBER: 140:105261

TITLE: Activated checkpoint therapy and methods of use thereof

INVENTOR(S): Li, Chiang J.; Li, You-Zhi; Pardee, Arthur B.

PATENT ASSIGNEE(S): Cyclis Pharmaceuticals, Inc., USA; Dana-Farber Cancer Institute, Inc.; Beth Israel Deaconess Medical Center PCT Int. Appl., 61 pp.

SOURCE: CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 13

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004007531	A2	20040122	WO 2003-US22631	20030717 <--
WO 2004007531	A3	20040812		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2492772	A1	20040122	CA 2003-2492772	20030717 <--
AU 2003254029	A1	20040202	AU 2003-254029	20030717 <--
EP 1545507	A2	20050629	EP 2003-764816	20030717
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
JP 2005538981	T	20051222	JP 2004-521982	20030717
CA 2506340	A1	20040603	CA 2003-2506340	20031118 <--
EP 1567515	A2	20050831	EP 2003-786941	20031118
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK			
BR 2003016296	A	20051213	BR 2003-16296	20031118
CN 1729183	A	20060201	CN 2003-80106617	20031118
JP 2006508147	T	20060309	JP 2004-553999	20031118
PRIORITY APPLN. INFO.:			US 2002-396360P	P 20020717
			US 2002-427283P	P 20021118
			US 2003-622854	A 20030717
			WO 2003-US22631	W 20030717
			WO 2003-US37219	W 20031118
AB	Disclosed herein are novel methods and compns. for Activated Checkpoint TherapyTM. Also disclosed are methods of treating cancer and apoptosis-associated disorders using cell cycle checkpoint activation modulators. The invention further discloses methods for screening for			

cell cycle checkpoint activation modulators and the cell cycle checkpoint activation modulators identified by those screening methods.  
 OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 32 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2004:41226 CAPLUS  
 DOCUMENT NUMBER: 140:105321  
 TITLE: Methods and compositions relating to isoleucine boroproline compounds  
 INVENTOR(S): Adams, Sharlene; Miller, Glenn T.; Jesson, Michael I.; Jones, Barry  
 PATENT ASSIGNEE(S): Point Therapeutics, Inc., USA  
 SOURCE: PCT Int. Appl., 152 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2004004658	A2	20040115	WO 2003-US21405	20030709 <--
WO 2004004658	A3	20050804		
W:				
AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZW				
RW:				
GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2491466	A1	20040115	CA 2003-2491466	20030709 <--
AU 2003265264	A1	20040123	AU 2003-265264	20030709 <--
US 20040077601	A1	20040422	US 2003-616694	20030709 <--
US 20050084490	A1	20050421	US 2003-616409	20030709
EP 1578434	A2	20050928	EP 2003-763380	20030709
R:				
AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2006507352	T	20060302	JP 2004-562634	20030709
CN 1802090	A	20060712	CN 2003-821282	20030709
CN 1826129	A	20060830	CN 2003-821281	20030709
IN 2005KN00151	A	20050916	IN 2005-KN151	20050208
PRIORITY APPLN. INFO.:				
			US 2002-394856P	P 20020709
			US 2002-414978P	P 20021001
			US 2003-466435P	P 20030428
			WO 2003-US21405	W 20030709

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 140:105321

AB A method for treating subjects with, inter alia, abnormal cell proliferation or infectious disease using agents of formula (I), AmNHCH(CH(CH3)CH2CH3)COA1R (where Am and A1 are amino acids and R = organo boronates, organo phosphonates, fluoroalkyl ketones, aliphatic ketones, N-peptidyl-O-(acylhydroxylamines), azapeptides, azetidines, fluoroolefins, dipeptide isosteres, peptidyl ( $\alpha$ -aminoalkyl) phosphonate esters, aminoacyl pyrrolidine-2-nitriles and 4-cyanothiazolidides) is claimed. Methods for stimulating an immune response using the compds. of the

invention are also claimed. Compns. containing Ile-boroPro compds. are also provided as are kits containing the compns. The invention embraces the use of these compds. alone or in combination with other therapeutic agents.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)  
REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 33 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:892567 CAPLUS

DOCUMENT NUMBER: 139:386334

TITLE: Production of monomeric calicheamicin derivative  
cytotoxic drug/carrier conjugates

INVENTOR(S): Kunz, Arthur; Moran, Justin Keith; Rubino, Joseph  
Thomas; Jain, Neera; Vidunas, Eugene Joseph; Simpson,  
John McLean; Robbins, Paul David; Merchant, Nishith;  
Dijoseph, John Francis; Ruppen, Mark Edward; Damle,  
Nitin Krishnaji; Popplewell, Andrew George; et al.

PATENT ASSIGNEE(S): Wyeth Holdings Corporation, USA

SOURCE: PCT Int. Appl., 186 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003092623	A2	20031113	WO 2003-US13910	20030502 <--
WO 2003092623	A3	20040318		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2483552	A1	20031113	CA 2003-2483552	20030502 <--
AU 2003231293	A1	20031117	AU 2003-231293	20030502 <--
EP 1507556	A2	20050223	EP 2003-724432	20030502
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, HU, SK				
JP 2005524700	T	20050818	JP 2004-500808	20030502
CN 1665532	A	20050907	CN 2003-815260	20030502
CN 100482277	C	20090429		
BR 2003009868	A	20051018	BR 2003-9868	20030502
NO 2004004663	A	20050125	NO 2004-4663	20041028
MX 2004010792	A	20050307	MX 2004-10792	20041029
IN 2004KN01802	A	20060106	IN 2004-KN1802	20041129
IN 2007KN01141	A	20080801	IN 2007-KN1141	20070402
AU 2009202609	A1	20090716	AU 2009-202609	20090626
PRIORITY APPLN. INFO.:			US 2002-377440P	P 20020502
			AU 2003-231293	A3 20030502
			WO 2003-US13910	W 20030502
			IN 2004-KN1802	A3 20041129

AB The present invention relates to methods for the production of monomeric cytotoxic drug/carrier conjugates (the "conjugates") with higher drug loading and substantially reduced low conjugate fraction (LCF). Cytotoxic



drug derivative/antibody conjugates, compns. comprising the conjugates and uses of the conjugates are also described. Particularly, the invention relates to anti-CD22 antibody-monomeric calicheamicin conjugates. The invention also relates to the conjugates of the invention, to methods of purification of the conjugates, to pharmaceutical compns. comprising the conjugates, and to uses of the conjugates.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)  
REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 34 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:813738 CAPLUS

DOCUMENT NUMBER: 140:368192

TITLE: Inhibition of primary colon carcinoma growth and liver metastasis by the A3 adenosine receptor agonist CF101

AUTHOR(S): Ohana, G.; Bar-Yehuda, S.; Arich, A.; Madi, L.; Dreznick, Z.; Rath-Wolfson, L.; Silberman, D.; Slosman, G.; Fishman, P.

CORPORATE SOURCE: Rabin Medical Center, Department of Surgery A/B, Sackler Faculty of Medicine Tel-Aviv University, Petach-Tikva, 49100, Israel

SOURCE: British Journal of Cancer (2003), 89(8), 1552-1558

CODEN: BJCAAI; ISSN: 0007-0920

PUBLISHER: Nature Publishing Group

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Adenosine is a purine nucleoside that acts as a regulatory mol. by binding to specific G-protein-coupled A1, A2A, A2B, and A3 cell surface receptors. We have recently demonstrated that adenosine inhibits tumor cell growth and concomitantly stimulates bone marrow cell proliferation via activation of the A3 adenosine receptor (A3AR). In the present study, we show that a synthetic agonist to the A3AR, CF101, at the low nanomolar concentration range, inhibits HCT-116 human colon carcinoma cell growth. This effect was reversed by the selective A3AR antagonist MRS1523, demonstrating the specificity of the response. CF101 (given orally) was efficacious in inhibiting the development of primary tumors in xenograft and syngeneic models in which mice were inoculated s.c. with human HCT-116 or murine CT-26 colon carcinoma cells, resp. Moreover, CF101 suppressed (50%,  $P < 0.01$ ) colon cancer liver metastases in syngeneic mice inoculated to the spleen with CT-26 cells. The mechanism of action entailed upregulation of interleukin-12 production in the CF101-treated groups and potentiation of NK cell activity. In the HCT-116 xenograft model in which a combined therapy of CF101 and 5-fluorouracil (5-FU) was examined, an additive antitumor effect was demonstrated. Moreover, CF101 prevented the 5-FU-induced myelotoxicity, resulting in normal values of white blood cell and neutrophil counts. We conclude that the A3AR agonist CF101, a small orally bioavailable mol., exerts systemic anticancer, antimetastatic, and myeloprotective effects in colon carcinoma-bearing mice, and may serve as an adjuvant treatment to enhance the chemotherapeutic index and prevent myelotoxicity.

OS.CITING REF COUNT: 25 THERE ARE 25 CAPLUS RECORDS THAT CITE THIS RECORD (25 CITINGS)

REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 35 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:780113 CAPLUS

DOCUMENT NUMBER: 140:104669

TITLE: Mislocalization of membrane proteins associated with multidrug resistance in cisplatin-resistant cancer cell lines

AUTHOR(S): Liang, Xing-Jie; Shen, Ding-Wu; Garfield, Susan; Gottesman, Michael M.

CORPORATE SOURCE: National Cancer Institute, Laboratory of Cell Biology, National Institutes of Health, Bethesda, MD, 20892-4254, USA

SOURCE: Cancer Research (2003), 63(18), 5909-5916  
CODEN: CNREA8; ISSN: 0008-5472

PUBLISHER: American Association for Cancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The accumulation of [14C]carboplatin and [3H]methotrexate is reduced in single-step KB epidermoid adenocarcinoma (KB-CP) cells, which are cross-resistant to carboplatin, methotrexate, and sodium arsenite. In these KB-CP cells, multidrug resistance is accompanied by mislocalization of multidrug resistance associated protein (MRP) 1 and other membrane proteins such as folate-binding protein. MRP1 was not decreased in amount in single-step variants but accumulates in a cytoplasmic fraction, and its apparent mol. weight was altered probably because of reduced glycosylation in resistant cells. This low-d. compartment was partially labeled with antibodies to lectin-GSII (a Golgi marker) and Bip/GRP78 (an endoplasmic reticulum marker). Pulse-chase labeling of MRP1 with 35S-methionine and 35S-cysteine and pulse-chase biotinylation of cell surface MRP1 suggests that membrane protein mislocalization is caused mainly by a defect of plasma membrane protein recycling, manifested also as a defect in acidification of lysosomes. The reduced accumulation of cytotoxic compds. in the KB-CP cells is presumed to result from the failure of carrier proteins and/or transporters to localize to the plasma membrane.

OS.CITING REF COUNT: 35 THERE ARE 35 CAPLUS RECORDS THAT CITE THIS RECORD (35 CITINGS)

REFERENCE COUNT: 30 THERE ARE 30 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 36 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:691408 CAPLUS

DOCUMENT NUMBER: 139:254928

TITLE: Topical vidarabine or 5-fluorouracil treatment against persistent HPV infection in genital (pre)cancerous lesions

AUTHOR(S): Niwa, Kenji; Tagami, Keiko; Lian, Zenglin; Gao, Jingchun; Mori, Hideki; Tanaya, Teruhiko

CORPORATE SOURCE: Departments of Obstetrics and Gynecology, Gifu University School of Medicine, Gifu-city, 500-8705, Japan

SOURCE: Oncology Reports (2003), 10(5), 1437-1441  
CODEN: OCRFEW; ISSN: 1021-335X

PUBLISHER: Oncology Reports

DOCUMENT TYPE: Journal

LANGUAGE: English

AB In the present study, effectiveness of topical vidarabine or subsequent 5-fluorouracil (5-FU) administration was examined against persistent genital human papillomavirus (HPV) infection after local surgery. Thirty patients underwent local eradication treatment of uterine cervical intra-epithelial neoplasia (CIN) and stage Ial uterine cervical cancers. HPV typing was performed by PCR-RFLP anal. HPV infection was detected pre-operatively in 29 of 30 patients. Of these, HPV was still present in the 20 patients within two months after the therapy. Topical administration of vidarabine or subsequent 5-FU once a week for four weeks was performed to the post-operative persistent

HPV-pos. cases. HPV infection was abolished in 1 of 10 (10%) with topical vidarabine, and in 2 of 4 vidarabine-resistant cases (50%) with topical 5-FU. Topical vidarabine or 5-FU treatment is beneficial for HPV-pos. cases after local surgical excision.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

REFERENCE COUNT: 22 THERE ARE 22 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 37 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:590597 CAPLUS

DOCUMENT NUMBER: 139:144951

TITLE: Preparation of fusion genes encoding streptavidin and single chain antibody and methods of therapeutic use thereof

INVENTOR(S): Goshorn, Stephen Charles; Graves, Scott Stoll; Schultz, Joanne Elaine; Lin, Yukang; Sanderson, James Allen; Reno, John M.; Dearstyne, Erica A.

PATENT ASSIGNEE(S): NeoRx Corporation, USA

SOURCE: U.S. Pat. Appl. Publ., 89 pp., Cont.-in-part of U.S. Ser. No. 150,762.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030143233	A1	20030731	US 2002-244821	20020916 <--
US 20030095977	A1	20030522	US 2001-13173	20011207 <--
US 7144991	B2	20061205		
US 20030103948	A1	20030605	US 2002-150762	20020517 <--
WO 2003050260	A2	20030619	WO 2002-US39429	20021206 <--
WO 2003050260	A3	20041125		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
AU 2002353095	A1	20030623	AU 2002-353095	20021206 <--
EP 1499630	A2	20050126	EP 2002-790070	20021206
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
PRIORITY APPLN. INFO.:				
			US 1999-137900P	P 19990607
			US 1999-168976P	P 19991203
			US 2000-589870	A2 20000605
			US 2001-13173	A2 20011207
			US 2002-150762	A2 20020517
			US 2002-244821	A 20020916
			WO 2002-US39429	W 20021206

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention provides vectors for expressing genomic streptavidin fusion cassettes and therapeutic uses. In the various embodiments, fusion proteins produced from these vectors are provided. In particular embodiments, fusion proteins comprising a single chain antibody and

genomic streptavidin are provided as are vectors encoding the same. Also provided, are methods of using the fusion proteins of the present invention, in the absence and presence of a radiation-sensitizing agent, and in particular, the use of scFvSA fusion proteins as diagnostic markers or as a cell specific targeting agents.

L21 ANSWER 38 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:472615 CAPLUS

DOCUMENT NUMBER: 139:30800

TITLE: Streptavidin expressed gene fusions with single-chain antibodies and their use as targeting vehicles for diagnosis and treatment of cancer

INVENTOR(S): Goshorn, Stephen Charles; Graves, Scott Stoll; Schultz, Joanne Elaine; Lin, Yukang; Sanderson, James Allen; Reno, John M.; Dearstyne, Erica A.

PATENT ASSIGNEE(S): Neorx Corporation, USA  
SOURCE: PCT Int. Appl., 156 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003050260	A2	20030619	WO 2002-US39429	20021206 <--
WO 2003050260	A3	20041125		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 20030095977	A1	20030522	US 2001-13173	20011207 <--
US 7144991	B2	20061205		
US 20030103948	A1	20030605	US 2002-150762	20020517 <--
US 20030143233	A1	20030731	US 2002-244821	20020916 <--
AU 2002353095	A1	20030623	AU 2002-353095	20021206 <--
EP 1499630	A2	20050126	EP 2002-790070	20021206
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK			

PRIORITY APPLN. INFO.:  
US 2001-13173 A 20011207  
US 2002-150762 A 20020517  
US 2002-244821 A 20020916  
US 1999-137900P P 19990607  
US 1999-168976P P 19991203  
US 2000-589870 A2 20000605  
WO 2002-US39429 W 20021206

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention provides vectors for expressing genomic streptavidin fusion cassettes. In the various embodiments, fusion proteins produced from these vectors are provided. In particular embodiments, fusion proteins comprising a single-chain antibody and genomic streptavidin are provided as are vectors encoding the same. The single-chain antibodies are directed to cell surface antigens, or cell-associated stromal or matrix antigens, including, but not limited to, CD20, CD22, CD25, CD45, CD52, CD56, CD57, EGP40 (or EPCAM or KSA), N-CAM, CEA, TAG-72,

$\gamma$ -glutamyl transferase, mucins (MUC1 through MUC7), human  $\beta$ -chorionic gonadotropin, EGF receptor, interleukin-2 receptor, her2/neu, Lewis Y, gangliosides GD2 and GM2, tenascin, sialylated tenascin, somatostatin, activated tumor stromal antigen, or neangiogenic antigens. Generically, a single-chain Fv/streptavidin (scFvSA) fusion protein is expressed from the genetic fusion of the single-chain antibody of the variable regions to the genomic streptavidin of *Streptomyces avidinii*. The scFv gene consists of the variable regions of the light and heavy chains separated by a DNA linker sequence. The streptavidin coding sequence is joined to the 3'-terminus of the scFv gene, and the two genes are separated in-frame by a second DNA linker sequence. The signal sequence from the streptavidin gene is fused at the 5'-terminus of the scFvSA gene to direct expression to the *Escherichia coli* periplasmic space. The scFvSA gene is under control of the lac promoter, and the expressed fusion protein is extracted and purified from *E. coli* and forms a soluble tetramer of .apprx.173,000 mol. weight. Also provided, are methods of using the fusion proteins of the present invention, in the absence and presence of a radiation-sensitizing agent (e.g., Gemcitabine), and in particular, the use of scFvSA fusion proteins as diagnostic markers or as cell-specific targeting agents.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L21 ANSWER 39 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2003:466122 CAPLUS

DOCUMENT NUMBER: 139:285604

TITLE: A Bayesian method for predicting 5-fluorouracil

pharmacokinetic parameters following short-term infusion in patients with colorectal cancer

AUTHOR(S): Climente-Marti, M.; Merino-Sanjuan, M.;

Almenar-Cubells, D.; Jimenez-Torres, N. V.

CORPORATE SOURCE: Pharmacy Service, Hospital Universitario Dr. Peset.

Avda. Gaspar Aguilar, Valencia, 46017, Spain

SOURCE: Journal of Pharmaceutical Sciences (2003),

92(6), 1155-1165

CODEN: JPMSAE; ISSN: 0022-3549

PUBLISHER: Wiley-Liss, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The objective of this study was to develop a population pharmacokinetic model and validate it using a Bayesian approach for predicting, a priori and a posteriori, the individual volume of distribution (Vd) and clearance (Cl) of 5-fluorouracil (5-FU) given as short-term i.v. infusion in weekly and multiple doses. Forty-four patients were divided in group A (5-FU weekly doses) including 27 patients with nonmetastatic colorectal adenocarcinoma treated with 450 mg/m<sup>2</sup> of 5-FU, 1 day per wk for 48 doses, plus oral levamisole (50 mg/8 h) for 3 days, every 15 days and group B (5-FU multiple doses) including 17 patients with metastatic colorectal adenocarcinoma, receiving 5-FU (425 mg/m<sup>2</sup>) plus i.v. folinic acid (20 mg/m<sup>2</sup>) over 5 consecutive days, every 4 wk for six cycles. In both groups 5-FU was administered as a 30-60-min infusion. A total of 176 plasma concns. were analyzed using a NONMEM program according to a linear one-compartment model. In group A, 5-FU population pharmacokinetic parameters were obtained and the covariables studied were age, gender, weight, ideal body weight, height, body surface area, creatinine clearance, and hepatic function tests. A priori and a posteriori validation of this model was carried out with plasma concns. obtained in day 1 in group B. In group B, population pharmacokinetic parameters of 5-FU following multiple doses were estimated using scale factors to identify differences in 5-FU Vd and Cl between days 1 and 4, and the interindividual, interoccasion, and residual variabilities studied. Vd

was 0.266 L/kg of ideal body weight and Cl was 1.21 L/h · kg of total weight following weekly doses. The plasma sample obtained at 10 min gave the best accuracy and precision predictions. When 5-FU was administered in multiple doses, the Cl of the drug in day 4 is reduced by 30.14% compared to day 1. The interoccasion variability was lower than interindividual variability for both Vd and Cl, suggesting that it could be feasible to individualize dosage of 5-FU for subsequent cycles from data obtained in a previous one in an attempt to improve the therapeutic index of colorectal cancer treatment.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)  
REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 40 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:454837 CAPLUS

DOCUMENT NUMBER: 139:41797

TITLE: Lipid vehicles for drug delivery

INVENTOR(S): Chancellor, Michael B.; Fraser, Matthew O.; Chuang, Yao-Chi; De Groat, William C.; Huang, Leaf; Yoshimura, Naoki

PATENT ASSIGNEE(S): University of Pittsburgh, USA

SOURCE: U.S. Pat. Appl. Publ., 31 pp., Cont.-in-part of U.S. Provisional Ser. No. 311,868.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030108597	A1	20030612	US 2002-218797	20020813 <--
US 7063860	B2	20060620		
US 20070003610	A1	20070104	US 2006-438912	20060522
US 20070122466	A1	20070531	US 2006-546025	20061011
PRIORITY APPLN. INFO.:			US 2001-311868P	P 20010813
			US 2002-218797	A3 20020813
			US 2005-701431P	P 20050720
			US 2005-725402P	P 20051011
			US 2006-438912	A2 20060522
			US 2006-489748	A2 20060719

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention relates to compns. and methods for the administration of lipid-based vehicles to treat various disorders, including bladder inflammation, infection, dysfunction, and cancer. In various aspects, the compns. and methods of the invention are useful for prolonged delivery of drugs, e.g., antibiotics, pain treatments, and anticancer agents, to the bladder, genitourinary tract, gastrointestinal system, pulmonary system, and other organs or body systems. In particular, the present invention relates to liposome-based delivery of vanilloid compds., such as resiniferatoxin, capsaicin, or tinyatoxin, and toxins, such as botulinum toxin, for the treatment of bladder conditions, including pain, inflammation, incontinence, and voiding dysfunction. Further related are methods of using these vehicles alone or in conjunction with antibodies, e.g., uroplakin antibodies, to improve duration of liposome attachment, and provide a long-term intravesical drug delivery platform. The present invention specifically relates to antibody-coated liposomes that are useful for targeting specific receptors for drug, peptide, polypeptide, or nucleic acid delivery. In one particular aspect, the present invention relates to liposomes coated with

antibodies against nerve growth factor (NGF) receptor and containing NGF antisense nucleic acids, which are used as a treatment for neurogenic bladder dysfunction. Liposomes are capable of highly effective delivery of at least one hydrophobic drug, CAP, as evidenced by a dramatic increase in bladder contraction frequency and subsequent desensitization. Moreover, liposomes alone had no effect on the micturition reflex in the unirritated state. In combination with other expts. that have demonstrated a protective effect of liposomes, this suggested that the liposome vehicle may partially protect against the compromise of urothelial barrier function due to the neuro-inflammatory response caused by irritants.

OS.CITING REF COUNT: 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)  
 REFERENCE COUNT: 25 THERE ARE 25 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 41 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:454119 CAPLUS

DOCUMENT NUMBER: 139:17567

TITLE: Aryl urea compounds in combination with other cytostatic or cytotoxic agents for treating human cancers and other raf kinase-mediated diseases  
 INVENTOR(S): Carter, Christopher A.; Dumas, Jacques; Gibson, Neil; Hibner, Barbara; Humphrey, Rachel W.; Trail, Pamela; Vincent, Patrick W.; Zhai, Yifan; Riedl, Bernd; Khire, Uday; Lowinger, Timothy B.; Scott, William J.; Smith, Roger A.; Wood, Jill E.; Monahan, Mary-Katherine; Natero, Reina; Renick, Joel; Sibley, Robert N.

PATENT ASSIGNEE(S): Bayer Corporation, USA; Bayer AG

SOURCE: PCT Int. Appl., 52 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003047579	A1	20030612	WO 2002-US38439	20021203 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG				
CA 2468463	A1	20030612	CA 2002-2468463	20021203 <--
AU 2002351196	A1	20030617	AU 2002-351196	20021203 <--
US 20030232765	A1	20031218	US 2002-308187	20021203 <--
EP 1450799	A1	20040901	EP 2002-786842	20021203 <--
EP 1450799	B1	20061115		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2005511658	T	20050428	JP 2003-548834	20021203
AT 345130	T	20061215	AT 2002-786842	20021203
EP 1769795	A2	20070404	EP 2006-23696	20021203
EP 1769795	A3	20080312		
R: AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT,				

LI, LU, MC, NL, PT, SE, SI, SK, TR, AL, LT, LV, MK, RO

ES 2275931	T3	20070616	ES 2002-786842	20021203
RU 2316326	C2	20080210	RU 2004-120785	20021203
IN 2004DN01420	A	20070316	IN 2004-DN1420	20040526
IN 233603	A1	20090403		
MX 2004005137	A	20050603	MX 2004-5137	20040528
ZA 200404225	A	20050829	ZA 2004-4225	20040528
US 20060247186	A1	20061102	US 2006-480360	20060705
IN 2008DN07086	A	20080926	IN 2008-DN7086	20080820

PRIORITY APPLN. INFO.:

US 2001-334609P	P	20011203
EP 2002-786842	A3	20021203
US 2002-308187	B1	20021203
WO 2002-US38439	W	20021203
IN 2004-DN1420	A3	20040526

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 139:17567

AB The invention discloses aryl urea compds. in combination with cytotoxic or cytostatic agents for use in treating raf kinase-mediated diseases, e.g. cancer.

OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)

REFERENCE COUNT: 10 THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 42 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:435061 CAPLUS

DOCUMENT NUMBER: 139:21033

TITLE: Vectors expressing soluble form of single chain antibody and streptavidin (scFvSA) fusions and uses thereof as diagnostic markers or as cell specific targeting agents

INVENTOR(S): Goshorn, Stephen Charles; Graves, Scott Stoll; Schultz, Joanne Elaine; Lin, Yukang; Sanderson, James Allen; Reno, John M.; Dearstyne, Erica A.

PATENT ASSIGNEE(S): NeoRx Corporation, USA

SOURCE: U.S. Pat. Appl. Publ., 84 pp., Cont.-in-part of U.S. Ser. No. 13,173.

CODEN: USXXCO

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 5

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 20030103948	A1	20030605	US 2002-150762	20020517 <--
US 20030095977	A1	20030522	US 2001-13173	20011207 <--
US 7144991	B2	20061205		
US 20030143233	A1	20030731	US 2002-244821	20020916 <--
WO 2003050260	A2	20030619	WO 2002-US39429	20021206 <--
WO 2003050260	A3	20041125		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SI, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			



AU 2002353095 A1 20030623 AU 2002-353095 20021206 <--  
 EP 1499630 A2 20050126 EP 2002-790070 20021206  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK  
 PRIORITY APPLN. INFO.:  
 US 1999-137900P P 19990607  
 US 1999-168976P P 19991203  
 US 2000-589870 A2 20000605  
 US 2001-13173 A2 20011207  
 US 2002-150762 A2 20020517  
 US 2002-244821 A 20020916  
 WO 2002-US39429 W 20021206

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention provides vectors for expressing Streptomyces avidinii genomic streptavidin (SA) fusion cassettes. A genomic streptavidin expressed gene fusion is expressed as a soluble protein into the periplasmic space of bacteria and undergoes spontaneous folding. Such expression offers the advantage that the periplasm is a low biotin environment and one need not purify and refold the protein under harsh denaturing conditions that may prove fatal to the polypeptide encoded by a heterologous nucleic acid mol. fused to the genomic streptavidin nucleic acid mol. In the various embodiments, fusion proteins produced from these vectors are provided. In particular embodiments, fusion proteins comprising a single chain antibody and streptavidin (scFvSA) are provided as are vectors encoding the same. The single chain antibodies are directed to cell surface antigens or cell-associated stromal or matrix proteins such as CD20, CD45, CD22, CD52, CD56, CD57, EGP40, NCAM, CEA, TAG-72, mucins (MUC1-7), 13HCG, EGF receptor, IL-2 receptor, her2/neu, Lewis Y, GD2, GM2, tenascin, sialylated tenascin, somatostatin, activated tumor stromal antigen or neoangiogenic antigens. Also provided, are methods of using the fusion proteins of the present invention, in the absence and presence of a radiation-sensitizing agent, and in particular, the use of scFvSA fusion proteins as diagnostic markers or as a cell specific targeting agents.

L21 ANSWER 43 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:417704 CAPLUS  
 DOCUMENT NUMBER: 139:981  
 TITLE: Clusianone isomers and use thereof for the treatment of tumors and viral diseases  
 INVENTOR(S): Seeber, Siegfried; Hilger, Ralf Axel; Diaz-Carballo, David  
 PATENT ASSIGNEE(S): Germany  
 SOURCE: PCT Int. Appl., 64 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: German  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003043966	A2	20030530	WO 2002-EP12968	20021120 <--
WO 2003043966	A3	20031023		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES,			

FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF,  
 CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG  
 DE 10157031 A1 20030605 DE 2001-10157031 20011121 <--  
 AU 2002356679 A1 20030610 AU 2002-356679 20021120 <--  
 EP 1448504 A2 20040825 EP 2002-803387 20021120 <--  
 R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,  
 IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK  
 JP 2005509670 T 20050414 JP 2003-545607 20021120  
 US 20050090562 A1 20050428 US 2004-496592 20041029  
 US 7135501 B2 20061114

PRIORITY APPLN. INFO.: DE 2001-10157031 A 20011121  
 WO 2002-EP12968 W 20021120

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention discloses a clusianone isomer and to the use thereof, in particular as a pharmaceutical or medically active ingredient, in particular for producing medicaments for the prophylaxis and/or treatment of tumors and viral diseases. The compound of the invention can be used in cytostatics and antiviral agents. The compound acts, in particular, as a topoisomerase and telomerase inhibitor and as a regulator in the MAP kinase signal transduction pathway. The compound can thus intervene at the cellular level in the proliferation mechanism of tumor or cancer cells and viruses.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
 (1 CITINGS)

L21 ANSWER 44 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:406590 CAPLUS

DOCUMENT NUMBER: 139:977

TITLE: Substituted bicyclo[3.3.1]nonane-2,4,9-triones as pharmaceutically active substances for the treatment of cancer and viral diseases

INVENTOR(S): Seeber, Siegfried; Hilger, Ralf Axel; Diaz-Carballo, David

PATENT ASSIGNEE(S): Germany

SOURCE: Ger. Offen., 46 pp.

CODEN: GWXXBX

DOCUMENT TYPE: Patent

LANGUAGE: German

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
DE 10157033	A1	20030528	DE 2001-10157033	20011121 <--
CA 2506616	A1	20030530	CA 2002-2506616	20021120 <--
WO 2003043622	A1	20030530	WO 2002-EP12967	20021120 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GO, GW, ML, MR, NE, SN, TD, TG				
AU 2002366225	A1	20030610	AU 2002-366225	20021120 <--
AU 2002366225	B2	20080131		
EP 1448179	A1	20040825	EP 2002-790404	20021120 <--
EP 1448179	B1	20060809		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				

IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK

CN 1615127	A	20050511	CN 2002-827411	20021120
CN 100415219	C	20080903		
JP 2005515979	T	20050602	JP 2003-545303	20021120
AT 335474	T	20060915	AT 2002-790404	20021120
ES 2269791	T3	20070401	ES 2002-790404	20021120
IN 2004MN00337	A	20050218	IN 2004-MN337	20040615
IN 239087	A1	20100312		
ZA 2004004887	A	20050908	ZA 2004-4887	20040621
US 20050090693	A1	20050428	US 2004-496593	20041029
US 7199161	B2	20070403		

PRIORITY APPLN. INFO.: DE 2001-10157033 A 20011121  
WO 2002-EP12967 W 20021120

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT  
OTHER SOURCE(S): MARPAT 139:977

AB The invention describes the use of substituted Bicyclo[3.3.1]nonane-2,4,9-triones, in particular clusianone and clusianone derivs., as pharmaceutically active substances, in particular for the production of drugs for the prevention and/or treatment of cancers and viral diseases. They act in particular as inhibitors of topoisomerases and telomerases, as well as regulators within the MAP kinase signal transduction pathway and can in this way intervene at the cellular level in the multiplication mechanism of tumor cells and of viruses.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L21 ANSWER 45 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2003:376654 CAPLUS

DOCUMENT NUMBER: 138:390922

TITLE: Arsenide compound system for selective targeting of apoptotic cells

INVENTOR(S): Hogg, Philip John

PATENT ASSIGNEE(S): Unisearch Limited, Australia

SOURCE: PCT Int. Appl., 85 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003039564	A1	20030515	WO 2002-AU1523	20021108 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SC, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2466303	A1	20030515	CA 2002-2466303	20021108 <--
AU 2002340631	A1	20030519	AU 2002-340631	20021108 <--
AU 2002340631	B2	20060810		
EP 1453525	A1	20040908	EP 2002-774165	20021108 <--
EP 1453525	B1	20090930		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK				
JP 2005511598	T	20050428	JP 2003-541855	20021108

AT 444083	T	20091015	AT 2002-774165	20021108
ES 2330203	T3	20091207	ES 2002-774165	20021108
ZA 2004003803	A	20060329	ZA 2004-3803	20040518
US 20050101524	A1	20050512	US 2004-494822	20041124
US 7635464	B2	20091222		
US 20090311179	A1	20091217	US 2009-433401	20090430

PRIORITY APPLN. INFO.:

AU 2001-8746	A	20011108
WO 2002-AU1523	W	20021108
US 2004-494822	A3	20041124

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 138:390922

AB The invention discloses a method of selectively targeting an active agent (or agent capable of becoming an active agent) to apoptotic cells in a vertebrate, comprising administering to the vertebrate a system comprising an arsenoxide (or arsenoxide equivalent) compound and the agent, wherein the system selectively targets apoptotic cells. Preparation of e.g. 4-[N-(S-glutathionylacetyl)amino]phenylarsenoxide is described.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 46 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2003:362680 CAPLUS

DOCUMENT NUMBER: 139:97397

TITLE: Randomized comparison of photodynamic therapy with topical 5-fluorouracil in Bowen's disease

AUTHOR(S): Salim, A.; Leman, J. A.; McColl, J. H.; Chapman, R.; Morton, C. A.

CORPORATE SOURCE: Department of Dermatology, Falkirk Royal Infirmary, Falkirk, FK1 5QE, UK

SOURCE: British Journal of Dermatology (2003), 148(3), 539-543

CODEN: BJDEAZ; ISSN: 0007-0963

PUBLISHER: Blackwell Publishing Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Bowen's disease (BD; intraepithelial squamous cell carcinoma) is therapeutically challenging because lesions, which may be multiple, are frequently located at sites that heal poorly. There is a small risk of progression to invasive carcinoma. Photodynamic therapy (PDT) is an effective treatment for certain non melanoma skin cancers, but comparison studies with other, better-established therapies are limited. The aim was to compare the efficacy and tolerability of PDT and topical 5-fluorouracil (5-FU) in BD. Forty patients from two centers were randomized to either topical PDT or 5-FU. The PDT group was treated with 20% 5-aminolaevulinic acid (ALA) applied 4 h before illumination with 100 J cm<sup>-2</sup> narrowband red light (630 ± 15 nm). 5-FU was applied to lesions for 4 wk. A repeat treatment cycle was performed after 6 wk if required. Twenty-nine of 33 (88%) lesions treated with PDT initially responded completely, compared with 22 of 33 (67%) after 5-FU. After 12 mo, two recurrences in the PDT group and six in the 5-FU group reduced complete clin. clearance rates to 82% and 48%, resp. PDT was significantly more effective (P = 0.006, odds ratio 4.78, 95% confidence interval 1.56-14.62). In the 5-FU group, severe eczematous reactions developed around seven lesions, ulceration in three and erosions in two. No such reactions occurred following PDT. There was no difference in overall pain experienced during each therapy. Topical ALA-PDT is more effective than topical 5-FU in the treatment of BD, with fewer adverse events. ALA-PDT should be considered one of the first-line therapeutic options for BD.

OS.CITING REF COUNT: 31 THERE ARE 31 CAPLUS RECORDS THAT CITE THIS  
RECORD (31 CITINGS)  
REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 47 OF 124 CAPLUS COPYRIGHT 2010 ACS ON STN  
ACCESSION NUMBER: 2003:202527 CAPLUS  
DOCUMENT NUMBER: 138:210373  
TITLE: Potentiator of antitumoral agents in the treatment of  
cancer  
INVENTOR(S): Pichette, Andre; Legault, Jean  
PATENT ASSIGNEE(S): F.P.L. Pharma Inc., Can.  
SOURCE: PCT Int. Appl., 17 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2003020371	A2	20030313	WO 2002-CA1359	20020905 <--
WO 2003020371	A3	20030717		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2356438	A1	20030305	CA 2001-2356438	20010905 <--
CA 2458805	A1	20030313	CA 2002-2458805	20020905 <--
AU 2002325721	A1	20030318	AU 2002-325721	20020905 <--
EP 1423169	A2	20040602	EP 2002-759977	20020905 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR, BG, CZ, EE, SK			
JP 2005501128	T	20050113	JP 2003-524674	20020905
US 20040235785	A1	20041125	US 2004-488682	20040406 <--
US 20090286865	A1	20091119	US 2009-510196	20090727
PRIORITY APPLN. INFO.:			CA 2001-2356438	A 20010905
			WO 2002-CA1359	W 20020905
			US 2004-488682	A1 20040406

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention relates to a potentiator composition for enhancing  
therapeutic effect of an antitumoral agent, said composition comprising a  
terpene or derivative thereof in association with a pharmaceutically acceptable  
carrier. Synergistic efficacy of a mixture of paclitaxel and  
β-caryophyllene was shown on human breast adenocarcinoma.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD  
(2 CITINGS)  
REFERENCE COUNT: 9 THERE ARE 9 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 48 OF 124 CAPLUS COPYRIGHT 2010 ACS ON STN  
ACCESSION NUMBER: 2003:9768 CAPLUS  
DOCUMENT NUMBER: 138:50256  
TITLE: Phase II clinical trial of local use of GM-CSF for  
prevention and treatment of chemotherapy- and

concomitant chemoradiotherapy-induced severe oral mucositis in advanced head and neck cancer patients: an evaluation of effectiveness, safety and costs  
 AUTHOR(S): Mantovani, Giovanni; Massa, Elena; Astara, Giorgio; Murgia, Viviana; Gramignano, Giulia; Lusso, Maria Rita; Camboni, Paolo; Ferrelli, Luca; Mocci, Miria; Perboni, Simona; Mura, Loredana; Madeddu, Clelia; Maccio, Antonio  
 CORPORATE SOURCE: Department of Medical Oncology, University of Cagliari, Policlinico Universitario, Cagliari, Italy  
 SOURCE: Oncology Reports (2003), 10(1), 197-206  
 CODEN: OCRFEW; ISSN: 1021-335X  
 PUBLISHER: Oncology Reports  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB In the present open non-randomized phase II study we looked for effectiveness, safety, tolerability and costs of locally applied GM-CSF in preventing or treating mucositis in patients receiving chemotherapy or chemoradiotherapy for head and neck cancer. In addition to clin. mucositis scoring system, the effects of treatment with GM-CSF were evaluated by its impact on patient quality of life and by laboratory immunol. assays such as serum pro-inflammatory cytokines, IL-2 and leptin. The trial was designed to assess the effectiveness of local GM-CSF treatment in two different settings: (i) prophylaxis of mucositis; (ii) treatment of mucositis. Prophylaxis was chosen for chemoradiotherapy treatments of high mucosatoxic potential, while curative treatment was reserved for chemotherapy or chemoradiotherapy treatments of lesser potential of inducing mucositis. From Jan. 1998 to Dec. 2001, 68 patients entered the study. The great majority of patients of both groups had head and neck cancer, were stage IV, PS ECOG 0-1, were habitual smokers and were treated with chemotherapy and concomitant (or sequential) chemoradiotherapy. Forty-six patients were included in the 'prophylactic' setting and 22 patients in the 'curative' setting. The main findings of our study are: only 50% of patients included in the 'prophylactic' setting developed mucositis; the duration of oral mucositis from appearance until complete remission was significantly shorter in the 'prophylactic' than in the 'curative' setting; the mean grade of oral mucositis at baseline, on day 3 of therapy and on day 6 of therapy was significantly lower in the 'prophylactic' than in the 'curative' setting; 24 (55.82%) patients in the 'prophylactic' setting had grade 3/4 oral mucositis at baseline compared to 25 (80.60%) patients in the 'curative' setting ( $p=0.048$ ). Thirteen (30.23%) patients in the 'prophylactic' setting had grade 3/4 oral mucositis on day 3 of therapy compared to 19 (61.29%) patients in the 'curative' setting ( $p=0.015$ ); 'prophylactic' setting was able to shorten grade 3/4 oral mucositis to grade 0/1 more effectively than the 'curative' one on day 6 of therapy ( $p=0.05$ ). The present clin. trial is to date by far the largest study assessing the effectiveness of topical GM-CSF and it is the first study comparing the efficacy of topical GM-CSF in the 'prophylactic' setting, i.e., with the aim to prevent the chemoradiotherapy-induced oral mucositis, with that in the 'curative' treatment, i.e., the therapy for established oral mucositis. The topical application of GM-CSF was demonstrated to be effective for oral mucositis induced by chemotherapy and chemoradiotherapy regimens. Moreover, the 'prophylactic' setting was demonstrated to be more effective than the 'curative' one.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)  
 REFERENCE COUNT: 14 THERE ARE 14 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

ACCESSION NUMBER: 2002:906491 CAPLUS  
 DOCUMENT NUMBER: 137:379912  
 TITLE: Effect of topical morphine for mucositis-associated pain following concomitant chemoradiotherapy for head and neck carcinoma  
 AUTHOR(S): Cerchietti, Leandro C. A.; Navigante, Alfredo H.; Bonomi, Marcelo R.; Zaderajko, Mariel A.; Menendez, Pablo R.; Pogany, Catalina E.; Roth, Berta M. C.  
 CORPORATE SOURCE: Supportive Care Division, Department of Medical Oncology, Angel H. Roffo Cancer Institute, University of Buenos Aires, Buenos Aires, Argent.  
 SOURCE: Cancer (New York, NY, United States) (2002), 95(10), 2230-2236  
 CODEN: CANCAR; ISSN: 0008-543X  
 PUBLISHER: John Wiley & Sons, Inc.  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Oral mucositis is the dose-limiting toxicity for patients receiving concurrent chemoradiotherapy regimens for tumors of the head and neck area. Currently, the management of established mucositis includes the use of topical anesthetics and systemic analgesics. Based on the clin. evidence of pain alleviation by topical morphine in patients with some inflammatory and painful conditions, a clin. study was undertaken to determine this effect on mucositis-associated pain. Twenty-six patients with head and neck malignancies treated with concomitant chemoradiotherapy for head and neck carcinoma who had severe painful mucositis (World Health Organization Grade 2 or higher) were enrolled. Patients were randomly assigned to morphine mouthwash (MO; 14 patients) or magic mouthwash (MG), a mixture of equal parts of lidocaine, diphenhydramine, and magnesium aluminum hydroxide (12 patients). The duration of severe pain was 3.5 days less in the MO group compared with the MG group (P = 0.032). The intensity of oral pain was also significantly lower in the MO group compared with the MG group (P = 0.038). No patient in the MO group required third-step opiates for alleviation of the mouth pain. There was a significant difference in duration of severe functional impairment (P = 0.017). Five patients in the MG group complained of local side effects and only one in the MO group (P = 0.007). For patients with head and neck carcinomas receiving concomitant chemoradiotherapy, MO is a simple and effective treatment to decrease the severity and duration of pain and the duration of functional impairment.  
 OS.CITING REF COUNT: 15 THERE ARE 15 CAPLUS RECORDS THAT CITE THIS RECORD (15 CITINGS)  
 REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 50 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2002:850321 CAPLUS  
 DOCUMENT NUMBER: 137:342158  
 TITLE: Fluorouracil-containing formulation  
 INVENTOR(S): Singh, B. Sandhya; Saxena, Subhash J.  
 PATENT ASSIGNEE(S): A. P. Pharma, Inc., USA  
 SOURCE: U.S. Pat. Appl. Publ., 6 pp.  
 CODEN: USXXCO  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

US 20020165198 A1 20021107 US 2001-799792 20010305 <--  
US 6670335 B2 20031230

PRIORITY APPLN. INFO.: US 2001-799792 20010305

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Oil-in-water emulsion formulations contain both free fluorouracil and fluorouracil impregnated in porous microparticles are described. The formulations are suitable for topical administration, and are useful for the treatment of solar (actinic) keratosis and superficial basal cell carcinomas. For example, fluorouracil-impregnated microparticles were prepared containing fluorouracil 15.0%, Dimethicone 200 40.0%, and microparticles 45.0%. Fluorouracil-impregnated microparticles were then used to prepare oil-in-water emulsion containing 1.5% fluorouracil

and  
3.5% fluorouracil-impregnated microparticles.

L21 ANSWER 51 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:846443 CAPLUS

DOCUMENT NUMBER: 139:143268

TITLE: Human Pharmacokinetic Study of Heated Intraperitoneal Oxaliplatin in Increasingly Hypotonic Solutions after Complete Resection of Peritoneal Carcinomatosis  
AUTHOR(S): Elias, D.; El Otmány, A.; Bonnay, M.; Paci, A.; Ducreux, M.; Antoun, S.; Lasser, P.; Laurent, S.; Bourget, P.

CORPORATE SOURCE: Departments of Surgical Oncology, Clinical Biology, Pharmacy and Medical Oncology, Institut Gustave Roussy, Villejuif, F-94805, Fr.  
SOURCE: Oncology (2002), 63(4), 346-352  
CODEN: ONCOBS; ISSN: 0030-2414

PUBLISHER: S. Karger AG

DOCUMENT TYPE: Journal

LANGUAGE: English

AB We studied the pharmacokinetics of heated intraoperative i.p. oxaliplatin (LOHP) solution and its safety profile in increasingly hypotonic solns. This is the first clin. study of i.p. chemohyperthermia with hypotonic solns. Patients with peritoneal carcinomatosis (PC) underwent complete cytoreductive surgery followed by intraoperative i.p. chemohyperthermia (IPCH) with successive dextrose solns. of 300, 200, 150 and 100 mosm/L. LOHP (460 mg/m2) was administered in 2 L of solution/m2 at an i.p. temperature

of  
42-44°C for 30 min. IPCH was performed using an open procedure (skin pulled upward) with a continuous closed circuit. Patients received i.v. leucovorin (20 mg/m2) and 5-fluorouracil (400 mg/m2) just before IPCH to maximize the effect of LOHP. i.p. plasma and tissue samples were analyzed by means of atomic absorption spectrophotometry. Sixteen consecutive patients with PC of either gastrointestinal or peritoneal origin were treated. The safety of the procedure was studied. The mean duration of the entire procedure was 7.7 ± 2.6 h. Half the LOHP dose was absorbed within 30 min at all dose levels. Absorption was not higher with hypotonic solns. than with isotonic solns. The area under the curve of LOHP in plasma did not increase with decreasing osmolality of the i.p. solns. Intratumoral LOHP penetration was high; it was similar to that at the peritoneal surface, and about 18 times higher than that in non-bathed tissues. LOHP penetration was not significantly increased by using hypotonic solns. There was a very high incidence of unexplained post-operative peritoneal bleeding (50%) and unusually severe thrombocytopenia in the 150 and 100 mosm/L groups. Thus, contrary to exptl. studies, this clin. study showed no increase in tumoral or systemic penetration of LOHP with i.p. hypotonic solns. (200, 150 or 100 mosm/L) during IPCH. A high incidence of i.p. hemorrhage and thrombocytopenia was observed



OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD  
(8 CITINGS)  
REFERENCE COUNT: 24 THERE ARE 24 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 52 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2002:675858 CAPLUS  
DOCUMENT NUMBER: 137:222036  
TITLE: Compositions based on vanilloid-catechin synergies for  
prevention and treatment of cancer  
INVENTOR(S): Morre, Dorothy M.; Morre, James D.  
PATENT ASSIGNEE(S): Purdue Research Foundation, USA  
SOURCE: PCT Int. Appl., 51 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 2  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002067966	A1	20020906	WO 2002-US5295	20020222 <--
<p>W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZM, ZW</p> <p>RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG</p>				
AU 2002254007	A1	20020912	AU 2002-254007	20020222 <--
AU 2002254007	B2	20060921		
<p>PRIORITY APPLN. INFO.: US 2001-270557P P 20010222 WO 2002-US5295 W 20020222</p>				
<p>AB The invention described herein encompasses methods and compns. of preventing or treating cancer comprising the administration of a combination of catechins and vanilloids. Compns. of catechins include but not limited to, epigallocatechin gallate (EGCg), epicatechin (EC), epicatechin gallate (ECG), epigallocatechin (EGC). In a preferred embodiment the catechins have been treated with tannase. Compns. of vanilloids include, but are not limited to vanillylamine, the head group of capsaicin. The unique compns. of the invention contain various combinations of the catechins and vanilloids, in combination with each other or other therapeutic agents and are used to treat primary and metastatic cancers in humans. The invention also encompasses various modes of administration of the therapeutic compds., including formulations which may be used as a dietary or nutritional supplement or as a therapeutic compound. The effect of combinations of tea catechins (including tannase-treated Tegreen with and without gallic acid and EGCg) and the vanilloid vanillylamine, alone and in combination, was demonstrated on (i) cancer cell growth and (ii) NADH oxidase (tNOX) activity. The ratios of tea catechins and vanillylamine was varied to determine optimum ratios for the inhibition of cancer cell growth and the inhibition of tNOX activity. A synergy between tannase-treated Tegreen with gallic acid and vanillylamine in inhibiting the cell surface NADH oxidase was observed</p>				
REFERENCE COUNT:	3	THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT		

L21 ANSWER 53 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2002:521793 CAPLUS

DOCUMENT NUMBER: 137:77889  
 TITLE: Human antibodies to insulin-like growth factor I receptor  
 INVENTOR(S): Cohen, Bruce D.; Beebe, Jean; Miller, Penelope E.; Moyer, James D.; Corvalan, Jose R.; Gallo, Michael  
 PATENT ASSIGNEE(S): Pfizer Inc., USA; Abgenix, Inc.  
 SOURCE: PCT Int. Appl., 172 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002053596	A2	20020711	WO 2001-US51113	20011220 <--
WO 2002053596	A3	20040108		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2433800	A1	20020711	CA 2001-2433800	20011220 <--
AU 2002231368	A1	20020716	AU 2002-231368	20011220 <--
AU 2002231368	B2	20061123		
EE 200300318	A	20031015	EE 2003-318	20011220 <--
HU 2003002525	A2	20031028	HU 2003-2525	20011220 <--
EP 1399483	A2	20040324	EP 2001-991634	20011220 <--
EP 1399483	B1	20100414		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
ZA 2003005995	A	20040804	ZA 2003-5995	20011220 <--
JP 2004531217	T	20041014	JP 2002-555118	20011220 <--
CN 1564829	A	20050112	CN 2001-821808	20011220
CN 1330668	C	20070808		
IN 2001CA00696	A	20050311	IN 2001-CA696	20011220
BR 2001016728	A	20050412	BR 2001-16728	20011220
NZ 527302	A	20061027	NZ 2001-527302	20011220
CN 1854157	A	20061101	CN 2006-10059704	20011220
SG 138469	A1	20080128	SG 2005-6384	20011220
IL 156661	A	20081126	IL 2001-156661	20011220
AP 2072	A	20091231	AP 2001-2365	20011220
NZ 569856	A	20100326	NZ 2001-569856	20011220
US 20040086503	A1	20040506	US 2002-38591	20020104 <--
US 7037498	B2	20060502		
MX 2003006034	A	20050701	MX 2003-6034	20030703
NO 2003003074	A	20030704	NO 2003-3074	20030704 <--
KR 830082	B1	20080520	KR 2003-709063	20030705
BG 108037	A	20050430	BG 2003-108037	20030728
HR 2003000627	A2	20050630	HR 2003-627	20030804
IN 2003KN00994	A	20050708	IN 2003-KN994	20030804
US 20050244408	A1	20051103	US 2005-144248	20050602
US 20050281812	A1	20051222	US 2005-144222	20050602
US 7700742	B2	20100420		
HK 1072059	A1	20080118	HK 2005-104844	20050608
AU 2007200793	A1	20070315	AU 2007-200793	20070222

IN 2007KN02634	A	20080801	IN 2007-KN2634	20070713
JP 2009108055	A	20090521	JP 2008-267173	20081016
JP 2009297037	A	20091224	JP 2009-222178	20090928
PRIORITY APPLN. INFO.:			US 2001-259927P	P 20010105
			AU 2002-231368	A3 20011220
			CN 2001-821808	A3 20011220
			JP 2002-555118	A3 20011220
			WO 2001-US51113	W 20011220
			US 2002-38591	A3 20020104
			IN 2003-KN994	A3 20030804
			JP 2008-267173	A3 20081016

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The authors disclose the preparation and characterization of antibodies that specifically bind to human insulin-like growth factor I receptor (IGF-IR). The antibodies were prepared by immunization of XenoMouse with either the extracellular domain of human IGF-IR or with cells transformed for surface expression of the receptor. The isolated antibodies were shown to down-regulate IGF-IR, to prevent its phosphorylation induced by ligand, and to exhibit tumor growth inhibitory activities either alone or in combination with chemotherapeutic agents.

OS.CITING REF COUNT: 22 THERE ARE 22 CAPLUS RECORDS THAT CITE THIS RECORD (23 CITINGS)

REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 54 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:521462 CAPLUS

DOCUMENT NUMBER: 137:88442

TITLE: Incensole and furanogermacrens and compounds in treatment for inhibiting neoplastic lesions and microorganisms

INVENTOR(S): Shanahan-Pendergast, Elisabeth

PATENT ASSIGNEE(S): Ire.

SOURCE: PCT Int. Appl., 68 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002053138	A2	20020711	WO 2002-IE1	20020102 <--
WO 2002053138	A3	20020919		
W: AE, AG, AT, AU, BB, BG, CA, CH, CN, CO, CU, CZ, LU, LV, MA, MD, UA, UG, US, VN, YU, RU, TJ, TM				
RW: GH, GM, KE, LS, MW, SD, SL, SZ, UG, AT, BE, CH, CY, DE, ES, FI, ML, MR, NE, SN, TD, TG				
AU 2002219472	A1	20020716	AU 2002-219472	20020102 <--
EP 1351678	A2	20031015	EP 2002-727007	20020102 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 20040092583	A1	20040513	US 2004-250535	20040102 <--
PRIORITY APPLN. INFO.:			IE 2001-2	A 20010102
			WO 2002-IE1	W 20020102

OTHER SOURCE(S): MARPAT 137:88442

AB The invention discloses the use of incensole and/or furanogermacrens, derivs. metabolites and precursors thereof in the treatment of neoplasia, particularly resistant neoplasia and immunodysregulatory disorders. These compds. can be administered alone or in combination with conventional chemotherapeutic, antiviral, antiparasite agents, radiation and/or

surgery. Incensole and furanogermacren and their mixture showed antitumor activity against various human carcinomas and melanomas and antimicrobial activity against *Staphylococcus aureus* and *Enterococcus faecalis*.

OS.CITING REF COUNT: 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 55 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:428749 CAPLUS

DOCUMENT NUMBER: 137:28318

TITLE: Conjugates of glycosylated/galactosylated peptide, bifunctional linker, and nucleotidic monomers/polymers, and related compositions and methods of use

INVENTOR(S): Ts'o, Paul O. P.; Duff, Robert; Deamond, Scott

PATENT ASSIGNEE(S): Cell Works Inc., USA; Johns Hopkins University

SOURCE: PCT Int. Appl., 90 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002043771	A2	20020606	WO 2001-US44943	20011130 <--
WO 2002043771	A3	20030828		
WO 2002043771	A9	20040429		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2431839	A1	20020606	CA 2001-2431839	20011130 <--
AU 2002017980	A	20020611	AU 2002-17980	20011130 <--
US 20030032584	A1	20030213	US 2001-998497	20011130 <--
US 6906182	B2	20050614		
EP 1355672	A2	20031029	EP 2001-998366	20011130 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2004536027	T	20041202	JP 2002-545741	20011130 <--
US 20050250679	A1	20051110	US 2005-152135	20050614
US 7262177	B2	20070828		

PRIORITY APPLN. INFO.:

US 2000-250139P P 20001201  
 US 2001-998497 A1 20011130  
 WO 2001-US44943 W 20011130

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The invention discloses a conjugate A-L-P (A = glycosylated/galactosylated peptide that binds to cell-surface receptor; L = bifunctional linker, which does not comprise naturally occurring amino acid and is covalently bonded to A and P in regiospecific manner; P = monomer, homopolymer, or heteropolymer comprising at least one nucleotide, or analog thereof, which inhibits intracellular biosynthesis of nucleotides or nucleic acids in sequence-independent manner, wherein either or both of covalent bond between A and L and the covalent bond between L and P can be

cleaved intracellularly); a composition comprising such a conjugate; a method of inhibiting an abnormal cellular proliferation in a mammal; and a method of inhibiting replication of a virus in a mammal.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)

L21 ANSWER 56 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:368329 CAPLUS

DOCUMENT NUMBER: 136:374857

TITLE: Combination pharmaceuticals containing a biological response modifier and an anticancer agent

INVENTOR(S): Young, Aiping H.

PATENT ASSIGNEE(S): Lorus Therapeutics Inc., Can.

SOURCE: PCT Int. Appl., 62 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002038164	A1	20020516	WO 2001-CA1558	20011108 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MY, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG				
CA 2428145	A1	20020516	CA 2001-2428145	20011108 <--
AU 2002014876	A	20020521	AU 2002-14876	20011108 <--
EP 1333847	A1	20030813	EP 2001-983364	20011108 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
US 20040101511	A1	20040527	US 2004-416259	20040102 <--
US 20050192443	A1	20050901	US 2004-821649	20040408
US 20060034938	A1	20060216	US 2005-247026	20051011
US 20080279818	A1	20081113	US 2008-61274	20080402
PRIORITY APPLN. INFO.:			CA 2000-2325361	A 20001108
			WO 2001-CA1558	W 20011108
			US 2004-416259	A2 20040102
			US 2005-247026	B1 20051011

AB The present invention provides anticancer bile-derived biol. response modifier (BD-BRM) combinations. In accordance with an aspect of the present invention, there is provided a combination comprising: a composition comprising small mol. weight (<3000 daltons) components, and 1 or more anticancer agents. The modifier is extracted from animal bile, is capable of stimulating monocytes and/or macrophages in vitro and/or in vivo, is capable of modulating tumor necrosis factor production and/or release, contains no measurable levels of IL-1 $\alpha$ , IL-1 $\beta$ , TNF, IL-6, IL-8, IL-4, GM-CSF or IFN- $\gamma$ , is not cytotoxic to human peripheral blood mononuclear cells, and is not an endotoxin. The combination has therapeutic synergy or improves the therapeutic index in the treatment of cancer over the anticancer agents used alone. Another aspect of the present invention provides the use of this combination in the manufacture of a medicament or a pharmaceutical kit and in the treatment of cancer. The mouse xenograft model of neoplasia was used in these studies to demonstrate the effect of treatment with a BD-BRM composition on tumor growth

in mice. BD-BRM treatments always resulted in significant delay of tumor growth compared to saline control. Where a chemotherapeutic drug treatment group was included, the delay in tumor growth achieved with BD-BRM was typically superior to the inhibitory effects observed with the chemotherapeutic drug. Total regression of the tumor was also observed in some of the animals, when the animals were treated with a BRM composition alone or with a combination of the BD-BRM composition and a chemotherapeutic drug was used. In the remaining animals treated with a combination, significantly enhanced antitumor effects were observed

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)  
 REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 57 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2002:185083 CAPLUS  
 DOCUMENT NUMBER: 136:226783  
 TITLE: Chelating agent and method of prevention and treatment of cancer and other diseases in animals  
 INVENTOR(S): Fernandez-Pol, Jose A.  
 PATENT ASSIGNEE(S): Novactyl, Inc., USA  
 SOURCE: PCT Int. Appl., 73 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 9  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002020486	A2	20020314	WO 2001-US27578	20010905 <--
WO 2002020486	A3	20020704		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
US 6579891	B1	20030617	US 2000-657554	20000908 <--
AU 2001088788	A	20020322	AU 2001-88788	20010905 <--
PRIORITY APPLN. INFO.:			US 2000-657554	A 20000908
			US 1995-581351	A2 19951229
			US 1996-26992P	P 19960920
			US 1996-24221P	P 19961022
			US 1997-843157	B2 19970411
			US 1998-127620	A2 19980801
			US 2000-182608P	P 20000215
			WO 2001-US27578	W 20010905

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 136:226783

AB An antiproliferative, anti inflammatory, antiinfective, immunization agent of a metal ion chelating agent such as picolinic acid or derivs. thereof, and methods of using the same. The agents chelate metals in metal containing protein complexes and enzymes required for growth, replication or inflammatory response. The prepn. can be administered systemically or for topical use. The prepn. have antineoplastic, antiviral, antiinflammatory, analgesic antiangiogenic and antiproliferative effects and are used in the treatment of warts, psoriasis, acne, skin cancers,

sunburn, inflammatory responses, untoward angiogenesis and other diseases and in the prevention of sexually transmitted diseases such as genital warts, herpes and AIDS.

REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 58 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:171627 CAPLUS

DOCUMENT NUMBER: 136:226776

TITLE: Methods of treatment of a bcl-2 disorder using bcl-2 antisense oligomers

INVENTOR(S): Warrel, Raymond P., Jr.; Klem, Robert E.; Fingert, Howard

PATENT ASSIGNEE(S): Genta Incorporated, USA

SOURCE: PCT Int. Appl., 64 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002017852	A2	20020307	WO 2001-US26414	20010823 <--
WO 2002017852	A3	20030403		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
CA 2419480	A1	20020307	CA 2001-2419480	20010823 <--
AU 2001088373	A	20020313	AU 2001-88373	20010823 <--
EP 1313514	A2	20030528	EP 2001-968097	20010823 <--
EP 1313514	B1	20090603		
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
BR 2001013447	A	20030708	BR 2001-13447	20010823 <--
HU 2003003125	A2	20031229	HU 2003-3125	20010823 <--
JP 2004507480	T	20040311	JP 2002-522827	20010823 <--
EE 200300074	A	20041215	EE 2003-74	20010823 <--
AU 2001288373	B2	20060511	AU 2001-288373	20010823
AT 432717	T	20090615	AT 2001-968097	20010823
PT 1313514	E	20090901	PT 2001-968097	20010823
ES 2327904	T3	20091105	ES 2001-968097	20010823
EP 2135623	A1	20091223	EP 2009-7273	20010823
R:	AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC, NL, PT, SE, TR, AL, LT, LV, MK, RO, SI			
ZA 2003001161	A	20040225	ZA 2003-1161	20030212 <--
HR 2003000102	A2	20050430	HR 2003-102	20030213
MX 2003001575	A	20041101	MX 2003-1575	20030221 <--
NO 2003000858	A	20030424	NO 2003-858	20030224 <--
IN 2003CN00395	A	20050408	IN 2003-CN395	20030313
BG 107641	A	20040130	BG 2003-107641	20030318 <--
HK 1056505	A1	20090807	HK 2003-108693	20031128
PRIORITY APPLN. INFO.:			US 2000-227970P	P 20000825
			US 2000-237009P	P 20000929

US 2000-709170 A 20001110  
 EP 2001-968097 A3 20010823  
 WO 2001-US26414 W 20010823

AB The present invention is directed to the use of bcl-2 antisense oligomers to treat and prevent bcl-2 related disorders. These disorders include cancers, tumors, carcinomas and cell-proliferative related disorders. In one embodiment of the invention, a bcl-2 antisense oligomer is administered at high doses. The present invention is also directed to a method of preventing or treating a bcl-2 related disorder, in particular cancer, comprising administering a bcl-2 antisense oligomer for short periods of time. The present invention is further drawn to the use of bcl-2 antisense oligomers to increase the sensitivity of a subject to cancer therapeutics. The present invention also relates to pharmaceutical compns. comprising one or more bcl-2 antisense oligomers, which may comprise one or more cancer therapeutic agents.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)  
 REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 59 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:157495 CAPLUS

DOCUMENT NUMBER: 136:205412

TITLE: Oligopeptide-based prodrugs activated by plasmin and their use in cancer chemotherapy

INVENTOR(S): Trouet, Andre; Dubois, Vincent; Passioukov, Alexandre

PATENT ASSIGNEE(S): Coulter Pharmaceutical, Inc., USA

SOURCE: PCT Int. Appl., 67 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002015700	A1	20020228	WO 2001-US26476	20010823 <--
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG			
AU 2001086727	A	20020304	AU 2001-86727	20010823 <--
US 20040171562	A1	20040902	US 2003-362958	20031031 <--
US 7402556	B2	20080722		
US 20090076176	A1	20090319	US 2008-157575	20080611
PRIORITY APPLN. INFO.:			US 2000-227686P	P 20000824
			WO 2001-US26476	W 20010823
			US 2003-362958	A1 20031031

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

OTHER SOURCE(S): MARPAT 136:205412

AB A prodrug, cleavable by plasmin, comprises a therapeutic agent capable of entering a target cell, e.g., a tumor or inflammatory cell, an oligopeptide having a plasmin peptide substrate of 2-4 amino acids and mono- or di-peptide linkage, a stabilizing group and, optionally, a linker group. Also disclosed are methods of making and using the prodrug compds. For example, the activity of D-Ala-Leu-Lys-Leu-Leu-doxorubicin (I) (preparation



given) was evaluated in the B16-B16 murine melanoma model. The mice receiving the prodrug did not show any important weight loss during the experiment and no clin. signs of toxicity were observed. At the same time, the drug had a marked effect on the metastatic growth. At 34.5  $\mu\text{mol/kg}$ , I reduced the spread of lung metastases with a decrease of the ratio of the surface occupied by B16-B16 colonies to the non-affected one to  $8.2 \pm 1.8\%$  ( $P < 0.01$ ), compared to  $45.7 \pm 12.6\%$  and  $44.0 \pm 6.3\%$  for non-treated and doxorubicin ( $5.2 \mu\text{mol/kg}$ )-treated animals. The same prodrug at  $69.0 \mu\text{mol/kg}$  provided  $1.5 \pm 0.6\%$  of surface affected.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)  
 REFERENCE COUNT: 4 THERE ARE 4 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 60 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:71908 CAPLUS

DOCUMENT NUMBER: 136:112640

TITLE: Hyaluronan as a cytotoxic agent, drug pre-sensitizer

and chemo-sensitizer in the treatment of disease

INVENTOR(S): Brown, Tracey; Fox, Richard

PATENT ASSIGNEE(S): Mediatech Research Limited, Australia

SOURCE: PCT Int. Appl., 70 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002005852	A1	20020124	WO 2001-AU849	20010713 <--
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2382560	A1	20020124	CA 2001-2382560	20010713 <--
GB 2368525	A	20020508	GB 2002-4331	20010713 <--
GB 2368525	B	20040811		
EP 1301209	A1	20030416	EP 2001-951219	20010713 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
AU 760404	B2	20030515	AU 2001-72202	20010713 <--
NZ 517359	A	20031128	NZ 2001-517359	20010713 <--
JP 2004502789	T	20040129	JP 2002-511783	20010713 <--
ZA 2002001561	A	20031125	ZA 2002-1561	20020225 <--
US 20030180382	A1	20030925	US 2003-88774	20030313 <--
US 20050267069	A1	20051201	US 2005-191407	20050727
US 20060178342	A1	20060810	US 2005-198663	20050805
US 20060263395	A1	20061123	US 2006-415612	20060501
PRIORITY APPLN. INFO.:			AU 2000-8795	A 20000714
			WO 2001-AU849	W 20010713
			US 2003-88774	A2 20030313

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention relates to the enhancement of bioavailability of

chemotherapeutic agents for the treatment of disease. In particular the present invention relates to a method of enhancing the bioavailability of a chemotherapeutic agent comprising the step of administering to a subject in need thereof a therapeutically effective amount of hyaluronan. The present invention also relates to the treatment of a drug resistant disease whereby the drug resistance is overcome or alleviated with the use of hyaluronan either alone or in combination with a chemotherapeutic agent. One disease that is frequently affected by both cellular resistance and bioavailability problems is cancer. The present invention also provides a method of treating cancer cells comprising the step of administering to a patient in need thereof a therapeutically effective amount of hyaluronan.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD  
(1 CITINGS)  
REFERENCE COUNT: 6 THERE ARE 6 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 61 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2002:37903 CAPLUS

DOCUMENT NUMBER: 137:103494

TITLE: Treatment of human colon carcinoma cell lines with anti-neoplastic agents enhances their lytic sensitivity to antigen-specific CD8+ cytotoxic T lymphocytes

AUTHOR(S): Bergmann-Leitner, Elke S.; Abrams, Scott I.  
CORPORATE SOURCE: Laboratory of Tumor Immunology and Biology, Center for Cancer Research, National Cancer Institute, National Institutes of Health, Bethesda, MD, 20892-1402, USA  
SOURCE: Cancer Immunology Immunotherapy (2001), 50(9), 445-455

CODEN: CIIMDN; ISSN: 0340-7004

PUBLISHER: Springer-Verlag

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Certain anti-neoplastic agents at subtoxic doses may exert immunomodulatory effects, which alter the expression of specific tumor cell surface molecules. We reasoned that potential increases in tumor cell surface markers, such as those important for facilitating effector-target contact, as well as triggering cell death pathways, might then improve antigen (Ag)-specific T-cell-mediated tumor cytotoxicity. Here, in a human colon carcinoma cell model in vitro, we examined whether the anti-neoplastic agents 5-fluorouracil (5-FU), CPT-11 or cisplatin (CDDP) could upregulate the expression of specific tumor cell surface markers, which may then enhance productive lytic interactions between CD8+ CTL and Ag-bearing tumor cells. Based on our earlier studies, IFN- $\gamma$  treatment was included as a control for sensitization to CTL-mediated lysis. Pretreatment of the SW480 primary colon carcinoma cell line with IFN- $\gamma$ , 5-FU, CPT-11 or CDDP enhanced ICAM-1 and Fas expression, resulting in Ag-specific CTL-mediated lysis involving Fas-dependent and -independent mechanisms. In contrast, pretreatment of the SW620 metastatic isolate, derived from the same patient, with IFN- $\gamma$ , CPT-11 or CDDP, but not 5-FU, enhanced ICAM-1 expression, resulting in Ag-specific CTL-mediated lysis via Fas-independent mechanisms only. Flow cytometric-based assays were then developed to measure the effects of drug treatment on caspase signaling and apoptosis incurred by tumor targets after interaction with CTL. We found that the lytic enhancement caused by drug treatment of SW480 or SW620 targets was accompanied by an increase in caspase-3-like protease activity. A peptide-based caspase inhibitor abrogated CTL-mediated apoptosis, suggesting that "chemomodulation" involved regulation of the caspase pathway. These results revealed for the first time an important

role for components of the caspase pathway, such as caspase-3-like proteases, in the sensitization of human colon carcinoma cells by anti-neoplastic agents to Ag-specific CTL. Thus, certain anti-neoplastic agents may display unique immunoregulatory properties that facilitate human colon carcinoma death by engaging the lytic capacity of Ag-specific CTL, which may have implications for chemoinmunotherapy strategies.

OS.CITING REF COUNT: 33 THERE ARE 33 CAPLUS RECORDS THAT CITE THIS RECORD (33 CITINGS)  
 REFERENCE COUNT: 33 THERE ARE 33 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 62 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2002:31286 CAPLUS  
 DOCUMENT NUMBER: 136:90918  
 TITLE: Isolation of a cell-specific internalizing peptide that infiltrates tumor tissue for targeted drug delivery  
 INVENTOR(S): Clayman, Gary; Hong, Frank D.  
 PATENT ASSIGNEE(S): Board of Regents, the University of Texas System, USA  
 SOURCE: PCT Int. Appl., 104 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2002002147	A2	20020110	WO 2001-US21518	20010702 <--
WO 2002002147	A3	20020725		
W: CA, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR				
CA 2414650	A1	20020110	CA 2001-2414650	20010702 <--
US 20020102265	A1	20020801	US 2001-899376	20010702 <--
US 6919425	B2	20050719		
EP 1297002	A2	20030402	EP 2001-958886	20010702 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY, TR				
JP 2004501664	T	20040122	JP 2002-506768	20010702 <--
US 20060188437	A1	20060824	US 2005-53602	20050208
PRIORITY APPLN. INFO.:				
			US 2000-215491P	P 20000630
			US 2001-899376	A3 20010702
			WO 2001-US21518	W 20010702

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention provides a tumor-homing peptide that can target cancer and/or tumor tissues. The peptide is taken up by certain specific cancer cell types. The invention describes methods to achieve targeted delivery of anticancer drugs conjugated to this peptide for anticancer therapy. The invention also describes methods for using the peptide for the diagnosis and imaging of cancer and tumor tissues.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L21 ANSWER 63 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2002:662 CAPLUS  
 DOCUMENT NUMBER: 136:212879  
 TITLE: Alternating chemoradiotherapy versus partly accelerated radiotherapy in locally advanced squamous cell carcinoma of the head and neck: Results

from a phase III randomized trial

AUTHOR(S): Corvo, Renzo; Benasso, Marco; Sanguineti, Giuseppe; Lionetto, Rita; Bacigalupo, Almalina; Margarino, Giovanni; Pallestrini, Eugenio; Merlano, Marco; Vitale, Vito; Rosso, Riccardo

CORPORATE SOURCE: Oncologia Radioterapica, National Cancer Research Institute, Genoa, Italy

SOURCE: Cancer (New York, NY, United States) (2001), 92(11), 2856-2867  
CODEN: CANCAR; ISSN: 0008-543X

PUBLISHER: John Wiley & Sons, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The authors previously have found that in patients with locally advanced squamous cell carcinoma of the head and neck (SCC-HN), alternating chemoradiotherapy (ALT) was superior to low-total-dose conventional radiotherapy alone. The purpose of this randomized trial was to compare the same chemoradiotherapy approach with high-total-dose partly accelerated radiotherapy. During 6 yr, 136 consecutive patients with previously untreated unfavorable Stage II or Stage III-IV (International Union Against Cancer) SCC of the oral cavity, pharynx, and larynx were enrolled. They were randomly assigned to chemotherapy consisting of 4 cycles of i.v. cisplatin (20 mg/m<sup>2</sup> of body surface area per day for 5 consecutive days) and 5-fluorouracil (200 mg/m<sup>2</sup> per day for 5 consecutive days; weeks 1, 4, 7, and 10) alternated with three 2-wk courses of radiotherapy (20 grays [Gy] per course, 2 Gy per day, 5 days per wk; ALT, 70 patients) or to partly accelerated radiotherapy with final concomitant boost technique (75 Gy/40 fractions in 6 wk; partly accelerated radiotherapy [PA-RT], 66 patients). At the median follow-up of 60 mo (range, 30-102 mo), no statistical differences were observed in overall survival, progression free survival, or locoregional control between the 2 treatments. Actuarial 3-yr overall survival and progression free survival were 37% and 35%, resp., in the ALT group and 29% and 27%, resp., in PA-RT group. The median overall survival and progression free survival were 24 and 15 mo, resp., in the ALT arm and 18 and 11 mo, resp., in PA-RT arm. Actuarial 3-yr locoregional control rates were 32% in the ALT group and 27% in the PA-RT group. At multivariate anal., tumor classification was the only factor that emerged as a significant independent variable affecting overall survival. Patients treated in the PA-RT arm experienced higher Grade 3+ (World Health Organization) acute skin and mucosal reactions than patients in the ALT arm. Moreover, local late mucosal and skin toxicities occurred more often in patients treated with PA-RT. This trial failed to disclose statistically significant differences in the outcome of patients treated with either ALT or PA-RT. Therefore, definitive conclusions could not be made. However, acute skin effects and late mucosal and skin toxicities above the clavicles appeared to be significantly lower with chemoradiotherapy.

OS.CITING REF COUNT: 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

REFERENCE COUNT: 35 THERE ARE 35 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 64 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2001:894190 CAPLUS

DOCUMENT NUMBER: 136:334871

TITLE: Up-regulation of Fas expression by 5-fluorouracil in human colon cancer cells

AUTHOR(S): Hu, Shengliang; Ding, Erxun; Wang, Qiang; Chen, Xueyun; Fu, Zhiren

CORPORATE SOURCE: Department of General Surgery, Changzheng Hospital, Second Military Medical University, Shanghai, 200003,

SOURCE:                   Peop. Rep. China  
                           Dier Junyi Daxue Xuebao (2001), 22(9),  
                           809-811  
                           CODEN: DJXUE5; ISSN: 0258-879X  
 PUBLISHER:               Dier Junyi Daxue Xuebao Bianjibu  
 DOCUMENT TYPE:         Journal  
 LANGUAGE:               Chinese  
 AB   The role of 5-fluorouracil (5-Fu) in modulation of Fas expression in human colon carcinoma cells was studied. 6 Human colon cancer cell lines were examined for Fas cell surface protein expression and for 5-Fu modulation of Fas expression by flow cytometry method. Fas was expressed in 5 of the 6 cancer cell lines. Fas expression was up-regulated by 5-Fu at least in part of colon cancer cells. The results showed that 5-Fu may enhance Fas expression in tumor cells and may increase the sensibility of them to Fas-mediated immune surveillance of the host.

L21 ANSWER 65 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER:       2001:854288 CAPLUS  
 DOCUMENT NUMBER:       136:128733  
 TITLE:                   Efficacy of intraperitoneal and intravenous chemotherapy and left upper abdominal evisceration for advanced gastric cancer  
 AUTHOR(S):              Nomura, Eiji; Niki, Masami; Fujii, Keizou; Shinohara, Hisashi; Nishiguchi, Kanji; Sonoda, Toyooki; Tanigawa, Nobuhiko  
 CORPORATE SOURCE:       Department of General and Gastroenterological Surgery, Osaka Medical College, Takatsuki, 569-8686, Japan  
 SOURCE:                 Gastric Cancer (2001), 4(2), 75-82  
                           CODEN: GCANFO; ISSN: 1436-3291  
 PUBLISHER:               Springer-Verlag Tokyo  
 DOCUMENT TYPE:         Journal  
 LANGUAGE:               English  
 AB   The study was carried out to evaluate the efficacy of i.p. (IP) and i.v. (IV) chemotherapy, as well as left upper abdominal evisceration (LUAE), for patients with advanced gastric cancer. We carried out a retrospective study of 348 patients who underwent gastrectomy for advanced gastric carcinoma between 1978 and 1998 at our institution and who had macroscopic type 3 or 4 cancer (Japanese classification) with depth of invasion to the serosal surface, but no liver metastasis or lymph node metastasis around the abdominal aorta. Cumulative survival rates were compared in patients who underwent gastrectomy together with: (1) intraoperative IP chemotherapy alone, (2) postoperative IV chemotherapy alone, (3) both IP and IV, or (4) no chemotherapy. Then patients were stratified according to the presence of peritoneal dissemination (P+) and its absence (P-). In P+ patients, survival was compared between those who received IV chemotherapy and those who did not, and between those who received IP chemotherapy and those who did not. Then, survival was compared between patients with high and low immunosuppressive acidic protein (IAP) levels. Finally, we compared cumulative survival rates in patients (stratified as P+ and P-) who underwent LUAE with cumulative survival rates in those who underwent total gastrectomy combined with resection of the pancreatic body, tail, and spleen (PS). For P- patients, there was no survival advantage with adjuvant IP or IV therapy when compared with surgery alone. For P+ patients, however, there was an improvement in survival when patients received both IP and IV, compared with survival with surgery alone (P < 0.05). In P+ patients aged less than 60 yr, there was improvement in survival for those who underwent IP therapy together with surgery (P < 0.05), but not for those who had IV chemotherapy after surgery. When LUAE was examined, there was a survival advantage for this procedure when there

was no peritoneal dissemination. Four long-term survivors (surviving for more than 5 yr) were identified in our study. Three of the 4 patients were aged less than 60 yr, and all 4 had macroscopic type 4 gastric cancers. Although the prognosis for patients with invasive type gastric cancer remains poor, there have been a few long-term survivors, in whom this survival was associated with aggressive combination therapy, including surgery, IP, and IV therapy. P+ patients aged less than 60 yr and patients with type 4 gastric cancer may stand to benefit most from such therapy. For P- patients, the role of adjuvant IP or IV therapy continues to be ambiguous, although LUAE in this population may be superior to P5.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD  
(2 CITINGS)  
REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 66 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2001:713176 CAPLUS

DOCUMENT NUMBER: 135:262259

TITLE: Pharmaceutical comprising an agent that blocks the cell cycle and an antibody

INVENTOR(S): Stimmel, Julie Beth; Thurmond, Linda Margarite; Knick, Vincent Clark

PATENT ASSIGNEE(S): Glaxo Group Limited, UK

SOURCE: PCT Int. Appl., 74 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001070268	A1	20010927	WO 2001-US9368	20010322 <--
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
EP 1265635	A1	20021218	EP 2001-922610	20010322 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
JP 2003527441	T	20030916	JP 2001-568464	20010322 <--
US 20030138430	A1	20030724	US 2002-239278	20020920 <--
PRIORITY APPLN. INFO.:			US 2000-191336P	P 20000322
			WO 2001-US9368	W 20010322

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Pharmaceutical combinations comprise an agent that arrests target cells in the G2 and/or M phase of the cell cycle and another therapeutic agent that targets an internalizing cell surface structure such as an antigen. Manufacture of a medicament and methods of medical treatment, particularly in the treatment of diseases of cell cycle regulation such as cancer are disclosed.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD  
(3 CITINGS)

REFERENCE COUNT: 7 THERE ARE 7 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 67 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2001:617776 CAPLUS  
 DOCUMENT NUMBER: 135:175428  
 TITLE: Protection of the female reproductive system from natural and artificial insults  
 INVENTOR(S): Tilly, Jonathan L.; Kolesnick, Richard N.  
 PATENT ASSIGNEE(S): Sloan-Kettering Institute for Cancer Research, USA  
 SOURCE: PCT Int. Appl., 42 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 2  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001060318	A2	20010823	WO 2001-US4712	20010215 <--
WO 2001060318	A3	20020418		
W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
US 7195775	B1	20070327	US 2000-503852	20000215
CA 2400405	A1	20010823	CA 2001-2400405	20010215 <--
AU 2001038246	A	20010827	AU 2001-38246	20010215 <--
EP 1257245	A2	20021120	EP 2001-910661	20010215 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR			
US 20070157331	A1	20070705	US 2007-715795	20070307
PRIORITY APPLN. INFO.:			US 2000-503852	A 20000215
			WO 2001-US4712	W 20010215

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Described are methods for protecting the female reproductive system against natural and artificial insults by administering to women a composition comprising an agent that antagonizes one or more acid sphingomyelinase (ASMase) gene products. Specifically, methods disclosed herein serve to protect women's germline from damage resulting from cancer therapy regimens including chemotherapy or radiotherapy. In one aspect, the method preserves, enhances, or revives ovarian function in women, by administering to women a composition containing sphingosine-1-phosphate, or an analog thereof prior to therapy. Also disclosed are methods to prevent or ameliorate menopausal syndromes and to improve in vitro fertilization techniques.

REFERENCE COUNT: 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 68 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2001:380442 CAPLUS  
 DOCUMENT NUMBER: 135:9968  
 TITLE: Compositions and methods for treating disease utilizing a combination of radioactive therapy and cell-cycle inhibitors  
 INVENTOR(S): Hunter, William L.; Liggins, Richard  
 PATENT ASSIGNEE(S): Angiotech Pharmaceuticals, Inc., Can.  
 SOURCE: PCT Int. Appl., 270 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent

LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2001036007	A2	20010525	WO 2000-CA1333	20001114 <--
WO 2001036007	A3	20020704		
W: AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW				
RW: GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2388844	A1	20010525	CA 2000-2388844	20001114 <--
AU 2001013746	A	20010530	AU 2001-13746	20001114 <--
EP 1235598	A2	20020904	EP 2000-975703	20001114 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL, TR				
JP 2003513756	T	20030415	JP 2001-537997	20001114 <--
AU 2005203583	A1	20050908	AU 2005-203583	20050811
PRIORITY APPLN. INFO.:				
			US 1999-165259P	P 19991112
			US 2000-712047	A 20001113
			AU 2001-13746	A 20001114
			WO 2000-CA1333	W 20001114

AB Disclosed herein are therapeutic devices, compns. and methods for treating proliferative diseases. Within one aspect of the invention therapeutic devices are provided, comprising a device that locally administers radiation, and a cell-cycle inhibitor. Among examples provided are: paclitaxel-containing topical gels, thermally responsive pastes prepared from polycaprolactone and a cell cycle inhibitor such as vincristine, cast films containing paclitaxel in polyethylene vinyl acetate, cylindrically-shaped spacers from poly( $\epsilon$ -caprolactone) loaded with drug, and various devices such as wires and brachytherapy seeds coated with a cell cycle inhibitor.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)  
 REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 69 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 2001:279564 CAPLUS  
 DOCUMENT NUMBER: 134:285609  
 TITLE: Topical formulations containing hyaluronic acid  
 INVENTOR(S): Falk, Rudolf Edger; Asculai, Samuel Simon; Hochman, David; Purschke, Don; Klein, Ehud Shmuel; Harper, David William  
 PATENT ASSIGNEE(S): Hyal Pharmaceutical Corp., Can.  
 SOURCE: U.S., 49 pp., Cont.-in-part of U.S. Ser. No. 675,908.  
 CODEN: USXXAM  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 24  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----



US 6218373	B1	20010417	US 1995-467994	19950606 <--
WO 9104058	A2	19910404	WO 1990-CA306	19900918 <--
WO 9104058	A3	19910919		
W:	AT, AU, BB, BG, BR, CA, CH, DE, DK, ES, FI, GB, HU, JP, KP, KR, LK, LU, MC, MG, MW, NL, NO, RO, SD, SE, SU, US			
RW:	AT, BE, BF, BJ, CF, CG, CH, CM, DE, DK, ES, FR, GA, GB, IT, LU, ML, MR, NL, SE, SN, TD, TG			
CZ 288292	B6	20010516	CZ 1990-4598	19900921 <--
US 6069135	A	20000530	US 1991-675908	19910703 <--
CA 2061703	A1	19930821	CA 1992-2061703	19920220 <--
CA 2061703	C	20020702		
US 5827834	A	19981027	US 1994-286263	19940805 <--
US 6114314	A	20000905	US 1994-352697	19941201 <--
IN 182267	A1	19990227	IN 1995-CA270	19950313 <--
IN 182348	A1	19990327	IN 1995-CA271	19950313 <--
US 5811410	A	19980922	US 1995-465335	19950605 <--
US 5830882	A	19981103	US 1995-462615	19950605 <--
US 5852002	A	19981222	US 1995-462147	19950605 <--
US 5942498	A	19990824	US 1995-467171	19950606 <--
US 5977088	A	19991102	US 1995-467995	19950606 <--
US 6136793	A	20001024	US 1995-466715	19950606 <--
US 6147059	A	20001114	US 1995-468330	19950606 <--
US 6087344	A	20000711	US 1995-474732	19950607 <--
US 6194392	B1	20010227	US 1995-460978	19950807 <--
CA 2268476	A1	19980430	CA 1996-2268476	19961018 <--
WO 9817320	A1	19980430	WO 1996-CA700	19961018 <--
W:	AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN			
RW:	KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG			
AU 9672721	A	19980515	AU 1996-72721	19961018 <--
AU 739701	B2	20011018		
EP 952855	A1	19991103	EP 1996-934250	19961018 <--
EP 952855	B1	20050727		
R:	DE, FR, GB, IT, SE			
NZ 335259	A	20001222	NZ 1996-335259	19961018 <--
ZA 9608847	A	19970527	ZA 1996-8847	19961022 <--
IN 1996CA01848	A	20050304	IN 1996-CA1848	19961023 <--
US 6475795	B1	20021105	US 1997-860696	19970616 <--
AU 9742732	A	19980115	AU 1997-42732	19971020 <--
US 20030036525	A1	20030220	US 2002-234355	20020904 <--
PRIORITY APPLN. INFO.:			WO 1990-CA306	A2 19900918
			US 1991-675908	A2 19910703
			CA 1992-2061703	A 19920220
			US 1992-838674	B2 19920221
			US 1994-290840	A3 19941027
			CA 1989-612307	A 19890921
			CS 1990-4598	A 19900921
			CA 1992-2061566	A 19920220
			IN 1993-CA94	A1 19930216
			WO 1996-CA700	A 19961018
			US 1997-860696	A1 19970616

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Disclosed is a method of treating a disease of condition administered topically to the skin or exposed tissue of a human, a dosage amount of a pharmaceutical composition, said dosage comprising a therapeutically effective amount of a drug to treat said disease or condition and a form of hyaluronic acid characterized in that the composition is immediately available to

transport the drug percutaneously into the epidermis of the skin or exposed tissue to the site of trauma or pathol. of the disease or condition to be treated. A formulation containing glycerin 150, benzyl alc. 90, diclofenac sodium 90, sodium hyaluronate 75g, and water 2795 mL was prepared. The formulation was applied on the forehead of a patient having horny epithelium and some degree of ulceration.

OS.CITING REF COUNT: 24 THERE ARE 24 CAPLUS RECORDS THAT CITE THIS RECORD (46 CITINGS)  
REFERENCE COUNT: 382 THERE ARE 382 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 70 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2001:119520 CAPLUS

DOCUMENT NUMBER: 135:240848

TITLE: Influence of cytokines, monoclonal antibodies and chemotherapeutic drugs on epithelial cell adhesion molecule (EpCAM) and LewisY antigen expression  
AUTHOR(S): Flieger, D.; Hoff, A. S.; Sauerbruch, T.; Schmidt-Wolf, I. G. H.

CORPORATE SOURCE: Medizinische Klinik und Poliklinik I, Allgemeine Innere Medizin, Universitat Bonn, Bonn, D-53105, Germany

SOURCE: Clinical and Experimental Immunology (2001), 123(1), 9-14  
CODEN: CEXIAL; ISSN: 0009-9104

PUBLISHER: Blackwell Science Ltd.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB MoAbs against tumor-associated antigens (TAA) may be useful for the treatment of colorectal cancer. Since an increased expression of TAA may lead to enhanced antibody-dependent cellular cytotoxicity the authors examined whether the cytokines IL-2, IL-4, IL-6, IL-10, IL-12, interferon- $\alpha$  (IFN- $\alpha$ ), IFN- $\gamma$ , granulocyte-macrophage colony-stimulating factor, macrophage colony-stimulating factor, and tumor necrosis factor- $\alpha$  can influence EpCAM and LewisY expression on the surface of the colorectal carcinoma cell lines HT29, LoVo, and SW480. The authors found that only IFN- $\alpha$  increased whereas IL-4 decreased both EpCAM and LewisY expression. IFN- $\gamma$  increased LewisY expression only. When tumor cells were treated with MoAb, the LewisY-specific MoAb BR55-2 down-regulated LewisY antigen expression, whereas MoAb 17-1A, which binds to EpCAM, up-regulated this TAA after 3 days of culture. The cytokines IFN- $\alpha$  or IFN- $\gamma$  combined with MoAb 17-1A enhanced further slightly the expression of EpCAM. In addnl. expts. with chemotherapeutic drugs commonly used for the treatment of colorectal cancer, the authors found that 5-fluorouracil, mitomycin-C, and oxaliplatin upregulated EpCAM and LewisY antigen expression. Raltitrexed enhanced LewisY and down-regulated EpCAM expression, whereas CPT-11 had no influence at all. The highest expression for EpCAM on HT29 cells was achieved by the combination of IFN- $\alpha$ , 5-fluorouracil, and MoAb 17-1A. These results may thus be useful for defining combinations of biol. and chemotherapeutic drugs for the treatment of colorectal cancer.

OS.CITING REF COUNT: 20 THERE ARE 20 CAPLUS RECORDS THAT CITE THIS RECORD (20 CITINGS)

REFERENCE COUNT: 54 THERE ARE 54 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 71 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2001:70491 CAPLUS

DOCUMENT NUMBER: 135:86663

TITLE: In vitro antitumor activity of 9-nitrocarnitrothecin as a single agent and in combination with other antitumor drugs

AUTHOR(S): Bernacki, Ralph J.; Pera, Paula; Gambacorta, Peter; Brun, Yseult; Greco, William R.

CORPORATE SOURCE: Department of Pharmacology and Therapeutics, Roswell Park Cancer Institute, Buffalo, NY, 14263, USA

SOURCE: Annals of the New York Academy of Sciences ( 2000), 922(Carnitrothecins), 293-297  
CODEN: ANYAA9; ISSN: 0077-8923

PUBLISHER: New York Academy of Sciences

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Preclin. studies at Roswell Park Cancer Institute by Minderman, Cao, and Rustum (unpublished results) showed that a combination of SN-38 and 5-FU against HCT-8 human colon carcinoma cells in vitro was synergistic, with the best interaction occurring when the drugs were added sequentially, SN-38 first. Their in vivo studies using HCT-8 tumor xenografts implanted s.c. in nude athymic mice demonstrated superior efficacy for a sequential i.v. administration of CPT-11, 24 h before 5-FU. On the basis of these studies, our group has begun to evaluate effects of RFS2000 (9-nitro-20(S)-carnitrothecin) (9-NC) in combination with a series of other antitumor agents. Using a panel of human tumor cell lines including A121 ovarian cancer, HCT-8 colon cancer, H-460 NSCLC, HT-1080 fibrosarcoma, and MCF7 mammary cancer, we found that a 2-h exposure to 9-NC resulted in ID50 values of <1.0  $\mu$ M, whereas continuous exposure to drug resulted in ID50 values of <1.0 nM. Tumor growth inhibitory activities of 5-FU, gemcitabine, and paclitaxel were determined for comparison. Combinations of these agents were evaluated with 9-NC using the human HCT-8 colon tumor cell line. Concurrent and sequential combinations of 9-NC with 5-FU had some regions of the concentration-effect surface with local synergy and some with local antagonism. However, sequential combination of 9NC or SN-38 followed by 5-FU, 24 h later appeared to be highly synergistic at high dose-effect levels (i.e., ID90), suggesting that sequential drug administration may be more efficacious at high effect level and that the order of drug addition is very important. Overall, our results were similar to that found earlier by Rustum's group with CPT11 (or SN-38) and 5-FU, suggesting that sequential combination of 9-NC (or other carnitrothecin analogs) followed by 5-FU has potential for the treatment of cancer in man.

OS.CITING REF COUNT: 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 72 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2001:53269 CAPLUS

DOCUMENT NUMBER: 134:95249

TITLE: Adenovirus-mediated tumor-specific gene therapy using Cre/loxP system

AUTHOR(S): Ueda, Kentaro

CORPORATE SOURCE: Second Dep. Surg., Wakayama Med. Coll., Wakayama, Japan

SOURCE: Wakayama Igaku (2000), 51(4), 405-415  
CODEN: WKMIAO; ISSN: 0043-0013

PUBLISHER: Wakayama Igakkai

DOCUMENT TYPE: Journal

LANGUAGE: Japanese

AB Cre/loxP system has been employed to enhance the carcinoembryonic antigen (CEA) promoter activity and improve the suicide gene therapy using cytosine deaminase (CD)/5-fluorocytosine (5-FC). Four adenovirus vectors

were constructed; AxCEANCre expressing Cre recombinase under the control of CEA promoter, AxCALNLCD expressing CD gene under the control of the CAG (cytomegalovirus enhancer plus chicken  $\beta$ -actin) promoter by the Cre-mediated switching system, AxCEACD expressing CD gene driven by CEA promoter, AxCACD expressing CD gene driven by CAG promoter. In the orthotopic model of gastric carcinoma (each group: n=5), in which a tumor piece of MKN 45 (CEA producing gastric carcinoma cell) was fixed on the serosal surface of the glandular stomach of athymic BALB/c-nu/nu mice, Ad vectors (1 + 10<sup>9</sup> pfu/day + 3 days) were injected into the abdominal cavity 4 days after tumor implantation. Then, 5-FC (500 mg/kg) was administered i.p. once daily for the next ten days. Animals were sacrificed at the end of 4 wk, and tumor volume was measured. Tumor vols. in mice treated with AxCEANCre plus AxCALNLCD/5-FC, or AxCACD/5-FC were significantly reduced as compared to those in mice treated with AxCEACD/5-FC, Mock/PBS, or Mock/5-FC (p<0.0001). However, two of five mice treated with AxCACD/5-FC had died until the end of 4 wk. The median survival periods of mice treated with AxCEANCre plus AxCALNLCD/5-FC were significantly longer compared to those of mice treated with Mock/PBS, Mock/5-FC, or AxCEACD/5-FC (p<0.01). These results suggested that CEA specific suicide gene therapy enhanced by Cre/loxP system could be a very useful strategy for the treatment of patients with advanced gastric carcinoma.

L21 ANSWER 73 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 2000:880923 CAPLUS  
DOCUMENT NUMBER: 134:37055  
TITLE: Methods and compositions using FGF inhibitors and agonists for modulating cell proliferation and cell death  
INVENTOR(S): Au, Jessie L. S.; Wientjes, M. Guillaume  
PATENT ASSIGNEE(S): USA  
SOURCE: PCT Int. Appl., 143 pp.  
CODEN: PIXXD2  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000074634	A2	20001214	WO 2000-US40103	20000605 <--
WO 2000074634	A3	20010823		
WO 2000074634	A9	20020926		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, SZ, BE, CY, FR, GR, IE, IT, MC, NL, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2377385	A1	20001214	CA 2000-2377385	20000605 <--
EP 1206234	A2	20020522	EP 2000-943429	20000605 <--
R:	AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL			
JP 2003503313	T	20030128	JP 2001-501171	20000605 <--
US 6599912	B1	20030729	US 2000-587559	20000605 <--
AU 780454	B2	20050324	AU 2000-57903	20000605
IL 146872	A	20061031	IL 2000-146872	20000605

KR 903243	B1	20090617	KR 2001-715591	20011203
US 20040010001	A1	20040115	US 2003-464018	20030618 <--
US 7625860	B2	20091201		

PRIORITY APPLN. INFO.:

US 1999-137345P	P	19990603
US 1999-165983P	P	19991117
US 1999-172031P	P	19991223
US 2000-187445P	P	20000307
US 2000-587559	A3	20000605
WO 2000-US40103	W	20000605

AB Methods and compns. for modulating the FGF effect on the sensitivity of malignant and normal cells to anticancer agents are provided. In particular, methods and compns. for inhibiting FGF-induced resistance to a broad spectrum of anticancer agents in solid and soft-tissue tumors, metastatic lesions, leukemia and lymphoma are provided. Preferably, the compns. include at least one FGF inhibitor in combination with a cytotoxic agents, e.g., antimicrotubule agents, topoisomerase I inhibitors, topoisomerase II inhibitors, antimetabolites, mitotic inhibitors, alkylating agents, intercalating agents, agents capable of interfering with a signal transduction pathway (e.g., g., a protein kinase C inhibitor, e.g., an anti-hormone, e.g., an antibody against growth factor receptors), an agent that promote apoptosis and/or necrosis, an interferon, an interleukin, a tumor necrosis factor, and radiation. In other embodiments, methods and composition for protecting a cell in a subject, from one or more of killing, inhibition of growth or division or other damage caused, e.g., by a cytotoxic agent, are provided. Preferably, the method includes administering to the subject an effective amount of at least one FGF agonist, thereby treating the cell, e.g., protecting or reducing the damage to the dividing cell from said cytotoxic agent. FGF gene expression-based methods for diagnosis of proliferative disorders are also disclosed.

OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (13 CITINGS)

REFERENCE COUNT: 2 THERE ARE 2 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 74 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:861712 CAPLUS

DOCUMENT NUMBER: 134:25368

TITLE: C-CAM as an angiogenesis inhibitor

INVENTOR(S): Lin, Sue-Hwa; Luo, Weiping; Logothetis, Christopher

PATENT ASSIGNEE(S): Board of Regents, the University of Texas System, USA

SOURCE: PCT Int. Appl., 133 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 2000073340	A1	20001207	WO 2000-US14597	20000526 <--
W: AU, CA, JP				
RW: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
CA 2374990	A1	20001207	CA 2000-2374990	20000526 <--
EP 1183273	A1	20020306	EP 2000-936343	20000526 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
JP 2003501362	T	20030114	JP 2001-500664	20000526 <--
US 6517828	B1	20030211	US 2000-580043	20000526 <--

PRIORITY APPLN. INFO.: US 1999-136563P P 19990528

## ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The present invention relates generally to the fields hyperproliferative disease and angiogenesis. More particularly, the present invention demonstrates that a C-CAM1 cytoplasmic domain is necessary and sufficient for inhibiting angiogenesis. In particular embodiments, it relates to inhibiting hyperproliferative cell growth by administering to a cell a C-CAM1 cytoplasmic domain or an expression construct encoding a C-CAM1 cytoplasmic domain. In other embodiments, angiogenesis is inhibited by administering to a subject a C-CAM1 polypeptide or an expression construct encoding a C-CAM1 polypeptide.

REFERENCE COUNT: 8 THERE ARE 8 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 75 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:784015 CAPLUS

DOCUMENT NUMBER: 134:320585

TITLE: Involvement of caspases in 5-FU induced apoptosis in an oral cancer cell line

AUTHOR(S): Ohtani, Tadashi; Hatori, Masashi; Ito, Hidetoshi; Takizawa, Kunio; Kamiyo, Ryutaro; Nagumo, Masao

CORPORATE SOURCE: Second Department of Oral and Maxillofacial Surgery, Showa University, Tokyo, 145-8515, Japan

SOURCE: Anticancer Research (2000), 20(5A), 3117-3121

CODEN: ANTRD4; ISSN: 0250-7005

PUBLISHER: International Institute of Anticancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Although many anticancer drugs have been reported to induce apoptosis in cancer cells, the underlying mechanism remains unclear. Recent studies have revealed that the caspase family of cysteine proteases have been shown to play an important role in the regulation of several apoptotic processes. Thus, the present study investigated whether apoptosis induced by anticancer drugs is mediated by the activation of caspase cascade. NA cells, a squamous cell carcinoma cell line, were exposed to cisplatin (CDDP) or 5-fluorouracil (5-FU) with or without inhibitors of caspase 1, 3 and 8. Anal. of DNA fragmentation revealed that caspase inhibitors consistently inhibited DNA fragmentation induced by 5-FU. During the early stages of apoptosis, phosphatidylserine (PS) is translocated from the inner side of the plasma membrane to the cell surface. This PS externalization was markedly inhibited by treatment with caspase-8 inhibitor. These findings suggested that 5-FU induced apoptosis was mediated by the activation of a caspase cascade involving caspase 1, 3 and 8.

OS.CITING REF COUNT: 17 THERE ARE 17 CAPLUS RECORDS THAT CITE THIS RECORD (17 CITINGS)

REFERENCE COUNT: 21 THERE ARE 21 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 76 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:717375 CAPLUS

DOCUMENT NUMBER: 134:275301

TITLE: Dose and time dependencies of 5-fluorouracil pharmacokinetics

AUTHOR(S): Terret, Catherine; Erdociain, Eric; Guimbaud, Rosine; Boisdron-Celle, Michele; McLeod, Howard L.;

Fety-Deporte, Regine; Lafont, Thierry; Gamelin, Erick;

Bugat, Roland; Canal, Pierre; Chatelut, Etienne

CORPORATE SOURCE: Institut Claudius-Regaud and Universite Paul-Sabatier, Toulouse, Fr.

SOURCE: Clinical Pharmacology & Therapeutics (St. Louis) ( 2000), 68(3), 270-279

CODEN: CLPTAT; ISSN: 0009-9236

PUBLISHER: Mosby, Inc.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Objectives: The purpose of this study was to examine the interpatient and inpatient variability of the Michaelis-Menten plasma parameters of 5-fluorouracil administered according to a schedule combining a bolus of 400 mg/m<sup>2</sup> followed by 22-h infusion of 600 mg/m<sup>2</sup> for 2 consecutive days. Patients: A pharmacokinetic population approach was used to analyze the data from 21 patients with colorectal cancer. Results: The 5-fluorouracil plasma concns. vs. time were best described by a two-compartment model with nonlinear elimination from the central compartment. The relationships between the pharmacokinetic parameters and patient characteristics were tested. On day 1 the mean values (with interindividual variability as expressed by the coefficient of variation) were 1390 mg · h<sup>-1</sup> (20%), and 5.57 mg · L<sup>-1</sup> (22%) for the maximum rate of elimination, and the half-saturating plasma concentration. The maximum rate of elimination was pos. correlated to the body surface area and the percentage of liver involvement by metastatic disease determined by tomodesitometric examination. The model was successfully tested with independent data sets corresponding to other schedules. The anal. of this inpatient variability showed that the half-saturating plasma concentration increased from day 1 to day 2, especially in the patients with low lymphocyte cell dihydropyrimidine dehydrogenase activity. Conclusion: The pharmacokinetic parameters obtained in this study would be useful to predict the 5-fluorouracil plasma concns. following other schedules of administration of 5-fluorouracil and to study the possible pharmacokinetic interactions between 5-fluorouracil and other drugs.

OS.CITING REF COUNT: 17 THERE ARE 17 CAPLUS RECORDS THAT CITE THIS RECORD (17 CITINGS)

REFERENCE COUNT: 36 THERE ARE 36 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 77 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:569908 CAPLUS

DOCUMENT NUMBER: 133:168401

TITLE: Therapeutic formulations containing hyaluronic acid  
INVENTOR(S): Falk, Rudolf Edgar; Asculai, Samuel Simon; Klein, Ehud Shmuel; Harper, David William; Hochman, David; Purschke, Don

PATENT ASSIGNEE(S): Hyal Pharmaceutical Corporation, Can.

SOURCE: U.S., 45 pp., Cont.-in-part of U.S. Ser. No. 675,908.

CODEN: USXXAM

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 24

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6103704	A	20000815	US 1993-18754	19930217 <--
CZ 288292	B6	20010516	CZ 1990-4598	19900921 <--
US 6069135	A	20000530	US 1991-675908	19910703 <--
US 5639738	A	19970617	US 1992-838675	19920221 <--
US 5827834	A	19981027	US 1994-286263	19940805 <--
US 5811410	A	19980922	US 1995-465335	19950605 <--
US 5830882	A	19981103	US 1995-462615	19950605 <--
US 5852002	A	19981222	US 1995-462147	19950605 <--

US 5972906	A	19991026	US 1995-503919	19950719 <--
US 6194392	B1	20010227	US 1995-460978	19950807 <--
CA 2268476	A1	19980430	CA 1996-2268476	19961018 <--
WO 9817320	A1	19980430	WO 1996-CA700	19961018 <--
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE,				
ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS,				
LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,				
SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR,				
IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML,				
MR, NE, SN, TD, TG				
AU 9672721	A	19980515	AU 1996-72721	19961018 <--
AU 739701	B2	20011018		
EP 952855	A1	19991103	EP 1996-934250	19961018 <--
EP 952855	B1	20050727		
R: DE, FR, GB, IT, SE				
NZ 335259	A	20001222	NZ 1996-335259	19961018 <--
ZA 9608847	A	19970527	ZA 1996-8847	19961022 <--
IN 1996CA01848	A	20050304	IN 1996-CA1848	19961023
US 6475795	B1	20021105	US 1997-860696	19970616 <--
US 20030036525	A1	20030220	US 2002-234355	20020904 <--
PRIORITY APPLN. INFO.:				
			US 1991-675908	A2 19910703
			US 1992-838675	A2 19920221
			CA 1989-612307	A 19890921
			WO 1990-CA306	W 19900918
			CS 1990-4598	A 19900921
			CA 1992-2061566	A 19920220
			US 1992-838674	B2 19920221
			US 1993-18508	A2 19930217
			US 1993-18754	A2 19930217
			US 1994-290848	A2 19940819
			US 1994-290840	A2 19941027
			WO 1996-CA700	A 19961018
			US 1997-860696	A1 19970616
ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT				
AB A method of accumulating a drug and a form of hyaluronic acid in the skin and/or exposed tissue of a human includes topically administering a therapeutically effective dosage amount of a formulation which comprises at least 5 mg/cm <sup>2</sup> of the form of hyaluronic acid and a therapeutically effective amount of the drug. A topical gel contained glycerin 3, benzyl alc. 1.5, liquid wax 3, diclofenac sodium 1, sodium hyaluronate 3, and water q.s. 100%. Permeation of diclofenac sodium from the excised human skin was studied.				
OS.CITING REF COUNT:	4	THERE ARE 4 CAPLUS RECORDS THAT CITE THIS RECORD (4 CITINGS)		
REFERENCE COUNT:	344	THERE ARE 344 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT		
L21 ANSWER 78 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN				
ACCESSION NUMBER: 2000:281965 CAPLUS				
DOCUMENT NUMBER: 151:366909				
TITLE: Method for automatic dosing of drugs for controlled microdelivery				
INVENTOR(S): Jacobsen, Stephen C.; Zentner, Gaylen M.				
PATENT ASSIGNEE(S): Sarcos Lc, USA				
SOURCE: U.S., 21pp., Cont. -in-part of U.S. 5,782,799.				
CODEN: USXXAM				
DOCUMENT TYPE: Patent				
LANGUAGE: English				
FAMILY ACC. NUM. COUNT: 1				



PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
US 6056734	A	20000502	US 1998-97950	19980616 <--
PRIORITY APPLN. INFO.:			US 1997-797296	A2 19970207

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB The method for automatic dosing of drugs utilizes a microdelivery device which may be implanted in or otherwise administered to an animal or human. A microdelivery device is configured to have at least one compartment containing at least one drug so that a plurality of doses of the drug(s) are held within the device. In accordance with the present invention, the microdelivery device selectively actuates a compartment to selectively release doses of the drug(s) to provide an efficacious dosing pattern. One primary function of the present invention is to release two or more pesticides in such a pattern that parasites are effectively controlled while preventing the development of tolerance to the drugs within the parasites. Preferably, the microdelivery device is programmable to effectuate the release of the drug(s) at a desired time to maintain efficacious levels of the drug while minimizing the amount of drug which must be used. More particularly, the present invention relates to a method for using electromech. mechanisms and micromachines for dosing of drugs to maximize the effectiveness of the drugs and to prevent the development of drug tolerance and resistance. Thus, the electromech. microdelivery system is used to automatically administer the anticoagulant enoxaparin sodium for prevention of deep vein thrombosis which may lead to pulmonary embolism; the electromech. micropump is programmed to deliver 0.6 mL/day of a sterile solution containing 60 mg of enoxaparin sodium; the

drug is administered s.c. in two divided doses through either an indwelling catheter or freshly inserted small gauge hypodermic needle; typically, the doses are spaced every twelve hours; sufficient drug solution is contained in the attached drug reservoir for 2 to 6 doses (1 to 3 days) depending on local medical protocol for change-out of infusion sets; once the reservoir is exhausted, the entire assembly is discarded and a new assembly is positioned and switched on; the duration of use of a given drug delivery device is presently limited by the potency period of the indwelling catheter; as advances occur in indwelling catheter technol. that permit longer duration catheter use, the drug delivery technol. is fully capable of unattended use for periods of several months; the automation of parenteral anticoagulant drug delivery using small, lightweight, inexpensive electromech. micropumps that provide instrument level precision and accuracy permits patients to leave high-cost hospital environments and return home without compromising the quality of therapy.

OS.CITING REF COUNT: 5 THERE ARE 5 CAPLUS RECORDS THAT CITE THIS RECORD (5 CITINGS)

REFERENCE COUNT: 16 THERE ARE 16 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 79 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:135476 CAPLUS

DOCUMENT NUMBER: 132:160967

TITLE: The predictive value of body protein for chemotherapy-induced toxicity

AUTHOR(S): Aslani, Alireza; Smith, Ross C.; Allen, Barry J.; Pavlakis, Nicholas; Levi, John A.

CORPORATE SOURCE: Center for In Vivo Body Composition Studies, Royal North Shore Hospital, St. Leonards, Australia

SOURCE: Cancer (New York) (2000), 88(4), 796-803  
CODEN: CANCAR; ISSN: 0008-543X

PUBLISHER: John Wiley & Sons, Inc.

DOCUMENT TYPE: Journal  
LANGUAGE: English

AB The use of body surface area in determining chemotherapy dosing, particularly in the obese, remains controversial. Total body nitrogen (TBN) measurement in patients with serious illness has been suggested to be an accurate predictor of clin. course. The ability of TBN to predict chemotherapy-induced neutropenia was examined in the current study. TBN measurements were performed in 31 female outpatients with breast carcinoma who were undergoing standard cyclophosphamide, methotrexate, and 5-fluorouracil (CMF)-based chemotherapy (median age, 48 yr; range, 26-77 yr). TBN was measured using the in vivo neutron capture anal. technique on Day 1 of Cycles 2-6. The chemotherapy toxicity index used was the absolute neutrophil count nadir (ANCN). Neutropenia was defined as an ANCN < 1.0+109/L. The nitrogen index (NI) (TBN expressed as a percentage of age-, gender-, and height-matched healthy patients) then was compared with the corresponding ANCN values. Using receiver operating characteristics anal., a "cut-off" value of NI = 0.89 was found. In this group of patients, when the NI was < 0.89, 11 of 13 courses in 7 patients (85%) led to an ANCN of < 1.0+109/L, and when the NI was > 0.89, 29 of 109 courses (27%) led to an ANCN of < 1.0+109/L (P < 0.0001). In this small group of breast carcinoma patients, the NI was found to be the most powerful predictor of neutropenia after CMF-based chemotherapy. The authors conclude that NI may be a useful clin. tool in identifying patients at a higher risk of chemotherapy-induced toxicity when widely distributed drug combinations such as CMF are used, and warrants further study with other commonly used drugs or drug regimens.

OS.CITING REF COUNT: 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

REFERENCE COUNT: 46 THERE ARE 46 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 80 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 2000:84648 CAPLUS

DOCUMENT NUMBER: 132:141941

TITLE: Conjugates and fusion proteins for treating secondary tissue damage and other inflammatory conditions and disorders

INVENTOR(S): McDonald, John R.; Coggins, Philip J.

PATENT ASSIGNEE(S): Osprey Pharmaceuticals Limited, Can.

SOURCE: PCT Int. Appl., 204 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 2

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	-----	-----	-----	-----
WO 2000004926	A2	20000203	WO 1999-CA659	19990721 <--
WO 2000004926	A3	20001102		
W:	AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZA, ZW			
RW:	GH, GM, KE, LS, MW, SD, SL, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
CA 2335105	A1	20000203	CA 1999-2335105	19990721 <--
AU 9948918	A	20000214	AU 1999-48918	19990721 <--
EP 1098664	A2	20010516	EP 1999-932572	19990721 <--

EP 1098664	B1	20030806		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO				
JP 2002521019	T	20020716	JP 2000-560919	19990721 <--
JP 4454152	B2	20100421		
AT 246517	T	20030815	AT 1999-932572	19990721 <--
EP 1346731	A1	20030924	EP 2003-76150	19990721 <--
EP 1346731	B1	20061206		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, SI, LT, LV, FI, RO, MK, CY, AL				
ES 2205849	T3	20040501	ES 1999-932572	19990721 <--
AT 347378	T	20061215	AT 2003-76150	19990721
ES 2275999	T3	20070616	ES 2003-76150	19990721
US 7157418	B1	20070102	US 1999-360242	19990722
US 7166702	B1	20070123	US 1999-453851	19991202
US 20020168370	A1	20021114	US 2001-792793	20010222 <--
US 7192736	B2	20070320		
HK 1037133	A1	20031107	HK 2001-107546	20011030 <--
US 20030215421	A1	20031120	US 2003-375209	20030224 <--
AU 2004202331	A1	20040624	AU 2004-202331	20040527 <--
US 20060198820	A1	20060907	US 2006-361977	20060224
AU 2007200061	A1	20070125	AU 2007-200061	20070105
AU 2007200061	B2	20070920		
AU 2007201749	A1	20070510	AU 2007-201749	20070419
AU 2007201749	B2	20080529		
AU 2007201753	A1	20070510	AU 2007-201753	20070419
AU 2007201755	A1	20070510	AU 2007-201755	20070419
AU 2007201755	B2	20070920		
AU 2007201759	A1	20070517	AU 2007-201759	20070419
AU 2007201759	B2	20080529		
JP 2010075194	A	20100408	JP 2009-264178	20091119
PRIORITY APPLN. INFO.:				
US 1998-120523 A2 19980722				
US 1998-155186P P 19980722				
AU 1999-48918 A3 19990721				
EP 1999-932572 A3 19990721				
JP 2000-560919 A3 19990721				
WO 1999-CA659 W 19990721				
US 1999-360242 A3 19990722				
US 1999-453851 A3 19991202				
US 2001-792793 A1 20010222				
US 2003-375209 A1 20030224				
AU 2004-202331 A3 20040527				
AU 2007-200061 A3 20070105				

# ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Conjugates containing as a ligand a chemokine receptor-targeting agent, such as chemokines, and a targeted agent, such as a toxin are provided. These conjugates are used to treat inflammatory responses associated with activation, proliferation and migration of immune effector cells, including leukocyte cell types, neutrophils, macrophages, and eosinophils. The conjugates provided herein are used to lessen or inhibit these processes to prevent or at least lessen the resulting secondary effects. In particular, the conjugates are used to target toxins to receptors on secondary tissue damage-promoting cells. The ligand moiety can be selected to deliver the cell toxin to such secondary tissue damage-promoting cells as mononuclear phagocytes, leukocytes, natural killer cells, dendritic cells, and T and B lymphocytes, thereby suppressing the proliferation, migration, or physiologic activity of such cells. Among preferred conjugates are fusion proteins having a chemokine, or a biologically active fragment thereof, as the ligand moiety linked to a cell toxin via a peptide linker of from 2 to about 60 amino acid residues.

OS.CITING REF COUNT: 13 THERE ARE 13 CAPLUS RECORDS THAT CITE THIS

REFERENCE COUNT: 10 RECORD (19 CITINGS)  
THERE ARE 10 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 81 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 2000:14232 CAPLUS  
DOCUMENT NUMBER: 132:273960  
TITLE: Role for  $\alpha$ 1,2-fucosyltransferase and histo-blood  
group antigen H type 2 in resistance of rat colon  
carcinoma cells to 5-fluorouracil  
AUTHOR(S): Cordel, Sandrine; Goupille, Caroline; Hallouin,  
Florence; Meflah, Khaled; Le Pendu, Jacques  
CORPORATE SOURCE: INSERM U419, Institute of Biology, Nantes, F-44035,  
Fr.  
SOURCE: International Journal of Cancer (2000),  
85(1), 142-148  
CODEN: IJCNAM; ISSN: 0020-7136  
PUBLISHER: Wiley-Liss, Inc.  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB 5-Fluorouracil (5-FU) is a drug of standard use in chemotherapy of colon  
carcinoma. However, its efficacy is limited by inherent and  
acquired cell resistance. Major changes in histo-blood group antigenic  
expression, at times associated with poor prognosis, occur on colon cancer  
cells. To assess whether these antigens might play a role in the  
resistance to 5-FU, a rat model of colon carcinoma was used. We  
observed that in vivo treatment of tumors with the drug increased expression  
of antigen H type 2. The increase was also observed after in vitro  
short-term exposure to 5-FU, as well as on a cell-resistant variant  
selected by continuous exposure to the drug, and was accompanied by an  
increase in  $\alpha$ 1,2-fucosyltransferase activity, the key enzyme  
involved in synthesis of H antigens. Transfection of cells devoid of this  
enzymic activity by an  $\alpha$ 1,2-fucosyltransferase cDNA allowed  
expression of H type 2 antigen and increased resistance to 5-FU.  
Inversely, transfection of cells which possess enzymic activity by a cDNA  
in anti-sense orientation reduced both H type 2 cell-surface  
antigen and resistance to the drug. These results demonstrate that, in  
this exptl. model,  $\alpha$ 1,2-fucosyltransferase and H type 2 antigen are  
involved in cellular resistance to 5-FU.  
OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD  
(9 CITINGS)  
REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 82 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1999:733436 CAPLUS  
DOCUMENT NUMBER: 132:245934  
TITLE: Vaginal 5-fluorouracil for high-grade cervical  
dysplasia in human immunodeficiency virus infection: a  
randomized trial  
AUTHOR(S): Maiman, M.; Watts, D. H.; Andersen, J.; Clax, P.;  
Merino, M.; Kendall, M. A.  
CORPORATE SOURCE: Division of AIDS, Adult AIDS Clinical Trials Group,  
National Institute of Allergy and Infectious Diseases,  
Bethesda, MD, USA  
SOURCE: Obstetrics & Gynecology (New York) (1999),  
94(6), 954-961  
CODEN: OBGNAS; ISSN: 0029-7844  
PUBLISHER: Elsevier Science Inc.  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB Objective: To compare the efficacy and toxicity of topical vaginal 5-fluorouracil (5-FU) maintenance therapy against the effects of observation after standard treatment for high-grade cervical dysplasia in human immunodeficiency virus (HIV)-infected women and to evaluate the association between baseline CD4 count and time to recurrence. Methods: In a phase III unmasked, randomized, multicenter, outpatient clin. trial, 101 HIV-pos. women either received 6 mo of biweekly treatment with vaginal 5-FU cream (2 g) or underwent 6 mo of observation after standard excisional or ablative cervical treatment for cervical intraepithelial neoplasia (CIN). Papanicolaou smears and colposcopy were scheduled at regular intervals during the ensuing 18 mo, with the primary end point being the time at which CIN of any grade recurred. Results: Thirty-eight percent of women developed recurrence: 14 (28%) of 50 in the 5-FU therapy group and 24 (47%) of 51 in the observation group. Treatment with 5-FU was significantly associated with prolonged time to CIN development ( $P = .04$ ). Observation subjects were more likely to have high-grade recurrences, with 31% developing CIN 2-3 compared with 8% in the 5-FU treatment arm ( $P = .014$ ), and disease recurred more quickly in observation subjects as well. Baseline CD4 count was related significantly to time to recurrence ( $P = .04$ ), with 46% of subjects with CD4 counts less than 200 cells/mm<sup>3</sup> developing recurrence compared with 33% of subjects with CD4 counts at least 200 cells/mm<sup>3</sup>. Disease recurred more slowly in subjects who had received antiretroviral therapy than in antiretroviral therapy-naïve subjects. There were no instances of grade 3 or 4 toxicity, and compliance with 5-FU treatment was generally good. Conclusion: Adjunctive maintenance intravaginal 5-FU therapy after standard surgery for high-grade lesions safely and effectively reduced recurrence of cervical intraepithelial neoplasia in HIV-infected women.

OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)

REFERENCE COUNT: 20 THERE ARE 20 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 83 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1999:506037 CAPLUS

DOCUMENT NUMBER: 132:73298

TITLE: Fluorouracil releasing pattern from fluorouracil-polyglycolic acid composite in the peritoneal cavity of rat

AUTHOR(S): Noh, Seung-Moo; Chung, Kyeong-Soo; Oh, Jung-Yeon; Kim, Jin-Hyang; Yang, Joon-Mook; Kang, Dae-Young; Song, Kyu-Sang; Choi, Jung-Mok; Choi, Sun-Woong; Lee, Jin-Ho; Cho, June-Sik; Min, Byung-Moo; Kim, Yong-Baek; Kim, Chang-Sik; Park, Keun-Sung; Kim, Seung-Young; Kim, Hak-Yong; In, Hyun-Bin

CORPORATE SOURCE: Dep. General Surgery, Chungnam Natl. Univ., S. Korea

SOURCE: Chungnam Uidae Chapchi (1998), 25(1), 39-46

CODEN: CUCHDS; ISSN: 0253-6307

PUBLISHER: Chungnam National University, College of Medicine

DOCUMENT TYPE: Journal

LANGUAGE: Korean

AB A common form of relapse in adenocarcinoma of the stomach is i.p. dissemination, in fact, among gastric adenocarcinoma patients who have undergone surgery intended to cure, approx. 50% of the patients develop initial recurrence in the peritoneal cavity regardless of the anat. site of the primary tumor within the stomach. The efficacy of systemic postoperative chemotherapy to prevent peritoneal recurrence of gastric adenocarcinoma is not satisfactory. There is still a great need for improved therapeutic strategies on the disseminated microscopic disease and small miliary nodules remaining on the peritoneal surface or lymphatics after operation. The authors have made

fluorouracil-polyglycolic acid composite disks (Fu-PGA disks) with fluorouracil and biodegradable polymer: polyglycolic acid for more effective i.p. chemotherapy. We inserted the Fu-PGA disk(s) in the peritoneal cavity of rat and pharmacokinetic study was performed to measure fluorouracil concentration in the peritoneal fluid, plasma, liver, kidney and heart tissue at 24 h, 72 h and 168 h after insertion of Fu-PGA disk(s). Myelosuppressive action of this composite also was determined following its administration. The data of this study suggested that Fu-PGA composite will be a new device releasing drugs in a controlled manner and having target-ability to peritoneum, and this device will be improving the efficacy of i.p. chemotherapy for gastric adenocarcinoma.

L21 ANSWER 84 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1999:348369 CAPLUS

DOCUMENT NUMBER: 130:334740

TITLE: Alterations of intratumoral pharmacokinetics of 5-fluorouracil in head and neck carcinoma during simultaneous radiochemotherapy

AUTHOR(S): Schlemmer, Heinz-Peter; Becker, Markus; Bachert, Peter; Dietz, Andreas; Rudat, Volker; Vanselow, Bernhard; Wollensack, Petra; Zuna, Iwan; Knopp, Michael V.; Weidauer, Hagen; Wannenmacher, Michael; Van Kaick, Gerhard

CORPORATE SOURCE: Research Program Radiological Diagnostics and Therapy, German Cancer Research Center (Deutsches Krebsforschungszentrum), University of Heidelberg, Heidelberg, 69120, Germany

SOURCE: Cancer Research (1999), 59(10), 2363-2369

CODEN: CNREA8; ISSN: 0008-5472

PUBLISHER: AACR Subscription Office

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The kinetics of local drug uptake and metabolism of the anticancer drug 5-fluorouracil (5-FU) has been monitored by means of <sup>19</sup>F NMR spectroscopy in 17 patients with neck tumors during concurrent radiochemotherapy. All of the patients underwent an accelerated hyperfractionated, concomitant-boost radiochemotherapy with 5-FU (600 or 1000 mg/m<sup>2</sup> of body surface (b.s.)) and carboplatin (70 mg/m<sup>2</sup> of b.s.). Serial <sup>19</sup>F NMR spectra were obtained during and after the administration of 5-FU in a 1.5-T scanner with the use of a 5-cm diameter surface coil positioned on a cervical lymph node metastasis. Exams. were performed at day 1 of therapy and, in 13 patients, also after 43.5 Gy of irradiation at day 1 of the second chemotherapy cycle. Resonances of 5-FU and the catabolites 5,6-dihydro-5-fluorouracil (DHFU) and  $\alpha$ -fluoro- $\beta$ -alanine (FBAL) were resolved in the tumor spectra. The median of the 5-FU and FBAL levels was significantly higher (more than 2-fold) at the second compared with the first examination, whereas the level of DHFU did not change. This effect could indicate an increased delivery of 5-FU into the interstitial space of the tumor in the course of the combined treatment, which would result in an enhanced exposure of the tumor cells to the drug. A potential mechanism for synergy between radio- and chemotherapy is discussed, but alternative mechanisms are also being considered. The findings indicate that a method is available to rationally address the design of dosing schedules in concurrent therapy regimens.

OS.CITING REF COUNT: 28 THERE ARE 28 CAPLUS RECORDS THAT CITE THIS RECORD (28 CITINGS)

REFERENCE COUNT: 32 THERE ARE 32 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 85 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1999:276365 CAPLUS  
 DOCUMENT NUMBER: 130:322426  
 TITLE: Concurrent cisplatin-based radiotherapy and chemotherapy for locally advanced cervical cancer  
 AUTHOR(S): Rose, Peter G.; Bundy, Brian N.; Watkins, Edwin B.; Thigpen, J. Tate; Deppe, Gunther; Maiman, Mitchell A.; Clarke-Pearson, Daniel L.; Insalaco, Sam  
 CORPORATE SOURCE: Division of Gynecologic Oncology, Department of Reproductive Biology, University Hospitals of Cleveland and Case Western Reserve University, Cleveland, USA  
 SOURCE: New England Journal of Medicine (1999), 340(15), 1144-1153  
 CODEN: NEJMAG; ISSN: 0028-4793  
 PUBLISHER: Massachusetts Medical Society  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB On behalf of the Gynecol. Oncol. Group, we performed a randomized trial of radiotherapy in combination with three concurrent chemotherapy regimens - cisplatin alone; cisplatin, fluorouracil, and hydroxyurea; and hydroxyurea alone - in patients with locally advanced cervical cancer. Women with primary untreated invasive squamous-cell carcinoma, adenocarcinoma, or adenocarcinoma of the cervix of stage IIB, III, or IVA, without involvement of the para-aortic lymph nodes, were enrolled. The patients had to have a leukocyte count of at least 3000 per cubic millimeter, a platelet count of at least 100,000 per cubic millimeter, a serum creatinine level no higher than 2 mg per dL (177 µmol per L), and adequate hepatic function. All patients received external-beam radiotherapy according to a strict protocol. Patients were randomly assigned to receive one of three chemotherapy regimens: 40 mg of cisplatin per square meter of body-surface area per wk for six weeks (group 1); 50 mg of cisplatin per square meter on days 1 and 29, followed by 4 g of fluorouracil per square meter given as a 96-h infusion on days 1 and 29, and 2 g of oral hydroxyurea per square meter twice weekly for six weeks (group 2); or 3 g of oral hydroxyurea per square meter twice weekly for six weeks (group 3). The anal. included 526 women. The median duration of follow-up was 35 mo. Both groups that received cisplatin had a higher rate of progression-free survival than the group that received hydroxyurea alone ( $P < 0.001$  for both comparisons). The relative risks of progression of disease or death were 0.57 (95 % confidence interval, 0.42 to 0.78) in group 1 and 0.55 (95 % confidence interval, 0.40 to 0.75) in group 2, as compared with group 3. The overall survival rate was significantly higher in groups 1 and 2 than in group 3, with relative risks of death of 0.61 (95 % confidence interval, 0.44 to 0.85) and 0.58 (95 % confidence interval, 0.41 to 0.81), resp. Regimens of radiotherapy and chemotherapy that contain cisplatin improve the rates of survival and progression-free survival among women with locally advanced cervical cancer.  
 OS.CITING REF COUNT: 266 THERE ARE 266 CAPLUS RECORDS THAT CITE THIS RECORD (266 CITINGS)  
 REFERENCE COUNT: 64 THERE ARE 64 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 86 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1999:262135 CAPLUS  
 DOCUMENT NUMBER: 130:276741  
 TITLE: Compositions and methods for the treatment of primary and metastatic neoplastic diseases using arsenic compounds

INVENTOR(S): Ellison, Ralph M.; Mermelstein, Fred H.  
 PATENT ASSIGNEE(S): Polarx Biopharmaceuticals, Inc., USA  
 SOURCE: PCT Int. Appl., 58 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9918798	A1	19990422	WO 1998-US21782	19981015 <--
W: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, UZ, VN, YU, ZW				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
CA 2307208	A1	19990422	CA 1998-2307208	19981015 <--
AU 9910893	A	19990503	AU 1999-10893	19981015 <--
AU 751932	B2	20020829		
EP 1022951	A1	20000802	EP 1998-953552	19981015 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
BR 9813085	A	20000822	BR 1998-13085	19981015 <--
TR 2000001959	T2	20001121	TR 2000-1959	19981015 <--
NZ 503973	A	20010928	NZ 1998-503973	19981015 <--
JP 2001519366	T	20011023	JP 2000-515442	19981015 <--
US 20020183385	A1	20021205	US 1998-173531	19981015 <--
US 6875451	B2	20050405		
EP 1374875	A2	20040102	EP 2003-19595	19981015 <--
EP 1374875	A3	20040107		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1378240	A2	20040107	EP 2003-19594	19981015 <--
EP 1378240	A3	20040114		
EP 1378240	B1	20080910		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1378241	A1	20040107	EP 2003-19629	19981015 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1391206	A1	20040225	EP 2003-19628	19981015 <--
EP 1391206	B1	20080702		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1419778	A1	20040519	EP 2003-29713	19981015 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1466607	A1	20041013	EP 2004-7847	19981015 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI, CY				
EP 1621077	A1	20060201	EP 2005-77338	19981015
EP 1621077	B1	20080924		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO, MK, CY, AL				
AT 399560	T	20080715	AT 2003-19628	19981015
PT 1391206	E	20080813	PT 2003-19628	19981015
AT 407683	T	20080915	AT 2003-19594	19981015



AT 409043	T	20081015	AT 2005-77338	19981015
PT 1621077	E	20081104	PT 2005-77338	19981015
PT 1378240	E	20081114	PT 2003-19594	19981015
ES 2309258	T3	20081216	ES 2003-19628	19981015
EP 2018858	A1	20090128	EP 2008-16665	19981015
R: AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LI, LU, MC,				
NL, PT, SE, AL, LT, LV, MK, RO, SI				
ES 2312701	T3	20090301	ES 2003-19594	19981015
ES 2313201	T3	20090301	ES 2005-77338	19981015
NO 2000001977	A	20000613	NO 2000-1977	20000414 <--
MX 2000003655	A	20010731	MX 2000-3655	20000414 <--
HK 1061198	A1	20081017	HK 2004-104292	20010129
HK 1061199	A1	20090403	HK 2004-104293	20010129
AU 2002308909	A1	20030320	AU 2002-308909	20021128 <--
AU 2002308909	B2	20060427		
US 20050100611	A1	20050512	US 2003-640399	20030814
US 7163703	B2	20070116		
US 20050191367	A1	20050901	US 2003-640403	20030814
US 7132116	B2	20061107		
US 20040047916	A1	20040311	US 2003-649944	20030828 <--
US 20040115283	A1	20040617	US 2003-649776	20030828 <--
US 7179493	B2	20070220		
US 20040096518	A1	20040520	US 2003-698625	20031103 <--
US 7205001	B2	20070417		
US 20040161475	A1	20040819	US 2004-776504	20040212 <--
US 20040197420	A1	20041007	US 2004-789604	20040227 <--
US 20040197421	A1	20041007	US 2004-789628	20040227 <--
HK 1085886	A1	20090605	HK 2006-108235	20060725

PRIORITY APPLN. INFO.:

US 1997-62375P	P	19971015
AU 1999-10893	A3	19981015
EP 1998-953552	A3	19981015
EP 2005-77338	A3	19981015
US 1998-173531	A3	19981015
WO 1998-US21782	W	19981015
HK 2001-100623	A3	20010129

ASSIGNMENT HISTORY FOR US PATENT AVAILABLE IN LSUS DISPLAY FORMAT

AB Arsenic compds. are used to treat a variety of neoplastic diseases, including metastatic neoplastic diseases.

OS.CITING REF COUNT: 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD (7 CITINGS)

REFERENCE COUNT: 5 THERE ARE 5 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 87 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1998:794134 CAPLUS

DOCUMENT NUMBER: 130:162776

TITLE: Therapeutic evaluation of compounds in the SCID-RA papillomavirus model

AUTHOR(S): Lobe, David C.; Kreider, John W.; Phelps, William C.

CORPORATE SOURCE: Department of Virology, Glaxo Wellcome, Research

Triangle Park, NC, 27709, USA

SOURCE: Antiviral Research (1998), 40(1-2), 57-71

CODEN: ARSRDR; ISSN: 0166-3542

PUBLISHER: Elsevier Science B.V.

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A previous study by Kreider (Kreider et al., 1979) indicated that rabbit skin, which had been transplanted to immunodeficient nude mice, could be successfully infected with cottontail rabbit papillomavirus (CRPV). The authors have extended this observation in developing a rodent model for evaluation of compds. for activity against the papillomaviruses. In this

model (called the SCID-Ra model), rabbit ear skin is transplanted to the dorsum of SCID mice and allowed to heal for 3 wk. Infection with CRPV by scarification leads to the growth of warty lesions within 2-3 wk in >95% of the animals. Topical and/or systemic therapy can be initiated at various times post infection (PI). Weekly lesion scores are recorded and compds. are evaluated for their ability to suppress wart growth when compared to untreated control mice. Ribavirin, which has had a suppressive effect both in the clinic for the treatment of respiratory papillomatosis and on the growth of warts in the rabbit back model, was evaluated and showed significant anti-proliferative activity with oral dosing. Both antiviral and antiproliferative compds. including podophyllin and 5-fluorouracil, which have been used clin. for the treatment of human papillomavirus (HPV) infections, were evaluated in this model. The anti-mitotic compound, Navelbine (vinorelbine tartrate), which is used for the treatment of non-small cell lung carcinoma was evaluated in this system and showed significant inhibition of wart growth with somewhat less topical cytotoxicity when compared to podophyllotoxin.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)  
 REFERENCE COUNT: 48 THERE ARE 48 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 88 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 1998:634706 CAPLUS

DOCUMENT NUMBER: 129:339412

ORIGINAL REFERENCE NO.: 129:68989a,68992a

TITLE: Intratumoral conversion of 5-fluorocytosine to 5-fluorouracil by monoclonal antibody-cytosine deaminase conjugates: noninvasive detection of prodrug activation by magnetic resonance spectroscopy and spectroscopic imaging

AUTHOR(S): Aboagye, Eric O.; Artemov, Dmitri; Senter, Peter D.; Bhujwalla, Zaver M.

CORPORATE SOURCE: Department of Radiology, Oncology Section, Division of MR Research, The Johns Hopkins University School of Medicine, Baltimore, MD, 21205, USA

SOURCE: Cancer Research (1998), 58(18), 4075-4078  
 CODEN: CNREA8; ISSN: 0008-5472

PUBLISHER: American Association for Cancer Research

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The monitoring of antibody-directed enzyme-prodrug therapies requires evaluation of drug activation within the tissues of interest. We have demonstrated the feasibility of noninvasive magnetic resonance spectroscopy and spectroscopic imaging (chemical shift imaging) to detect activation of the prodrug 5-fluorocytosine (5-FCyt) to the cytotoxic species 5-fluorouracil (5-FU) by monoclonal antibody-cytosine deaminase (CD) conjugates. In vitro, L6-CD but not 1F5-CD selectively metabolized 5-FCyt to 5-FU on H2981 human lung adenocarcinoma cells because of the presence and absence of cell surface L6 and CD20 antigens, resp. After pretreatment of H2981 tumor-bearing mice with L6-CD, in vivo metabolism of 5-FCyt to 5-FU within the tumors was detected by 19F magnetic resonance spectroscopy; the chemical shift separation between 5-FCyt

and 5-FU resonances was approx.1.2 ppm. 5-FU levels were 50-100% of 5-FCyt levels in tumors 10-60 min after 5-FCyt administration. Whole body 19F chemical shift imaging (6+6 mm in-plane resolution) of tumor-bearing mice demonstrated the highest signal intensity of 5-FU within the tumor region. This study supports further development of noninvasive magnetic resonance methods for preclin. and clin. monitoring of CD enzyme-prodrug

therapies.  
 OS.CITING REF COUNT: 30 THERE ARE 30 CAPLUS RECORDS THAT CITE THIS  
 RECORD (30 CITINGS)  
 REFERENCE COUNT: 17 THERE ARE 17 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 89 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1998:527214 CAPLUS  
 DOCUMENT NUMBER: 129:131244  
 ORIGINAL REFERENCE NO.: 129:26693a,26696a  
 TITLE: Method of treating cancer using alkylglycerols in  
 conjunction with chemotherapy  
 INVENTOR(S): Firshein, Richard N.  
 PATENT ASSIGNEE(S): USA  
 SOURCE: PCT Int. Appl., 38 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9832447	A1	19980730	WO 1998-US1411	19980127 <--
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE,				
ES, FI, GB, GE, HU, IL, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS,				
LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD,				
SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN				
RW: GH, GM, KE, LS, MW, SD, SZ, UG, ZW, AT, BE, CH, DE, DK, ES, FI,				
FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM,				
GA, GN, ML, MR, NE, SN, TD, TG				
US 6121245	A	20000919	US 1997-791757	19970129 <--
AU 9862490	A	19980818	AU 1998-62490	19980127 <--
EP 1011685	A1	20000628	EP 1998-904676	19980127 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT,				
IE, FI				
PRIORITY APPLN. INFO.:			US 1997-791757	A1 19970129
			WO 1998-US1411	W 19980127

OTHER SOURCE(S): MARPAT 129:131244  
 AB Tumor cell kill is increased and the sensitivity of tumors to  
 chemotherapeutic agents is increased by the administration of an  
 alkylglycerol together with the chemotherapeutic agent.  
 OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD  
 (2 CITINGS)  
 REFERENCE COUNT: 1 THERE ARE 1 CITED REFERENCES AVAILABLE FOR THIS  
 RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 90 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1998:293396 CAPLUS  
 DOCUMENT NUMBER: 129:3862  
 ORIGINAL REFERENCE NO.: 129:963a  
 TITLE: Enhancement of tumor cell chemosensitivity and  
 radiosensitivity using single chain intracellular  
 antibodies  
 INVENTOR(S): Buchsbaum, Donald J.; Curiel, David T.; Stackhouse,  
 Murray  
 PATENT ASSIGNEE(S): UAB Research Foundation, USA  
 SOURCE: PCT Int. Appl., 95 pp.  
 CODEN: PIXXD2  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English

FAMILY ACC. NUM. COUNT: 4  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9818489	A1	19980507	WO 1997-US19911	19971030 <--
W: AU, CA, JP				
RW: AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE				
AU 9852431	A	19980522	AU 1998-52431	19971030 <--
PRIORITY APPLN. INFO.:			US 1996-29673P	P 19961030
			WO 1997-US19911	W 19971030

AB The present invention provides a method of enhancing the chemosensitivity and radiosensitivity of a neoplastic cell expressing an oncoprotein that stimulates proliferation of the cell. Abrogation of tumor cell resistance is achieved by transfection with a nucleic acid mol. encoding an scFv antibody homolog, wherein the homolog is expressed intracellularly and binds to the oncoprotein in the endoplasmic reticulum. An intracellular single-chain antibody, directed to the erbB-2 oncoprotein, is shown to down-regulate its surface expression in breast and ovarian carcinoma cells lines and increase the sensitivity to cisplatin and x-irradiation

OS.CITING REF COUNT: 8 THERE ARE 8 CAPLUS RECORDS THAT CITE THIS RECORD (8 CITINGS)

REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 91 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1998:50154 CAPLUS

DOCUMENT NUMBER: 128:175771

ORIGINAL REFERENCE NO.: 128:34495a,34498a

TITLE: Effects of PALA on the pharmacokinetics of 5-fluorouracil

AUTHOR(S): Nassim, Mark Adel; Rouini, Mohammad R.; Cripps, M. Christine; Shirazi, Farshad H.; Veerasinghan, Shereeni; Molepo, J. Matshela; Obrocea, Micheal; Redmond, Diedre; Bates, Susan; Fry, Diane; Stewart, David J.; Goel, Rakesh

CORPORATE SOURCE: Ottawa Regional Cancer Centre, Ottawa, ON, K1Y 4K7, Can.

SOURCE: Oncology Reports (1998), 5(1), 217-221

CODEN: OCRPEW; ISSN: 1021-335X

PUBLISHER: Oncology Reports

DOCUMENT TYPE: Journal

LANGUAGE: English

AB N-(phosphonacetyl)-L-aspartate (PALA) modulates the activity of 5-fluorouracil (5-FU) by inhibiting pyrimidine biosynthesis. A cross-over study was conducted to determine whether PALA affects the pharmacokinetic parameters of 5-FU in patients given 5-FU/folinic acid (FA). Six patients (3 males, 3 females) aged 63±4.3 (mean ± SD) years (body surface area of 1.84±18 m2) with metastatic colorectal carcinoma were given two courses of treatment. The treatment consisted of 250 mg/m2 of PALA on day 1 followed by 20 mg/m2 FA and 400 mg/m2 5-FU (5 min i.v. bolus injection) on days 2-5 in one cycle of treatment (PALA+). In another treatment cycle, these doses of 5-FU and FA were given for all 5 days without PALA (PALA-). The two courses were given four weeks apart. It was determined by random selection whether the course with PALA was given before or after the course without PALA. Blood samples were collected over a period of three hours, starting from the beginning of 5-FU infusion on days 2 and 5 of both courses. Plasma concns. of 5-FU were determined by an HPLC technique. Pharmacokinetic parameters were calculated using a non-compartmental model. While there were

no significant differences between pharmacokinetic parameters in the PALA+ vs PALA- courses, there was a trend towards a decreasing area under the curve (AUC) and increasing clearance (Cl) in PALA+ courses of treatment.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

REFERENCE COUNT: 31 THERE ARE 31 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 92 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1997:511962 CAPLUS

DOCUMENT NUMBER: 127:117382

ORIGINAL REFERENCE NO.: 127:22505a,22508a

TITLE: Oxidized glutathione, salts, and derivatives as enhancers of endogenous production of cytokines and hemopoietic factors, and methods of therapeutic use

INVENTOR(S): Balazovsky, Mark Borisovich; Kozhemyakin, Leonid Andreevich

PATENT ASSIGNEE(S): Balazovsky, Mark Borisovich, Russia; Kozhemyakin, Leonid Andreevich

SOURCE: PCT Int. Appl., 125 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 3

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
WO 9721444	A1	19970619	WO 1996-RU340	19961210 <--
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
RU 2089179	C1	19970910	RU 1995-120403	19951214 <--
WO 9721443	A1	19970619	WO 1996-RU226	19960808 <--
W: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, US, UZ, VN, RU				
RW: KE, LS, MW, SD, SZ, UG, AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG				
AP 928	A	20010115	AP 1998-1260	19961201 <--
W: KE, LS, MW, SD, SZ, UG				
AU 9711130	A	19970703	AU 1997-11130	19961210 <--
EP 869809	A1	19981014	EP 1996-941915	19961210 <--
EP 869809	B1	20020327		
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, FI				
RU 2153351	C2	20000727	RU 1998-108088	19961210 <--
JP 2000515111	T	20000114	JP 1997-521965	19961210 <--
AT 214936	T	20020415	AT 1996-941915	19961210 <--
US 6492329	B1	20021210	US 2000-702701	20001031 <--
PRIORITY APPLN. INFO.:			RU 1995-120403	A 19951214
			WO 1996-RU226	A 19960808
			US 1996-733886	A 19961018
			WO 1996-RU340	A 19961210
			US 1996-766557	A 19961211

AB A method for stimulating endogenous production of cytokines and hemopoietic factors comprises topical or parenteral administration of an effective amount of oxidized glutathione, and/or a pharmaceutically acceptable salt and/or derivative thereof, for a period sufficient to stimulate the endogenous production to obtain a therapeutic effect. The oxidized glutathione and/or pharmaceutically acceptable salt and/or derivative is introduced along with an extender of their half life. The compds. of the invention may be used in the treatment of neoplasms, immune diseases, etc.

OS.CITING REF COUNT: 7 THERE ARE 7 CAPLUS RECORDS THAT CITE THIS RECORD  
(7 CITINGS)  
REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS  
RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 93 OF 124 CAPLUS COPYRIGHT 2010 ACS on SIN

ACCESSION NUMBER: 1997:404609 CAPLUS

DOCUMENT NUMBER: 127:75654

ORIGINAL REFERENCE NO.: 127:14273a

TITLE: Sensitization of cancer cells treated with cytotoxic drugs to Fas-mediated cytotoxicity

AUTHOR(S): Micheau, Olivier; Solary, Eric; Hammann, Arlette; Martin, Francois; Dimanche-Boitrel, Marie-Therese

CORPORATE SOURCE: Unite de Formation et de Recherche de Medecine, Contrat Jeune Formation de l'Institut National de la Sante et de la Recherche Medicale (INSERM) 94-08, Dijon, 21033, Fr.

SOURCE: Journal of the National Cancer Institute (1997), 89(11), 783-789

CODEN: JNCIEQ; ISSN: 0027-8874

PUBLISHER: Oxford University Press

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The transmembrane receptor Fas, together with its protein-binding partner (Fas ligand), is a key regulator of programmed cell death (i.e., apoptosis). Fas and Fas ligand also influence the ability of cytotoxic T lymphocytes and natural killer cells to eliminate tumor cells. However, by inducing apoptosis in activated T cells, the Fas/Fas ligand system may protect some tumor cells from clearance by the immune system. Anticancer drugs enhance Fas ligand expression on the surface of Fas receptor-expressing leukemia cells, thus suggesting that apoptosis caused by these drugs may be mediated via the Fas/Fas ligand system. This study was conducted to further investigate the relationship between the modulation of Fas receptor gene and protein expression by treatment of cells with cytotoxic drugs and the immune clearance of tumor cells. Fas expression on human HT29 colon carcinoma cells treated with a variety of anticancer drugs (cisplatin, doxorubicin, mitomycin C, fluorouracil, and camptothecin) was analyzed by use of quant. flow cytometry. Human HCT8R and HCT116 colon carcinoma cells and human U937 leukemia cells were treated with cisplatin only and analyzed in the same way. Fas ligand mRNA and protein levels were studied by use of a reverse transcription-polymerase chain reaction assay and by flow cytometry. Fas gene expression and mRNA levels in cisplatin-treated HT29 cells were characterized by use of in vitro nuclear run-on and northern blot hybridization assays. The cytotoxic activities of agonistic anti-Fas antibodies, Fas ligand, and allogeneic peripheral blood leukocytes, in the absence or presence of Fas-blocking monoclonal antibodies, against tumor cells were assessed by methylene blue staining and chromium-51 release assays. Clin. relevant concns. of cisplatin, doxorubicin, mitomycin C, fluorouracil, or camptothecin enhanced Fas receptor expression on the plasma membrane of HT29 cells. Cisplatin-mediated increases in Fas expression were confirmed in HCT8R, HCT116, and U937 cells. The

enhancement of Fas protein expression was associated with an increased sensitivity of cisplatin-treated tumor cells to agonistic anti-Fas antibodies, to soluble Fas ligand, and to allogeneic peripheral blood leukocyte-mediated cytotoxicity. Each of these effects was blocked by co-treatment of the cells with antagonistic anti-Fas antibody. In addition to their direct cytotoxic effects, chemotherapeutic drugs sensitize tumor cells to Fas-mediated cytotoxicity and Fas-dependent immune clearance. On the basis of these findings, new strategies might be developed to improve the efficacy of these drugs.

OS.CITING REF COUNT: 183 THERE ARE 183 CAPLUS RECORDS THAT CITE THIS RECORD (183 CITINGS)  
 REFERENCE COUNT: 44 THERE ARE 44 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 94 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1997:395476 CAPLUS

DOCUMENT NUMBER: 127:85994

ORIGINAL REFERENCE NO.: 127:16425a,16428a

TITLE: Cytotoxicity of 5-fluorouracil released from a

bioadhesive patch into uterine cervical tissue

AUTHOR(S): McCarron, P. A.; Woolfson, A. D.; McCafferty, D. F.;

Price, J. H.; Sidhu, H.; Hickey, G. I.

CORPORATE SOURCE: School Pharmacy, Medical Biol. Centre, Queen's Univ.

Belfast, Belfast, BT9 7BL, UK

SOURCE: International Journal of Pharmaceutics (1997

), 151(1), 69-74

CODEN: IJPHDE; ISSN: 0378-5173

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB HeLa cells were used as a model cell line to evaluate the cytotoxic concentration

of 5-fluorouracil as a candidate drug for the topical treatment

of cervical intraepithelial neoplasia (CIN). Cytotoxicity was measured by

exposing cell suspensions to increasing concns. of drug and measuring the

decreased rate of cell growth. Results were confirmed by photographing

monolayers and estimating the ratio of cells entering mitosis. A drug

concentration

of 10-4M was cytotoxic. Cervical tissue samples were exposed for either 4

or 24 h periods to 5-fluorouracil released from a bioadhesive cervical

patch containing 20 mg of drug. The concentration distribution of

5-fluorouracil

through cervical tissue were estimated from the ams., as determined by HPLC, extracted

from tissue slices harvested at depths down to 5 mm from the

surface. Even at this depth, the tissue concentration following a 24-h

exposure to 5-fluorouracil was 100-fold that of the determined cytotoxic drug

concentration, indicating that the patch delivery system could result in clin.

effective drug concns. in those areas of the cervical stroma where

pre-cancerous lesions characteristic of cervical intraepithelial neoplasia

can occur.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

REFERENCE COUNT: 12 THERE ARE 12 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 95 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1997:222846 CAPLUS

DOCUMENT NUMBER: 126:258646

ORIGINAL REFERENCE NO.: 126:49901a,49904a

TITLE: Doxorubicin sensitizes human bladder carcinoma

cells to Fas-mediated cytotoxicity  
 AUTHOR(S): Mizutani, Youichi; Okada, Yusaku; Yoshida, Osamu; Fukumoto, Manabu; Bonavida, Benjamin  
 CORPORATE SOURCE: Department of Urology, Faculty of Medicine, Kyoto University, Kyoto, 606, Japan  
 SOURCE: Cancer (New York) (1997), 79(6), 1180-1189  
 CODEN: CANCAR; ISSN: 0008-543X  
 PUBLISHER: Wiley-Liss  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB The resistance of bladder carcinoma to anticancer chemotherapeutic agents remains a major problem. Hence, several immunotherapeutic approaches have been developed to treat the drug-resistant cancer cells. Fas antigen (Fas) and Fas ligand participate in cytotoxicity mediated by T lymphocytes and natural killer cells. Like Fas ligand, anti-Fas monoclonal antibody (MoAb) induces apoptosis of the cells expressing Fas. This study examined whether bladder carcinoma cells are sensitive to cytotoxicity mediated by anti-Fas MoAb and whether anticancer agents synergize with anti-Fas MoAb in cytotoxicity. Cytotoxicity was determined by a 1-day microculture tetrazolium dye assay. Synergy was assessed by isobolog. anal. The T24 human bladder carcinoma cell line constitutively expressed the Fas on the cell surface; however, T24 line was resistant to anti-Fas MoAb. Treatment of T24 cells with anti-Fas MoAb in combination with mitomycin C, methotrexate, or 5-fluorouracil did not overcome their resistance to these agents. However, treatment of T24 cells with a combination of anti-Fas MoAb and doxorubicin resulted in a synergistic cytotoxic effect. In addition, the doxorubicin-resistant T24 cells were sensitive to treatment with a combination of anti-Fas MoAb and doxorubicin. Synergy was also achieved in three other bladder carcinoma cell lines and four freshly derived human bladder carcinoma cells. Treatment with anti-Fas MoAb in combination with epirubicin or pirarubicin also resulted in a synergistic cytotoxic effect on T24 cells. The mechanisms of synergy were examined. Anti-Fas MoAb did not affect the intracellular accumulation of doxorubicin, the expression of P-glycoprotein, or the expression of the antioxidant glutathione S-transferase- $\pi$  mRNA. However, treatment with doxorubicin enhanced the expression of Fas on T24 cells. This study demonstrated that treatment of bladder carcinoma cells with doxorubicin sensitized the cells to lysis by anti-Fas MoAb. The synergistic effect obtained with established doxorubicin-resistant bladder carcinoma cells and freshly isolated bladder carcinoma cells suggests that drug-resistant bladder carcinoma cells can be sensitized by doxorubicin to Fas- and Fas ligand-mediated cytotoxicity by lymphocytes. Furthermore, the sensitization required low concns. of doxorubicin, thus supporting the in vivo application of a combination of chemotherapy and immunotherapy in the treatment of drug-resistant and/or immunotherapy-resistant bladder carcinoma.  
 OS.CITING REF COUNT: 72 THERE ARE 72 CAPLUS RECORDS THAT CITE THIS RECORD (72 CITINGS)

L21 ANSWER 96 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1997:222842 CAPLUS  
 DOCUMENT NUMBER: 126:233263  
 ORIGINAL REFERENCE NO.: 126:44969a, 44972a  
 TITLE: Bimonthly high dose leucovorin and 5-fluorouracil 48-hour infusion with interferon-alpha-2a in patients with advanced colorectal carcinoma  
 AUTHOR(S): Tournigand, Christophe; Louvet, Christophe; De Gramont, Aimery; Lucchi, Elisabeth; Seitz, Jean-Francois; Mal, Frederic; Raymond, Eric; Cady, Jean; Carola, Elisabeth; et al.



CORPORATE SOURCE: Hopital Saint-Antoine, Paris, Fr.  
SOURCE: Cancer (New York) (1997), 79(6), 1094-1099  
CODEN: CANCAR; ISSN: 0008-543X

PUBLISHER: Wiley-Liss

DOCUMENT TYPE: Journal

LANGUAGE: English

AB The rationale for the modulation of 5-fluorouracil (5-FU) with interferon-alpha (IFN) is inhibition of 5-FU catabolism and 5-FU resistance. Clin. trials have shown debatable results when IFN is given in high doses with 5-FU used as a bolus alone or in combination with leucovorin (LV). A first-line Phase II study was performed in 50 patients with metastatic colorectal carcinoma who were given a bimonthly combination of high dose LV, a high dose 48-h infusion of 5-FU, and a low dose of IFN. The regimen was comprised of a 2-h infusion of LV, 500 mg/m<sup>2</sup>, on each of 2 consecutive days, and a 48-h infusion of 5-FU, 1.5 to 2 g/m<sup>2</sup>/24 h, starting after Day 1 of LV treatment every 2 wk until there was evidence of disease progression. IFN was administered s.c. three times weekly at a dose of 3 MU (body surface area [BSA] < 1.75 m<sup>2</sup>) or 4.5 MU (BSA ≥ 1.75 m<sup>2</sup>). World Health Organization toxicity Grade 3-4 occurred in 21 patients (42%): diarrhea in 6%, mucositis in 12%, neutropenia in 30%, and alopecia in 8%. The overall response rate was 44%; 1 patient had a complete response (2%), 21 had partial responses (42%), 23 had stable disease (46%), and 5 had disease progression (10%). The median progression free survival was 9 mo, and median survival was 25 mo. Bimonthly high dose LV, a high dose 48-h infusion of 5-FU, and a low dose of IFN had good activity in patients with advanced colorectal carcinoma. However, as in other schedules of LV and 5-FU, IFN induces high grade toxicity.

L21 ANSWER 97 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1996:502370 CAPLUS

DOCUMENT NUMBER: 125:132013

ORIGINAL REFERENCE NO.: 125:24421a,24424a

TITLE: Regional chemotherapy for inoperable pancreatic carcinoma

AUTHOR(S): Muchmore, James H.; Preslan, Janet E.; George, William J.

CORPORATE SOURCE: School Medicine, Tulane University, New Orleans, LA, 70112, USA

SOURCE: Cancer (New York) (1996), 78(3, Suppl.), 664-673

CODEN: CANCAR; ISSN: 0008-543X

PUBLISHER: Wiley-Liss

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Survival for adenocarcinoma of the pancreatic remains unchanged over the last two decades. The majority of patients (85%) are diagnosed with an inoperable tumor. Patterns of failure reveal that pancreatic cancer involves three compartments: the pancreatic bed and regional lymph nodes, the liver and the peritoneal surfaces. Twelve patients with advanced, unresectable pancreatic cancer, Stage II/III, were treated with regional intra-arterial chemotherapy and extra-corporeal hemofiltration directed towards the pancreatic tumor-bearing area and the liver. Five patients had an arterial catheter/port system placed within the celiac axis; the rest had an angio. placed arterial catheter. All patients had a 16 Fr PFM filtration catheter inserted in the vena cava positioning the tip at the level of the diaphragm and then connected to a hemofiltration unit. Mitomycin C was infused over 25 min followed by 5-FU over 10 min. The hemofiltration was begun before the drug infusion and continued for 70 min. The twelve patients underwent 33 cycles of regional chemotherapy plus hemofiltration. Five patients had a partial response

(45.5%), five had stable disease (45.5%), and one had progression (9%). Four patients were re-explored with one patient undergoing a curative resection. The average survival for patients with unresectable pancreatic adenocarcinoma is 13 mo. Tumor implantation and progression on the peritoneal surfaces remains the major site of treatment failure. Regional chemotherapy plus hemofiltration with MMC and 5-FU appears to improve the response of Stage II/III inoperable pancreatic cancer and can convert some patients to resectability without significant complications and with no mortality.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD (3 CITINGS)

L21 ANSWER 98 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1995:697820 CAPLUS

DOCUMENT NUMBER: 123:104894

ORIGINAL REFERENCE NO.: 123:18523a,18526a

TITLE: Use of the fluorescence-activated cell sorter (FACS) for in vitro assays of developmental toxicity

AUTHOR(S): Hooghe, R. J.; Ooms, D.

CORPORATE SOURCE: Environment Div., Flemish Inst. Technological Res., Mol, B-2400, Belg.

SOURCE: Toxicology in Vitro (1995), 9(3), 349-54

CODEN: TIVIEQ; ISSN: 0887-2333

PUBLISHER: Elsevier

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Our objective is to predict embryotoxicity with reliable in vitro techniques. In several exptl. systems, differentiation is accompanied by changes in the glycosylation pattern of cell-surface glycoconjugates. This is also the case with embryonal carcinoma cells. We have monitored the expression of receptors for wheat germ agglutinin (WGA). Murine embryonal carcinoma cells (P19 and F9) were exposed in vitro to xenobiotics for 1-3 days, then incubated successively with WGA-biotin (15 µg/mL) and streptavidin-phycoerythrin (SA-PE) (20 µg/mL), each for 30 min at room temperature. Cell-surface fluorescence was then analyzed using a fluorescence-activated cell sorter (FACS). Exposure to 1 µM retinoic acid, a known inducer of differentiation, altered glycosylation as indicated by changes in WGA binding. Clear-cut effects were also observed after exposure to salts of arsenic (20 µM), or nickel (50 µM), and to methotrexate (1 µg/mL), fluorouracil (1.3 µg/mL) or actinomycin D (0.04 µg/mL). These compds. affected the percentage of pos. cells, the intensity of labeling, or both. Two non-teratogenic compds. (metronidazole and sulfonilamide) have also been tested and had no effect. Lectin histochem. of embryonal carcinoma cells exposed to potentially toxic agents holds promise as a method for predicting embryotoxicity. FACS anal. allows rapid quantification.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L21 ANSWER 99 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1995:625445 CAPLUS

DOCUMENT NUMBER: 123:47488

ORIGINAL REFERENCE NO.: 123:8283a,8286a

TITLE: Low doses of anticancer drugs increase susceptibility

of tumor cells to lysis by autologous killer cells  
Matsuoka, Hiroaki; Eura, Masao; Chikamatsu, Kazuaki;  
Nakano, Koji; Kanzaki, Yuichi; Masuyama, Keisuke;  
Ishikawa, Takeru

CORPORATE SOURCE: Department Otolaryngology, Kumamoto University School  
Medicine, Kumamoto, 860, Japan

SOURCE: Anticancer Research (1995), 15(1), 87-92  
 CODEN: ANTRD4; ISSN: 0250-7005  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Pretreatment of squamous cell carcinoma (SCC) cells from four patients with low doses of cisplatin, carboplatin or 5-fluorouracil increased the susceptibility to lysis by autologous killer cells in vitro. Exposure of two SCC cell lines to low doses of these drugs increased the cell surface expression of both HLA class I and intercellular adhesion mol.-1 (ICAM-1). HLA class II, neural cell adhesion mol., and B7 were not expressed on the cell surface before or after such treatment. The results suggest that these drugs increase the susceptibility of tumor cells to autologous cell-mediated cytotoxicity, at least in part, by enhancing the expression of HLA class I and ICAM-1.  
 OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L21 ANSWER 100 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1995:534226 CAPLUS  
 DOCUMENT NUMBER: 122:281603  
 ORIGINAL REFERENCE NO.: 122:51063a,51066a  
 TITLE: Differential effects of recombinant interferon-alpha and 5-fluorouracil against colon cancer cells or against peripheral blood mononuclear cells  
 AUTHOR(S): Filippi, Rosaria De; Prete, Salvatore P.; Giuliani, Anna; Silvi, Enrico; Yamaue, Hiroki; Nieroda, Carol A.; Greiner, John W.; Vecchis, Liana De; Bonmassar, Enzo  
 CORPORATE SOURCE: Laboratory Tumor Immunology and Biology, National Cancer Institute, Bethesda, MD, USA  
 SOURCE: Anticancer Research (1994), 14(5A), 1767-73  
 CODEN: ANTRD4; ISSN: 0250-7005  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB Comparative studies on the suppressive effects of recombinant interferon-alpha (IFN- $\alpha$ ), 5-fluorouracil (5-FU), or IFN- $\alpha$  + 5-FU have been performed in vitro on colon carcinoma cells (HT-29 cell line) and PHA-stimulated mononuclear cells (MNC) of peripheral blood obtained from healthy donors. IFN- $\alpha$  was used at 500 U/mL against HT-29 cells and at 1000 U/mL against MNC on day 1 of culture; 5-FU was used at 250  $\mu$ M against HT-29 and at 1400  $\mu$ M against MNC on day 2 of culture. The results show that: (a) IFN- $\alpha$  inhibited MNC and HT-29 cells by 13.4% and 32.9%, resp.; (b) 5-FU inhibited MNC and HT-29 cells by 54.7% and 87.0%, resp.; (c) IFN- $\alpha$  + 5-FU resulted in a stronger inhibition of HT-29 cells (i.e., 96.1%). In contrast, that combination was significantly less suppressive than 5-FU alone when MNC were used as targets (i.e., 35.9% inhibition). Natural cell-mediated cytotoxic activity relative to 106 MNC was not markedly altered by all agents alone or in combination. Moreover, treatment with IFN- $\alpha$ , 5-FU or IFN- $\alpha$  + 5-FU resulted in a marked increase in the number of HT-29 cells pos. for the CEA surface antigen. These data seem to provide further rational support of the clin. use of IFN- $\alpha$  + 5-FU in colorectal cancer, based on the differential toxicity of this drug combination on tumor vs. normal immunocompetent cells.

L21 ANSWER 101 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1995:524394 CAPLUS  
 DOCUMENT NUMBER: 122:256405  
 ORIGINAL REFERENCE NO.: 122:46537a,46540a  
 TITLE: Prevention and control of cancer with antiinflammatory agents and hyaluronic acid

INVENTOR(S): Falk, Rudolf E.; Asculai, Samuel S.  
PATENT ASSIGNEE(S): Norpharmco Inc., Can.  
SOURCE: Can. Pat. Appl., 213 pp.

CODEN: CPXXEB  
DOCUMENT TYPE: Patent  
LANGUAGE: English  
FAMILY ACC. NUM. COUNT: 1  
PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2097892	A1	19941207	CA 1993-2097892	19930607 <--
PRIORITY APPLN. INFO.:			CA 1993-2097892	19930607

AB A method of conditioning the immune system in humans to resist the formation of  $\geq 1$  cancerous tissue types comprises administering a nontoxic dosage amount of a composition comprising pharmaceutical excipients, a nonsteroidal antiinflammatory agent, hyaluronic acid and/or salts or derivs. thereof, and optionally vitamin C. Thus, repeated topical application of a 2.5% Na hyaluronate gel containing 3% Na diclofenac to basal cell carcinomas of the skin resulted in regression and disappearance of the lesions.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD  
(2 CITINGS)

L21 ANSWER 102 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1995:316085 CAPLUS

DOCUMENT NUMBER: 122:89434

ORIGINAL REFERENCE NO.: 122:16771a,16774a

TITLE: Formulations containing hyaluronic acid for facilitation of drug transport

INVENTOR(S): Falk, Rudolf E.; Asculai, Samuel S.; Klein, Ehud S.; Harper, David W.; Hochman, David; Furschke, Don

PATENT ASSIGNEE(S): Norpharmco Inc., Can.

SOURCE: Can. Pat. Appl., 117 pp.

CODEN: CPXXEB

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2089621	A1	19940817	CA 1993-2089621	19930216 <--
PRIORITY APPLN. INFO.:			CA 1993-2089621	19930216

AB Pharmaceutical compns. are provided from which effective nontoxic (to the patient) dosage amts. may be taken and applied to the skin and/or exposed tissue of a human, each effective dosage amount comprising pharmaceutical excipients suitable for topical application, an effective nontoxic dosage amount of a drug to treat a disease and/or condition of the skin and/or exposed tissue, and an effective nontoxic dosage amount of hyaluronic acid or its salts, homologs, analogs, derivs., complexes, esters, fragments, and/or subunits sufficient to facilitate or cause transport of the drug to a site in the skin, including epidermis or exposed tissue, resulting in its accumulation for a prolonged period of time. Thus, a gel containing glycerin 150, PhCH2OH 90, diclofenac Na 90, Na hyaluronate 75 g, and water 2795 mL, applied topically to cutaneous basal cell carcinoma several times a day for several wk, caused disappearance of the carcinoma.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD  
(3 CITINGS)

L21 ANSWER 103 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1995:309098 CAPLUS  
 DOCUMENT NUMBER: 122:64428  
 ORIGINAL REFERENCE NO.: 122:12191a,12194a  
 TITLE: Treatment of disease employing hyaluronic acid to facilitate transport of nonsteroidal antiinflammatory drugs (NSAIDs)  
 INVENTOR(S): Falk, Rudolf E.; Asculai, Samuel S.  
 PATENT ASSIGNEE(S): Norpharmco Inc., Can.  
 SOURCE: Can. Pat. Appl., 116 pp.  
 CODEN: CPXXEB  
 DOCUMENT TYPE: Patent  
 LANGUAGE: English  
 FAMILY ACC. NUM. COUNT: 1  
 PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
CA 2089635	A1	19940817	CA 1993-2089635	19930216 <--
PRIORITY APPLN. INFO.:			CA 1993-2089635	19930216

AB A pharmaceutical composition comprises a plurality of effective nontoxic dosage amts. of a NSAID for topical administration to the site of pathol. and/or trauma of skin and/or exposed tissue of a human patient, combined with an effective nontoxic dosage amount of hyaluronic acid and/or its salts, homologs, analogs, derivs., complexes, esters, fragments, and/or subunits to facilitate or cause transport of the drug to the site of the pathol. and/or trauma. Thus, application of a formulation containing glycerin 150, PhCH2OH 90, diclofenac Na 90, Na hyaluronate 75 g, and water 2795 mL to an actinic keratosis lesion 3 times daily for 7 days resulted in complete resolution of the lesion.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L21 ANSWER 104 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1994:289565 CAPLUS  
 DOCUMENT NUMBER: 120:289565  
 ORIGINAL REFERENCE NO.: 120:50751a,50754a  
 TITLE: The combined action of ICI-D1694, 5-fluoro-2'-deoxyuridine and 5-fluorouracil in inhibiting the growth of a human renal cell carcinoma cell line (RPMI-SE) in vitro  
 AUTHOR(S): Guimaraes, Manoel A.; Greco, William R.; Slocum, Harry K.; Huben, Robert P.; Rustum, Youcef M.  
 CORPORATE SOURCE: Dep. Urol. Oncol., Roswell Park Cancer Inst., Buffalo, NY, 14263, USA  
 SOURCE: International Journal of Oncology (1994), 4(1), 137-41  
 CODEN: IJONES; ISSN: 1019-6439  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 AB In order to investigate possible interactions among ICI-D1694 (a new folate-analog thymidylate synthase inhibitor), 5-fluoro-2'-deoxyuridine (FdUrd) and 5-fluorouracil (FUra), the effect of these agents alone and in 2-drug combinations against a human renal cell carcinoma cell line (RPMI-SE) in vitro was investigated. The median IC50's for cell growth inhibition for ICI-D1694, FdUrd and FUra were 4.00, 7.23 and 1,340 nM, resp. To quant. assess the degree of agent-combined action for 2-agent combinations of the 3 drugs, data from combination expts. were fitted with a response surface math. model (Greco et al, Cancer Res 50: 5318-5327, 1990). In 3 expts. for each combination, moderate Loewe synergism was consistently shown for ICI-D1694/FdUrd, less prominent

Loewe synergism was indicated for FdUrd/FUra; and Loewe additivity was shown for ICI-D1694/FUra. Studies in vitro to elucidate the mechanism of the interaction of ICI-D1694 + FdUrd, and in vivo to establish possible therapeutic advantages are warranted.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L21 ANSWER 105 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1993:462508 CAPLUS

DOCUMENT NUMBER: 119:62508

ORIGINAL REFERENCE NO.: 119:11029a,11032a

TITLE: Characterization of a human ovarian carcinoma cell line: UCI 101

AUTHOR(S): Fuchtnier, C.; Emma, D. A.; Manetta, A.; Gamboa, G.; Bernstein, R.; Liao, S. Y.

CORPORATE SOURCE: Dep. Obstet. Gynecol., Univ. California, Irvine, CA, 92668, USA

SOURCE: Gynecologic Oncology (1993), 48(2), 203-9

CODEN: GYNOA3; ISSN: 0090-8258

DOCUMENT TYPE: Journal

LANGUAGE: English

AB A new epithelial ovarian carcinoma cell line (UCI 101) was established from the ascitic fluid and solid tumor of a patient with progressive papillary ovarian adenocarcinoma refractory to combination chemotherapy with cyclophosphamide, adriamycin, and cisplatin as well as single-agent chemotherapy with taxol and high-dose cisplatin. The cell line grew well with an in vitro doubling time of 24 h. The cell line expressed the B 72.3 (Tag 72), CA125, MH99 (ESA), and E29 (EMA) cell surface antigens and AE1/AE3 cytokeratins. It overexpressed glycoprotein P and the epidermal growth factor receptor. The in vitro responses to single antitumor agents including adriamycin, cisplatin, dequalinium chloride, etoposide, 5-fluorouracil, taxol, and tumor necrosis factor was examined. I.p. transplantation of the cells into athymic mice resulted in foci of tumor on all peritoneal surfaces including the viscera and diaphragm, ultimately leading to solid bulky disease with a massive production of ascites. High levels of CA125 (>500 units/mL) were detected in the blood serum of tumor-bearing mice. Cytogenetic anal. of cultured cells showed several marker chromosomes containing deletions, duplications, and translocations. Cytol. and histol. evaluation of the xenograft revealed morphol. characteristics identical to those of the original tumor.

OS.CITING REF COUNT: 11 THERE ARE 11 CAPLUS RECORDS THAT CITE THIS RECORD (11 CITINGS)

L21 ANSWER 106 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1993:261030 CAPLUS

DOCUMENT NUMBER: 118:261030

ORIGINAL REFERENCE NO.: 118:45255a,45258a

TITLE: Pharmaceutical composition and method for treatment of premalignant and malignant lesions

INVENTOR(S): Klein, Edmund

PATENT ASSIGNEE(S): Cancer Immunology Research Corp., USA

SOURCE: PCT Int. Appl., 31 pp.

CODEN: PIXXD2

DOCUMENT TYPE: Patent

LANGUAGE: English

FAMILY ACC. NUM. COUNT: 1

PATENT INFORMATION:

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
-----	----	-----	-----	-----

WO 9306860 A1 19930415 WO 1992-US8349 19921007 <--  
W: AT, AU, BB, BG, BR, CA, CH, CS, DE, DK, ES, FI, GB, HU, JP, KP,  
KR, LK, LU, MG, MN, MW, NL, NO, PL, RO, RU, SD, SE, US  
RW: AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, SE, BF,  
BJ, CF, CG, CI, CM, GA, GN, ML, MR, SN, TD, TG  
AU 9227768 A 19930503 AU 1992-27768 19921007 <--  
PRIORITY APPLN. INFO.: US 1991-772413 A2 19911007  
WO 1992-US8349 A 19921007  
AB A pharmaceutical composition and method for control and treatment of lesions including tumors of the skin or deep-seated origin, e.g. liver, is disclosed. The method comprises administering to a patient  $\geq 2$  sensitizing agents in ams. sufficient to induce a delayed hypersensitivity response and applying or administering to the patient a min. amount of an immunolog. preparation sufficient to induce cell-mediated challenge response. The immunolog. preparation comprises  $\geq 2$  sensitizing agents in a pharmaceutically acceptable carrier. A 62 yr old female diagnosed to have mycosis fungoides was placed on topical immunotherapy with dinitrochlorobenzene and 5-fluorouracil and subsequent application of topical nitrogen mustard. The skin lesions regressed or disappeared within 3 mo of the therapy and 6 yr later she was free of the disease.  
REFERENCE COUNT: 3 THERE ARE 3 CITED REFERENCES AVAILABLE FOR THIS RECORD. ALL CITATIONS AVAILABLE IN THE RE FORMAT

L21 ANSWER 107 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1992:645002 CAPLUS  
DOCUMENT NUMBER: 117:245002  
ORIGINAL REFERENCE NO.: 117:42171a, 42174a  
TITLE: Relationships between the chromatographic retention data and the effects of nucleoside derivatives in highly metastatic 3LL cells  
AUTHOR(S): Pogany, G.; Cserhati, T.; Olah, J.; Valko, K.  
CORPORATE SOURCE: Jt. Res. Organ., Hung. Acad. Sci., Budapest, H-1086, Hung.  
SOURCE: Journal of Pharmaceutical and Biomedical Analysis (1992), 10(7), 495-500  
CODEN: JPBADA; ISSN: 0731-7085  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The effect of 21 nucleoside derivs. on the [3H]-thymidine cellular uptake and on the incorporation into DNA of highly metastatic 3LL (Lewis lung carcinoma) cells has been measured. Hydrophobic and hydrophilic mol. parameters (the adsorption capacity, specific adsorption surface, lipophilicity and specific hydrophobic surface area) have been determined by using TLC. Stepwise linear regression anal. and principal component anal. have been applied in order to reveal the relationships between the mol. parameters and the effect of the nucleoside derivs. on highly metastatic 3LL cells. The first principal component obtained from the measured activity data could be attributed to the change of [3H]-thymidine cellular uptake caused by the nucleoside, while the second principal component could be regarded as the measure of the effect on the DNA incorporation of [3H]-thymidine. The effect of nucleosides on the [3H]-thymidine uptake could be explained by the specific hydrophobic and adsorption surface area of the nucleoside, on the other hand the effect on the DNA incorporation could be described by the adsorption characteristics (specific hydrophilic surface area and adsorption capacity) of the derivs.

L21 ANSWER 108 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1992:34074 CAPLUS  
DOCUMENT NUMBER: 116:34074

ORIGINAL REFERENCE NO.: 116:5628h,5629a  
 TITLE: Adaptation to 5-fluorouracil of the heterogeneous human colon tumor cell line HT-29 results in the selection of cells committed to differentiation

AUTHOR(S): Lesuffleur, Thecla; Kornowski, Anne; Luccioni, Catherine; Muleris, Martine; Barbat, Alain; Beaumatin, Jacqueline; Dussaulx, Elisabeth; Dutrillaux, Bernard; Zweibaum, Alain

CORPORATE SOURCE: Unite Diff. Cell. Intest., Villejuif, 94807, Fr.  
 SOURCE: International Journal of Cancer (1991), 49(5), 721-30  
 CODEN: IJCNAW; ISSN: 0020-7136

DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB The HT-29 cell line contains a small proportion of differentiated, polarized, enterocytic and mucus-secreting cell types which can be selected under various conditions, e.g., glucose deprivation or methotrexate. The purpose of the present work was to investigate whether this also applied to 5-fluorouracil (FUra). Stepwise adaptation of exponentially growing cells to 1, 5, 10 and 20  $\mu$ M FUra resulted, after a phase of high mortality, in the emergence of adapted subpopulations with stable growth rates and curves, and IC50 values 6, 18, 37, and 110 times higher, resp., than in untreated cells. FUra-adapted cells were all differentiated, according to 2 phenotypes: (1) polarized dome-forming cells which express carcinoembryonic antigen at their apical surface and (2) goblet cells which secrete a mucus of colonic immunoreactivity. These phenotypes are present in the parental population and are different from those selected, e.g., by glucose deprivation or methotrexate. This differentiation pattern was maintained when the cells were subcultured in drug-free medium. Resistance to FUra is acquired through gene amplification, as substantiated by a 4-6-fold increase of thymidylate synthase gene copies in cells stably adapted to the drug. Whether the same mechanism or others are responsible for the 1st steps of resistance to FUra remains to be elucidated. These results support the hypothesis that some of the cells, which are present in the parental line and are committed to differentiation possess advantages which allow them to immediately resist and secondarily adapt to FUra. Comparison of the differentiation characteristics of FUra-adapted cells with those from cells selected under other pressure conditions suggests that resistance and adaptation to either type of pressure may depend on the differentiated phenotype to which the cells are committed.

OS.CITING REF COUNT: 38 THERE ARE 38 CAPLUS RECORDS THAT CITE THIS RECORD (38 CITINGS)

L21 ANSWER 109 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1991:622972 CAPLUS  
 DOCUMENT NUMBER: 115:222972  
 ORIGINAL REFERENCE NO.: 115:37771a,37774a  
 TITLE: Generation of 5-fluorouracil from 5-fluorocytosine by monoclonal antibody-cytosine deaminase conjugates

AUTHOR(S): Senter, Peter D.; Su, Peter C. D.; Katsuragi, Tohoru; Sakai, Takuo; Cosand, Wesley L.; Hellstrom, Ingegerd; Hellstrom, Karl Erik

CORPORATE SOURCE: Oncogen Div., Bristol-Myers Squibb Pharm. Res. Inst., Seattle, WA, 98121, USA

SOURCE: Bioconjugate Chemistry (1991), 2(6), 447-51  
 CODEN: BCCHEJ; ISSN: 1043-1802

DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB Cytosine deaminase (CDase) catalyzes the conversion of cytosine to uracil and is also able to convert the clin. used antifungal agent



5-fluorocytosine (5FC) into the anticancer drug 5-fluorouracil (5FU). The enzyme was purified from bakers' yeast CDase had a mol. weight of .apprx.32 kDa and was composed of 2 subunits of equal mol. wts. Monoclonal antibodies were covalently attached to CDase, forming conjugates that could bind to antigens on tumor cell surfaces. The combination of L6-CDase and 5FC was equivalent in cytotoxic activity to 5FU when tested against the H2981 human lung adenocarcinoma cell line (L6 pos., IF5 neg.). 5FC alone was noncytotoxic. The activation of 5FC was immunol. specific since IF5-CDase did not enhance 5FC activity.

OS.CITING REF COUNT: 44 THERE ARE 44 CAPLUS RECORDS THAT CITE THIS RECORD (44 CITINGS)

L21 ANSWER 110 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1991:549928 CAPLUS  
DOCUMENT NUMBER: 115:149928  
ORIGINAL REFERENCE NO.: 115:25439a,25442a  
TITLE: Effects of fluoropyrimidines on cellular proliferation and biological markers of a human salivary gland adenocarcinoma cell line  
AUTHOR(S): Kasai, Yasuo  
CORPORATE SOURCE: Sch. Dent., Univ. Tokushima, Tokushima, 770, Japan  
SOURCE: Shikoku Shigakkai Zasshi (1991), 4(1), 11-28  
CODEN: SSZAED; ISSN: 0914-6091  
DOCUMENT TYPE: Journal  
LANGUAGE: Japanese  
AB The fluoropyrimidine 5-fluoro-2'-deoxyuridine-5'-monophosphate (I) suppressed the growth and colony formation of HSG, a human salivary gland adenocarcinoma cell line. I induced differentiation of HSG into cells with phenotypes of myoepithelial cells and possessed antitumor activity in HSG-grafted nude mice. Growth inhibition was observed from 25 to 50 µg/mL of Tegafur (II), 1 to 5 µg/mL of I, and 0.1 to 0.5 µg/mL of 5-fluorouracil (III), 5'-fluoro-2'-deoxyuridine (IV), and 5-fluorouridine (V). II suppressed colony formation in soft agar and on plastic surfaces at concns. of 25 to 50 µg/mL. V suppressed this growth at 0.1 µg/mL. III, IV, and I suppressed the colony formation dose-dependently at concns. of more than 0.1 µg/mL, 0.1 µg/mL and 1 µg/mL, resp. HSG expressed myosin and S-100 protein β-chain when treated with 2 µg/mL of I. An ultrastructural study I-treated HSG cells showed myoepithelial cell-like structure. I suppressed the HSG growth in nude mice and elongation of the survival periods.

L21 ANSWER 111 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1990:434459 CAPLUS  
DOCUMENT NUMBER: 113:34459  
ORIGINAL REFERENCE NO.: 113:5725a,5728a  
TITLE: Characterization of adriamycin-resistant human breast cancer cells which display overexpression of a novel resistance-related membrane protein  
AUTHOR(S): Chen, Yi Nan; Mickley, Lyn A.; Schwartz, Arnold M.; Acton, Edward M.; Hwang, Jaulang; Fojo, Antonio T.  
CORPORATE SOURCE: Div. Cancer Treat., Natl. Cancer Inst., Bethesda, MD, 20892, USA  
SOURCE: Journal of Biological Chemistry (1990), 265(17), 10073-80  
CODEN: JBCHA3; ISSN: 0021-9258  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Development of multidrug resistance due to overexpression of P-glycoprotein (Pgp), a cell membrane drug efflux pump, occurs commonly during in vitro selections with adriamycin (Adr). Pgp-mediated drug

resistance can be overcome by the Ca<sup>2+</sup> channel blocker verapamil (Vp), which acts as a competitive inhibitor of drug binding and efflux. In order to identify other mechanisms of Adr resistance, an Adr-resistant subline was isolated by selecting the human breast cancer cell line MCF-7 with incremental increases of Adr in the presence of 10 µg Vp/mL. The resultant MCF-7/Adr Vp subline is 900-fold more resistant to Adr, does not overexpress Pgp, and does not exhibit a decrease in Adr accumulation. It exhibits a unique cross-resistance pattern: high cross-resistance to the potent Adr analog 3'-deamino-3'-(3-cyano-4-morpholinyl)doxorubicin, lower cross-resistance to the alkylating agent melphalan, and a sensitivity to vinblastine similar to that of the parental cell line. The levels of glutathione and glutathione S-transferase are similar in the parental line and the Adr-resistant subline. Topoisomerase II-DNA complexes measured by the K-Na dodecyl sulfate precipitation method showed a 2-3-fold decrease in the resistant subline. The MCF-7/Adr Vp cells overexpress a novel membrane protein with an apparent mol. mass of 95 kilodaltons. Polyclonal antibodies raised against the P-95 protein demonstrate a correlation between the level of expression and Adr resistance. Removal of Adr but not Vp from the selection media results in a decline in P-95 protein levels that parallels a restoration of sensitivity to Adr. Immunohistochem. demonstrates localization of the P-95 protein on the cell surface. The demonstration of high levels of the protein in clin. samples obtained from patients refractory to Adr suggests that this protein may play a role in clin. drug resistance.

OS.CITING REF COUNT: 107 THERE ARE 107 CAPLUS RECORDS THAT CITE THIS RECORD (107 CITINGS)

L21 ANSWER 112 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1990:191553 CAPLUS  
 DOCUMENT NUMBER: 112:191553  
 ORIGINAL REFERENCE NO.: 112:32185a,32188a  
 TITLE: Intraperitoneal tumor growth and chemotherapy in a rat model  
 AUTHOR(S): Los, Gerrit; Ruevekamp, Marjan; Bosnie, Nel; De Graaf, Peter W.; McVie, J. Gordon  
 CORPORATE SOURCE: Dep. Exp. Ther., Neth. Cancer Inst., Amsterdam, 1066 CX, Neth.  
 SOURCE: European Journal of Cancer & Clinical Oncology (1989), 25(12), 1857-66  
 CODEN: EJCODS; ISSN: 0277-5379  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB A new animal model is described, in which the effects of i.p. administration of cytostatic drugs on cancers restricted to the peritoneal cavity can be studied. The tumor cell line used is a chemical induced carcinoma (CC531), sensitive in vitro to cisplatin (cDDP), carboplatin, 5-fluorouracil, doxorubicin and mitoxantrone. Three to 5 wk after i.p. inoculation of 2 + 106 ML531 cells, 80% of Wag/Rij rats develop small tumor nodules on peritoneal surfaces. Both tumor size and localization at this time are comparable to the human situation, especially to cases of min. residual disease ovarian carcinoma. The model was used to determine the usefulness of i.p. treatment in comparison to i.v. Changing the route of administration of cDDP from i.v. to i.p. increases tumor Pt concns. and prolonged survival. The model offers the possibility to study drug pharmacokinetics and tumor drug penetration related to i.p. drug administration.

OS.CITING REF COUNT: 6 THERE ARE 6 CAPLUS RECORDS THAT CITE THIS RECORD (6 CITINGS)

L21 ANSWER 113 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1988:622069 CAPLUS

DOCUMENT NUMBER: 109:222069  
ORIGINAL REFERENCE NO.: 109:36557a, 36560a  
TITLE: Modulation of the cytotoxic effect of 5-fluorouracil by N-methylformamide on a human colon carcinoma cell line  
AUTHOR(S): Zupi, Gabriella; Marangolo, Maurizio; Arancia, Giuseppe; Greco, Claudia; Laudonio, Nina; Iosi, Francesca; Formisano, Giuseppe; Malorni, Walter  
CORPORATE SOURCE: Lab. Exp. Chemother., Regina Elena Inst. Cancer Res., Rome, 00161, Italy  
SOURCE: Cancer Research (1988), 48(21), 6193-200  
CODEN: CNREA8; ISSN: 0008-5472  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB The cytotoxic effect of the combination of N-methylformamide (NMF) with 5-fluorouracil (5-FU) on survival of the human colon cancer line HT29 was assessed. The differentiating activity of NMF was evidenced by morphol. maturation and conversion of cell culture characteristics to those consistent with a more benign phenotype. In combination expts., the noncytotoxic concentration of 1% NMF was chosen and concns. of 5-FU ranging 5-25 µg/mL were employed. Two main schedules were tested either on exponentially or stationarily growing cells: (a) NMF for 72 h followed by 12-h exposure to 5-FU; (b) 5-FU for 12 h followed by 72-h exposure to NMF. The 5-FU + NMF sequence reduced the surviving fraction of HT29 cells, while the reverse sequence did not increase the killing effect of 5-FU alone. Immunocytochem. and electron-microscopic studies seemed to confirm that the association in which the differentiating agent followed the 5-FU treatment strongly impaired cellular integrity and function and that cytoskeletal elements, particularly microfilaments, and surface structures could play an essential role in the mechanisms of cytotoxicity. Thus, the drug sequence is a critical factor for the optimal combination of 5-FU and NMF.  
OS.CITING REF COUNT: 9 THERE ARE 9 CAPLUS RECORDS THAT CITE THIS RECORD (9 CITINGS)  
L21 ANSWER 114 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1988:431425 CAPLUS  
DOCUMENT NUMBER: 109:31425  
ORIGINAL REFERENCE NO.: 109:5181a, 5184a  
TITLE: New chemotherapeutic drug sensitivity assay for colon carcinomas in monolayer culture  
AUTHOR(S): Schroy, Paul C., III; Cohen, Alfred; Winawer, Sidney J.; Friedman, Eileen A.  
CORPORATE SOURCE: Dep. Med., Memorial Sloan-Kettering Cancer Cent., New York, NY, 10021, USA  
SOURCE: Cancer Research (1988), 48(11), 3236-44  
CODEN: CNREA8; ISSN: 0008-5472  
DOCUMENT TYPE: Journal  
LANGUAGE: English  
AB Ten previously untreated human colon carcinomas were tested for chemotherapeutic drug sensitivity in primary monolayer culture. Colon carcinomas were partly digested to groups of epithelial cells which plated with a mean efficiency of 42% on a collagen I-bovine serum albumin substrate in serum-free medium, producing patches of tightly adherent epithelial cells. The cultured cells were judged epithelial by the presence of cytokeratins, an epithelial cell surface epitope, junctional complexes, and brush borders. Each carcinoma was plated in 40-60 Petri dishes (35 mm), yielding a mean of 28 colonies per dish (6832 cells). Drugs tested in duplicate plates were mitomycin C, cisplatin, streptozotocin, and 5-fluorouracil at 0.1, 1, 10, and 100


µg/mL, and at 0.1, 1, and 2+ the peak tolerated drug concentration in serum. At 24 h after plating, any nonadherent cells were removed, and the adherent tumor cells were continuously exposed to the drugs for 3 days. Each drug induced colony lysis in a dose-dependent manner in responsive tumors. Drug-resistant, cycling cells were identified by <sup>3</sup>H-thymidine incorporation in colonies which were not lysed by drug treatment. Each of the 10 carcinomas exhibited inherent resistance to ≥ 1 chemotherapy drug within the concentration ranges clin. achievable.

L21 ANSWER 115 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1986:95399 CAPLUS  
 DOCUMENT NUMBER: 104:95399  
 ORIGINAL REFERENCE NO.: 104:15021a,15024a  
 TITLE: Adhesive topical drug delivery system  
 AUTHOR(S): Nagai, Tsuneji  
 CORPORATE SOURCE: Fac. Pharm. Sci., Hoshi Univ., Tokyo, 142, Japan  
 SOURCE: Journal of Controlled Release (1985), 2,  
 121-34  
 CODEN: JCREEC; ISSN: 0168-3659  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English

AB A new topical dosage form containing hydroxypropyl cellulose (HPC) [9004-64-2] and Carbopol 934 [9007-16-3] swelled with body fluids sticking to the disease area with a good adhesiveness when placed on topical membranes. With preps. for carcinoma colli containing drugs such as bleomycin [11056-06-7] a high percentage of disappearance of cancerous foci was observed when they were placed on the portio vaginalis of humans. Although the oral mucosal dosage form for the absorption of insulin [9004-10-8] showed low bioavailability this was the 1st case showing that insulin could be absorbed through the oral mucosal membrane. The application of an adhesive tablet for aphthous stomatitis treatment is described. The tablet consists of 2 layers, 1 adhesive and the other supporting layer. The adhesive layer consists of HPC and Carbopol 934 containing triaminolone acetonide [76-25-5] and the supporting layer consists of lactose. Results from the clin. study showed that the average dose of triaminolone acetonide/day in the adhesive tablet was about one tenth of that obtained in the existing ointment. Side effects were not observed. An improvement in aphtha was observed after the tablet administration.

OS.CITING REF COUNT: 45 THERE ARE 45 CAPLUS RECORDS THAT CITE THIS RECORD (45 CITINGS)

L21 ANSWER 116 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
 ACCESSION NUMBER: 1985:172530 CAPLUS  
 DOCUMENT NUMBER: 102:172530  
 ORIGINAL REFERENCE NO.: 102:27061a,27064a  
 TITLE: Effect of implanted ethylene-vinyl alcohol copolymer matrixes containing 5-fluorouracil on Ehrlich ascites carcinoma  
 AUTHOR(S): Miyazaki, Shozo; Takeuchi, Shigem; Sugiyama, Mieko; Takada, Masahiko; Hosokawa, Masuo; Koga, Yutaka; Kobayashi, Hiroshi  
 CORPORATE SOURCE: Fac. Pharm. Sci., Higashi-Nippon-Gakuen Univ., Hokkaido, 061-02, Japan  
 SOURCE: Journal of Pharmacy and Pharmacology (1985), 37(1), 64-6  
 CODEN: JPPMAB; ISSN: 0022-3573  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



AB The antitumor activity of ethylene-vinyl alc. copolymer (EVA) [25067-34-9] matrices containing 5-fluorouracil (I) [51-21-8] was evaluated against Ehrlich ascites carcinoma in mice. A prolongation of the life-span of tumor-bearing mice following i.p. implantation of therapeutic matrices was noted. EVA matrices containing I may be effective in cancer chemotherapy. Matrices composed of EVA could be useful vehicles for implanted, inserted, or surface-applied delivery systems for anticancer agents.

OS.CITING REF COUNT: 1 THERE ARE 1 CAPLUS RECORDS THAT CITE THIS RECORD (1 CITINGS)

L21 ANSWER 117 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1984:563284 CAPLUS

DOCUMENT NUMBER: 101:163284

ORIGINAL REFERENCE NO.: 101:24539a,24542a

TITLE: Endoscopic intramural injection of antineoplastic emulsion

AUTHOR(S): Ohta, Hirotooshi; Takagi, Kunio; Noguchi, Yoshikazu; Ohashi, Ichiro; Takahashi, Tomoyuki; Watanabe, Susumu; Takekoshi, Takao; Ohashi, Kazuhiko; Kato, Yo  
Dep. Surg., Cancer Inst., Tokyo, 170, Japan

SOURCE: Gann (1984), 75(7), 641-9  
CODEN: GANNA2; ISSN: 0016-450X

DOCUMENT TYPE: Journal

LANGUAGE: English

AB With the aim of establishing a topical chemotherapy against stomach carcinoma, 5-fluorouracil (5-FU) [51-21-8] emulsion (oil/water type) for injection was developed. The drug distribution was analyzed by 5-FU bioassay and radiog. examination of soft parts, for which radiopaque Lipiodol was employed in an oil phase. In order to examine local toxicity, tissue retention, and transfer to lymph nodes of 5-FU emulsion, the drug was administered perorally to rats and injected intramurally through the gastric serosa into laparotomized dogs. Following this series of expts., which gave satisfactory results, the time courses of drug concentration in the gastric wall and regional lymph nodes were studied as a preclin. trial by giving endoscopic intramural injection of 5-FU emulsion or solution to dogs. The antimetastatic and antineoplastic effects of 5-FU emulsion were investigated in an exptl. model of lymph node metastasis in mice. The emulsion was more effective in subduing metastasis and tumor growth than the solution, and the effectiveness of the former was further augmented by the use of repeated injections rather than a single injection. This method of endoscopic injection of 5-FU emulsion should be of great value as a local therapeutic measure against stomach carcinoma itself as well as against metastatic lesions in the lymph nodes.


OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L21 ANSWER 118 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1983:59843 CAPLUS

DOCUMENT NUMBER: 98:59843

ORIGINAL REFERENCE NO.: 98:9109a,9112a  
 TITLE: Pharmaceutical application of biomedical polymers. Part VII. Antitumor effect of ethylene-vinyl acetate copolymer matrixes containing 5-fluorouracil on Ehrlich ascites carcinoma in mice  
 AUTHOR(S): Miyazaki, Shozo; Ishii, Kuniaki; Sugibayashi, Kenji; Morimoto, Yasunori; Takada, Masahiko  
 CORPORATE SOURCE: Fac. Pharm. Sci., Hagashi-Nippon-Gakuen Univ., Hokkaido, 061-20, Japan  
 SOURCE: Chemical & Pharmaceutical Bulletin (1982), 30(10), 3770-5  
 CODEN: CPBTAL; ISSN: 0009-2363  
 DOCUMENT TYPE: Journal  
 LANGUAGE: English  
 GI



AB ethylene-vinyl acetate copolymer (EVA) [24937-78-8] was evaluated as a carrier for controlled release of 5-fluorouracil (5-FU) (1) [51-21-8]. To study the effect of comonomer ratio modifications on the drug release kinetics, the release of 5-FU dispersed in polymer matrices composed of different ratios of ethylene and vinyl acetate was investigated. The vinyl acetate content of EVA was varied from 8 to 40% weight/weight. An increase in vinyl acetate comonomer content increased the drug release from the polymer matrix. The release rate could be controlled by modifying the ethylene/vinyl acetate ratios in the polymer matrices. The antitumor activity of EVA matrices containing 5-FU was evaluated against Ehrlich ascites carcinoma in mice on the basis of changes in body weight and animal survival data. Tumor cell injections were performed on Day 0 and matrix implantations of Day 4, both i.p. The suppressive effect of matrices containing 5-FU on the increase in body weight was higher than that of the free drug. A prolongation of the life-span of tumor-bearing mice following implantation of therapeutic matrices was also noted. Thus, EVA matrices containing 5-FU may be effective in cancer chemotherapy. Matrices composed of EVA could be useful vehicles for implanted, inserted, or surface-applied delivery systems for anticancer agents.

OS.CITING REF COUNT: 10 THERE ARE 10 CAPLUS RECORDS THAT CITE THIS RECORD (10 CITINGS)

L21 ANSWER 119 OF 124 CAPLUS COPYRIGHT 2010 ACS ON STN  
 ACCESSION NUMBER: 1982:484632 CAPLUS  
 DOCUMENT NUMBER: 97:84632  
 ORIGINAL REFERENCE NO.: 97:13881a,13884a  
 TITLE: Drug activity and therapeutic synergism in cancer treatment  
 AUTHOR(S): Carter, Walter H., Jr.; Wampler, Galen L.; Stablein, Donald M.; Campbell, Eleanor D.  
 CORPORATE SOURCE: Med. Coll. Virginia, Virginia Commonwealth Univ., Richmond, VA, 23298, USA  
 SOURCE: Cancer Research (1982), 42(8), 2963-71

CODEN: CNREA8; ISSN: 0008-5472

DOCUMENT TYPE:

Journal

LANGUAGE:

English

AB In work involving modeling of response surfaces to describe the effects of cancer chemotherapy treatments, it is important to define activity and therapeutic synergism in a statistically defendable manner. This requires the construction of confidence intervals around the estimated optimal treatment which has been achieved by use of an indirect method first proposed by Box and Hunter (1954). Activity for a drug or a combination can be claimed at  $100(1 - \alpha)\%$  level of confidence when the  $100(1 - \alpha)\%$  confidence interval about the optimal treatment excludes a zero dose. Results of treatment of B16 melanoma and Lewis lung carcinoma with 3,4-dihydroxybenzohydroxamic acid [69839-83-4] are used to demonstrate this definition. Extensions of this concept lead to a statistically valid definition of therapeutic synergism. If the confidence region about the optimum combination of  $k$  drugs does not contact any of the  $k - 1$  dimensional subspaces, then a  $k$  drug therapeutic synergism can be claimed. In the event that a  $k$  drug therapeutic synergism cannot be claimed, there may be subsets of the drugs which do combine with therapeutic synergy. These concepts are demonstrated by 2- and 3-drug combination expts. in L1210-bearing C57BL/6 + DBA/2 F1 (B6D2F1) mice. razoxane [21416-67-1] And dacarbazine [4342-03-4] show therapeutic synergism at a 95% confidence level. A 3-drug combination of 5-fluorouracil [51-21-8], Teniposide [29767-20-2], and mitomycin C [50-07-7] is considered. In this case, although the estimated optimum treatment includes 48.1 mg of 5-fluorouracil/kg, 15.9 mg of Teniposide/kg, and 3.9 mg of mitomycin C/kg, the confidence region generated failed to confirm at an 80% level of confidence that 5-fluorouracil was a necessary component of the best treatment.

OS.CITING REF COUNT: 3 THERE ARE 3 CAPLUS RECORDS THAT CITE THIS RECORD  
(3 CITINGS)

L21 ANSWER 120 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1982:135467 CAPLUS

DOCUMENT NUMBER: 96:135467

ORIGINAL REFERENCE NO.: 96:22069a,22072a

TITLE: Pharmacological interactions between 5-fluorouracil and methotrexate: new clinical applications  
Lingetti, M.; Belli, M.; Ciarimboli, M.; Guerriero, C.; Sorrentino, P.; Lingetti, E.

CORPORATE SOURCE: Div. Geriatr., Osp. Civile, Avellino, Italy  
SOURCE: Rassegna Internazionale di Clinica e Terapia (1981), 61(12), 843-52

CODEN: RICTA6; ISSN: 0370-548X



DOCUMENT TYPE:

Journal

LANGUAGE:

Italian

GI



AB Following a discussion of the antitumor mechanisms of 5-fluorouracil (I) [51-21-8] and methotrexate (II) [59-05-2] and of considerations indicating possible antagonism between them when given in the classical protocol of combination tumor therapy, a new cyclic dosage scheme is reported which was carried out on patients with metastasized mammary carcinoma. The treatment involved a 28-day cycle in which leucovorin [58-05-9] (3 mg) was given on the day preceding the initiation of antitumor therapy and on day 7; cyclophosphamide [50-18-0] (100 mg/m<sup>2</sup> body surface, orally) was given on days 1-14, I (600 mg/m<sup>2</sup>, orally) on days 2 and 9 and II (40 mg/m<sup>2</sup>, orally) on days 3 and 10. The percentage of complete remissions was 16%; overall, 69% of the patients showed some response. The mean duration of the remissions was 10 mo. These results were favorable in comparison with those of previous cyclophosphamide-I-II antitumor protocols.

L21 ANSWER 121 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1980:431739 CAPLUS

DOCUMENT NUMBER: 93:31739

ORIGINAL REFERENCE NO.: 93:5173a,5176a

TITLE: Pharmaceutical interactions in dosage forms and processing. Part XVII. Preparation and phase II clinical examination of topical dosage forms for the treatment of carcinoma colli containing bleomycin, carboquone, or 5-fluorouracil with hydroxypropyl cellulose

AUTHOR(S): Machida, Yoshiharu; Masuda, Hiroshi; Fujiyama, Norimasa; Iwata, Masanori; Nagai, Tsuneji

CORPORATE SOURCE: Hoshi Inst. Pharm. Sci., Tokyo, 142, Japan

SOURCE: Chemical & Pharmaceutical Bulletin (1980), 28(4), 1125-30

CODEN: CPBTAL; ISSN: 0009-2363

DOCUMENT TYPE: Journal

LANGUAGE: English

AB With the aim of developing a dosage form for the treatment of carcinoma colli, stick-like preps. containing bleomycin-HCl (I) [67763-87-5], carboquone (II) [24279-91-2], and 5-fluorouracil (III) [51-21-8] held in a mixture of hydroxypropyl cellulose (IV) [9004-64-2] and Carbopol 934 [9007-16-3] were prepared and clin. tested in volunteers suffering from carcinoma colli after various in vitro tests. The results of preliminary tests of drug release using the agar gel bed method indicated that the addition of Na lauryl sulfate enhanced the release of II, but the effect was not very great. Therefore, in order to enhance the release of II, the contents of IV in the base and II were increased. The preps. of I and III of 2 mm diameter showed faster drug release than those of 4 mm diameter according to the Kerami filter method.



In the preps. of 4 mm diameter, the release of II took place at almost the same rate as that of I, i.e., about 40% within 24 h, due to the modification of the formula for the preparation of II. In the case of the preparation of III, the release was so rapid that about 100% of the drug was released within 24 h. The present Kerami filter method seemed suitable and convenient for measuring the drug release from the present dosage forms. Clin. examination indicated the stick-like shape of the present dosage form to be favorable for the treatment of foci in the cervical canal. A high percentage of complete disappearance of the cancerous focus was obtained for patients of stage 0 in the cases of I and III, and a similar result was obtained for stage Ia in the case of II.

OS.CITING REF COUNT: 17 THERE ARE 17 CAPLUS RECORDS THAT CITE THIS RECORD (17 CITINGS)

L21 ANSWER 122 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1979:604361 CAPLUS

DOCUMENT NUMBER: 91:204361

ORIGINAL REFERENCE NO.: 91:32783a,32786a

TITLE: Clinical and pharmacological implications of cancer

cell differentiation and heterogeneity

AUTHOR(S): Calabresi, Paul; Dexter, Daniel L.; Heppner, Gloria H.

CORPORATE SOURCE: Dep. Med., Brown Univ., Providence, RI, 02912, USA

SOURCE: Biochemical Pharmacology (1979), 28(12), 1933-41

CODEN: BCPCA6; ISSN: 0006-2952

DOCUMENT TYPE: Journal

LANGUAGE: English

AB Starting 2 days after s.c. injection into mice of the tumor cell lines 68-H, 168, or 4.10 (each derived from the same single, autochthonous Balb/cf3H mouse mammary tumor), cyclophosphamide (I) [50-18-0] treatment (25-100 mg/kg, i.p., once a week for 4 wk) caused 67, 23, and 0% regression, resp. of these mammary tumor subpopulations in vivo, measured 3-4 mo after tumor cell injection. The corresponding values for in vivo regression by methotrexate (II) [59-05-2] (10-50 mg/kg) were 28, 4, and 0%, resp., and those for 5-fluorouracil (III) [51-21-8] (10-50 mg/kg) were 40, 0, and 0%, resp. When treatment (all 50 mg/kg) was started only after the appearance of palpable tumors, in vivo regression of 68-H, 168, and 4.10 subpopulations caused by I was 0, 24, and 32%, resp., by II was 43, 31, and 42%, resp., and by III was 17, 29, and 46%, resp. The molar concns. of III required to half the doublings of 68-H, 168, and 4.10 cells were  $3.2 \times 10^{-7}$ ,  $1.5 \times 10^{-8}$ , and  $4.3 \times 10^{-7}$ , resp., and those of II were  $7.0 \times 10^{-10}$ ,  $1.4 \times 10^{-10}$ , and  $2.9 \times 10^{-10}$ , resp. These subpopulations also showed marked variation in growth potential and surface antigens. N,N-Dimethylformamide [68-12-2] induced differentiation in rhabdomyosarcoma cells and in human colon carcinoma cells. The clin. implications of cancer cell differentiation and heterogeneity are discussed.

OS.CITING REF COUNT: 2 THERE ARE 2 CAPLUS RECORDS THAT CITE THIS RECORD (2 CITINGS)

L21 ANSWER 123 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN

ACCESSION NUMBER: 1977:577693 CAPLUS

DOCUMENT NUMBER: 87:177693

ORIGINAL REFERENCE NO.: 87:28014h,28015a

TITLE: Electrophoretic behavior and invasion of the drug

resistant sublines of Ehrlich ascites tumor cells

AUTHOR(S): Ku, Kuo-Yen; Liu, Li; Li, Mei-Fang; Hong, Long-Sun

CORPORATE SOURCE: Inst. Exp. Biol., Shanghai, Peop. Rep. China

SOURCE: Dongwu Xuebao (1977), 23(2), 207-11

CODEN: TWHPA3; ISSN: 0001-7302

DOCUMENT TYPE: Journal  
LANGUAGE: Chinese  
GI

\* STRUCTURE DIAGRAM TOO LARGE FOR DISPLAY - AVAILABLE VIA OFFLINE PRINT \*

AB Ehrlich ascites tumor cells became resistant to actinomycin D (I) [50-76-0], vinblastine (II) [865-21-4] and 5-fluorouracil (III) [51-21-8] after prolonged treatment in mice with these drugs. Electrophoretic mobility (as contributed by the total cell surface charges) of cells of II-resistant subline was significantly lower and those of I- and III-resistant sublines were significantly higher than that of sensitive one, indicating that there is no correlation between drug resistance and electrophoretic behavior. The mobility of cells of I-resistant subline remained unchanged after the interruption of the drug >1/2 years. On the other hand, in II- and III-resistant sublines, the mobility changed significantly after the discontinuation of drug treatment for 1 week or >1/2 year. The charge d. of PO<sub>4</sub>, NH<sub>2</sub> and SH groups on the cell surfaces changed in 1 way or the other. These indicate that electrophoretic behavior is determined by both the genetic and physiol. adaptations. The invasion of Ehrlich ascites tumor cells into fat tissues of the uterus in tumor bearing mice was not evident upon gross examination, but was noticed in histol. examns. in some cases. However, after the treatment of tumor cells with II for .apprx.40 weeks, the infiltrated fat tissue became enlarged, thickened and opalescent in appearance. This sort of invasion persisted for 40-50 generations and vanished gradually, despite the continuous presence of II. Invasion into the fat tissue was less apparent and prolonged in I- and III-resistant sublines. It seems that invasion of tumor cells is a rather complex phenomenon which may be related to both genetic and the physiol. adaptations, but in no way related to electrophoretic mobility.

L21 ANSWER 124 OF 124 CAPLUS COPYRIGHT 2010 ACS on STN  
ACCESSION NUMBER: 1967:74555 CAPLUS  
DOCUMENT NUMBER: 66:74555  
ORIGINAL REFERENCE NO.: 66:13959a,13962a  
TITLE: Prevention of skin cancer with topical 5-fluorouracil  
AUTHOR(S): Neldner, Kenneth H.  
CORPORATE SOURCE: Univ. of Colorado, Denver, CO, USA  
SOURCE: Rocky Mountain Medical Journal (1966), 63(11), 74-8  
CODEN: RMMJAK; ISSN: 0035-760X  
DOCUMENT TYPE: Journal  
LANGUAGE: English

AB An Aquaphor ointment containing 3% 5-fluorouracil (I) was effective in removing keratoses, particularly those caused by light and which often develop into cancer, but was ineffective against established carcinoma or psoriasis. I is thought to block intracellular synthesis of DNA and convert the RNA synthesis to fraudulent RNA. The rate of DNA synthesis in keratoses related to the rate in normal skin is a critical factor. This theory, however, does not account for the noneffectiveness of I in psoriasis nor the much lower effectiveness of other antimetabolites, particularly 6-mercaptopurine and methotrexate.

=> d his

(FILE 'HOME' ENTERED AT 13:27:42 ON 22 APR 2010)

FILE 'CAPLUS' ENTERED AT 13:29:20 ON 22 APR 2010  
S UBIQUINONE/CN

L1 FILE 'REGISTRY' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S UBIQUINONE/CN

L2 FILE 'CAPLUS' ENTERED AT 13:29:44 ON 22 APR 2010  
0 S L1  
S COENZYME Q10/CN

L3 FILE 'REGISTRY' ENTERED AT 13:30:10 ON 22 APR 2010  
1 S COENZYME Q10/CN

L4 FILE 'CAPLUS' ENTERED AT 13:30:10 ON 22 APR 2010  
5610 S L3

L5 FILE 'REGISTRY' ENTERED AT 13:30:20 ON 22 APR 2010  
1 S COENZYME Q10/CN

L6 FILE 'CAPLUS' ENTERED AT 13:30:34 ON 22 APR 2010  
41 S L5 AND CARCINOMA  
L7 16 S L6 AND PY<=2004  
L8 7 S L5 AND CARCINOMA AND TOPICAL  
L9 38 S L5 AND ?CARCINOMA  
L10 8 S L9 AND (TOPICAL OR SURFACE)  
L11 13 S L5 AND CANCER AND (TOPICAL OR SURFACE)  
L12 5 S L11 AND PY<=2004

L13 FILE 'REGISTRY' ENTERED AT 13:48:56 ON 22 APR 2010  
1 S FLUOROURACIL/CN  
L14 0 S L13 AND L5

L15 FILE 'CAPLUS' ENTERED AT 13:50:02 ON 22 APR 2010  
27 S L13 AND L5  
L16 4 S L15 AND CANCER  
L17 2 S L15 AND ?CARCINOMA  
L18 1153 S L13 AND (TOPICAL OR SURFACE)  
L19 388 S L18 AND CANCER  
L20 260 S L18 AND ?CARCINOMA  
L21 124 S L20 AND PY<=2004